

Optically Levitated Microspheres as a Probe for New Interactions

Alexander Rider, Charles Blakemore, Giorgio Gratta, Maxime
Louis, Marie Lu, David Moore, Sandip Roy


Stanford-Yale Collaboration

2017-07-24

Searching for New Interactions


- ▶ Particle accelerators discovered many interactions by producing and detecting their quanta.
- ▶ We are searching for the exchange of classical momentum transferred by an interaction. We are interested in short range ($\sim 1\mu m$) interactions coupled to mass usually parameterized by a Yukawa potential.

$$V(r) = -\frac{Gm_1m_2}{r} \left(1 + \alpha e^{-r/\lambda}\right)$$





Theory regions adapted from PRD **68**, 124021 (2003)

Optically Levitated Microspheres




- ▶ Difficult to measure momentum exchanges due to suspension.
- ▶ Laser radiation pressure suspension achieves excellent isolation.
- ▶ The optical force on the microsphere can be measured from the change in optical momentum flux.
- ▶ Measured force sensitivity of $\sim 5 \times 10^{-18} \text{ N}/\sqrt{\text{Hz}}$ with $5 \mu\text{m}$ spheres

Our Trap




- ▶ Trap must be isolated from residual gas and stray electric fields.



Search For Millicharged Particles



- ▶ Drive microsphere with oscillating electric field while removing electrons with Xe flash lamp.
- ▶ Moore et al., PRL **113** 251801 (2014)





Search For 'Chameleon' Interaction




- ▶ Drive microsphere with source mass.
- ▶ Microsphere is isolated from surrounding masses reducing screening effects.
- ▶ Rider et al., PRL **117**, 101101 (2016)

Current Work: Search for non-Newtonian Gravity




- ▶ Drive microsphere with source mass oscillating in density.
- ▶ Can probe into new parameter space.
- ▶ Fundamental backgrounds should not be significant.

Fundamental Backgrounds




- ▶ Casimir effect leads to attractive force from the cut off of long wavelength vacuum fluctuations.
- ▶ The effect is not entirely screened by a shield of finite thickness.
- ▶ Occurs with same periodicity as density modulation.

Apparatus



- ▶ Use interference to suppress backgrounds.
- ▶ Measure axial position of sphere with interferometer.

Interferometric Measurements



- ▶ Interferometer measurement of axial position calibrated into physical units by $\lambda = 1063.9nm$.
- ▶ Can fit PSD directly to Brownian motion of harmonic oscillator $S(\omega) = \frac{2k_b T}{M\Omega^2} \frac{\Omega^2 \Gamma_0}{(\Omega^2 - \omega^2)^2 + \omega^2 \Gamma_0^2}$.
- ▶ Get $T @ 1.5mbar = \sim 500K$.

Conclusion

- ▶ Optically levitated microspheres are a demonstrated force sensor with excellent isolation from the environment
- ▶ We have used optically levitated microspheres to search for millicharged particles as well as the chameleon interaction
- ▶ We expect to have a non-Newtonian gravity result soon