

Scale hierarchies and string cosmology

I. Antoniadis

Albert Einstein Center, University of Bern
and

LPTHE, UPMC/CNRS, Sorbonne Universités, Paris

BSM-2107 Beyond Standard Model: from Theory to Experiment


Hurghada, Egypt, 18-22 December 2017

Main predictions → inspirations for BSM physics

- Spacetime supersymmetry but arbitrary breaking scale
- Extra dimensions of space six or seven in M-theory
- Brane-world description of our Universe
matter and gauge interactions may be localised in less dimensions
- Landscape of vacua
- ...


Connect string theory to the real world

- Is it a tool for strong coupling dynamics or a theory of fundamental forces?
- If theory of Nature can it describe both particle physics and cosmology?



Problem of scales

- describe high energy (SUSY?) extension of the Standard Model
unification of all fundamental interactions
 - incorporate Dark Energy
simplest case: infinitesimal (tuneable) +ve cosmological constant
 - describe possible accelerated expanding phase of our universe
models of inflation (approximate de Sitter)
- ⇒ 3 very different scales besides M_{Planck} :



Relativistic dark energy 70-75% of the observable universe

negative pressure: $p = -\rho \Rightarrow$ cosmological constant

$$R_{ab} - \frac{1}{2}Rg_{ab} + \Lambda g_{ab} = \frac{8\pi G}{c^4} T_{ab} \Rightarrow \rho_\Lambda = \frac{c^4 \Lambda}{8\pi G} = -p_\Lambda$$

Two length scales:


- $[\Lambda] = L^{-2} \leftarrow$ size of the observable Universe

$$\Lambda_{obs} \simeq 0.74 \times 3H_0^2/c^2 \simeq 1.4 \times (10^{26} \text{ m})^{-2}$$

Hubble parameter $\simeq 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$

- $[\frac{\Lambda}{G} \times \frac{c^3}{h}] = L^{-4} \leftarrow$ dark energy length $\simeq 85 \mu\text{m}$

Problem of scales



① they are independent

② possible connections

- M_I could be near the EW scale, such as in Higgs inflation
but large non minimal coupling to explain
- M_{Planck} could be emergent from the EW scale
in models of low-scale gravity and TeV strings

What about M_I ? can it be at the TeV scale?

Can we infer M_I from cosmological data?

I.A.-Patil '14 and '15

- connect inflation and SUSY breaking scales

Inflation in supergravity: main problems

- slow-roll conditions: the eta problem \Rightarrow fine-tuning of the potential

$$\eta = V''/V, \quad V_F = e^K (|DW|^2 - 3|W|^2), \quad DW = W' + K'W$$

K : Kähler potential, W : superpotential

canonically normalised field: $K = X\bar{X} \Rightarrow \eta = 1 + \dots$

- trans-Planckian initial conditions \Rightarrow break validity of EFT
no-scale type models that avoid the η -problem
- stabilisation of the (pseudo) scalar companion of the inflaton
chiral multiplets \Rightarrow complex scalars
- moduli stabilisation, de Sitter vacuum, ...

Starobinsky model of inflation

$$\mathcal{L} = \frac{1}{2}R + \alpha R^2$$

$$\text{Lagrange multiplier } \phi \Rightarrow \mathcal{L} = \frac{1}{2}(1 + 2\phi)R - \frac{1}{4\alpha}\phi^2$$

Weyl rescaling \Rightarrow equivalent to a scalar field with exponential potential:


$$\mathcal{L} = \frac{1}{2}R - \frac{1}{2}(\partial\phi)^2 - \frac{M^2}{12} \left(1 - e^{-\sqrt{\frac{2}{3}}\phi}\right)^2 \quad M^2 = \frac{3}{4\alpha}$$

Note that the two metrics are not the same

supersymmetric extension:

add D-term $\mathcal{R}\bar{\mathcal{R}}$ because F-term \mathcal{R}^2 does not contain R^2

\Rightarrow brings two chiral multiplets



SUSY extension of Starobinsky model

$$K = -3 \ln(T + \bar{T} - C\bar{C}) \quad ; \quad W = MC(T - \frac{1}{2})$$

- T contains the inflaton: $\text{Re } T = e^{\sqrt{\frac{2}{3}}\phi}$
- $C \sim \mathcal{R}$ is unstable during inflation

⇒ add higher order terms to stabilize it

e.g. $C\bar{C} \rightarrow h(C, \bar{C}) = C\bar{C} - \zeta(C\bar{C})^2$ Kallosh-Linde '13

- SUSY is broken during inflation with C the goldstino superfield

→ model independent treatment in the decoupling sgoldstino limit

⇒ minimal SUSY extension that evades stability problem [13]

Non-linear supersymmetry \Rightarrow goldstino mode χ

Volkov-Akulov '73

Effective field theory of SUSY breaking at low energies

Analog of non-linear σ -model \Rightarrow constraint superfields

Rocek-Tseytlin '78, Lindstrom-Rocek '79, Komargodski-Seiberg '09

Goldstino: chiral superfield X_{NL} satisfying $X_{NL}^2 = 0 \Rightarrow$

$$\begin{aligned} X_{NL}(y) &= \frac{\chi^2}{2F} + \sqrt{2}\theta\chi + \theta^2 F & y^\mu &= x^\mu + i\theta\sigma^\mu\bar{\theta} \\ &= F\Theta^2 & \Theta &= \theta + \frac{\chi}{\sqrt{2}F} \end{aligned}$$

$$\mathcal{L}_{NL} = \int d^4\theta X_{NL}\bar{X}_{NL} - \frac{1}{\sqrt{2}\kappa} \left\{ \int d^2\theta X_{NL} + h.c. \right\} = \mathcal{L}_{Volkov-Akulov}$$

R-symmetry with $[\theta]_R = [\chi]_R = 1$ and $[X]_R = 2$

$$F = \frac{1}{\sqrt{2}\kappa} + \dots$$

Non-linear SUSY in supergravity

I.A.-Dudas-Ferrara-Sagnotti '14

$$K = -3 \log(1 - X\bar{X}) \equiv 3X\bar{X} \quad ; \quad W = fX + W_0 \quad \quad X \equiv X_{NL}$$

$$\Rightarrow \quad V = \frac{1}{3}|f|^2 - 3|W_0|^2 \quad ; \quad m_{3/2}^2 = |W_0|^2$$

- V can have any sign contrary to global NL SUSY
- NL SUSY in flat space $\Rightarrow f = 3 m_{3/2} M_p$
- R-symmetry is broken by W_0
- Dual gravitational formulation: $(\mathcal{R} - 6W_0)^2 = 0$ I.A.-Markou '15
↖ chiral curvature superfield
- Minimal SUSY extension of R^2 gravity

Non-linear Starobinsky supergravity [10]

$$K = -3 \ln(T + \bar{T} - X\bar{X}) \quad ; \quad W = MXT + fX + W_0 \quad \Rightarrow$$

$$\mathcal{L} = \frac{1}{2}R - \frac{1}{2}(\partial\phi)^2 - \frac{M^2}{12} \left(1 - e^{-\sqrt{\frac{2}{3}}\phi}\right)^2 - \frac{1}{2}e^{-2\sqrt{\frac{2}{3}}\phi}(\partial a)^2 - \frac{M^2}{18}e^{-2\sqrt{\frac{2}{3}}\phi}a^2$$

- axion a much heavier than ϕ during inflation, decouples:

$$m_\phi = \frac{M}{3}e^{-\sqrt{\frac{2}{3}}\phi_0} \ll m_a = \frac{M}{3}$$

- inflation scale M independent from NL-SUSY breaking scale f

\Rightarrow compatible with low energy SUSY

- however inflaton different from goldstino superpartner

- also initial conditions require trans-planckian values for ϕ ($\phi > 1$) [19]

Inflation from supersymmetry breaking

I.A.-Chatrabhuti-Isono-Knoops '16, '17

Inflaton : goldstino superpartner in the presence of a gauged R-symmetry


- linear superpotential $W = f X \Rightarrow$ no η -problem

$$\begin{aligned}V_F &= e^K (|DW|^2 - 3|W|^2) \\ &= e^K (|1 + K_X X|^2 - 3|X|^2) |f|^2 \quad K = X\bar{X} \\ &= e^{|X|^2} (1 - |X|^2 + \mathcal{O}(|X|^4)) |f|^2 = \mathcal{O}(|X|^4) \Rightarrow \eta = 0 + \dots\end{aligned}$$

- inflation around a maximum of scalar potential (hill-top) \Rightarrow small field
no large field initial conditions
- gauge R-symmetry: (pseudo) scalar absorbed by the $U(1)_R$
- vacuum energy at the minimum: tuning between V_F and V_D

Two classes of models

- Case 1: R-symmetry is restored during inflation (at the maximum)



- Case 2: R-symmetry is (spontaneously) broken everywhere
(and restored at infinity)

example: toy model of SUSY breaking [19] [28]

Case 1: R-symmetry restored during inflation

$$\mathcal{K}(X, \bar{X}) = \kappa^{-2} X \bar{X} + \kappa^{-4} A (X \bar{X})^2 \quad A > 0$$

$$W(X) = \kappa^{-3} f X \quad \Rightarrow$$

$$f(X) = 1 \quad (+\beta \ln X \text{ to cancel anomalies but } \beta \text{ very small})$$

$$\mathcal{V} = \mathcal{V}_F + \mathcal{V}_D$$

$$\mathcal{V}_F = \kappa^{-4} f^2 e^{X \bar{X} (1 + A X \bar{X})} \left[-3 X \bar{X} + \frac{(1 + X \bar{X} (1 + 2 A X \bar{X}))^2}{1 + 4 A X \bar{X}} \right]$$

$$\mathcal{V}_D = \kappa^{-4} \frac{q^2}{2} [1 + X \bar{X} (1 + 2 A X \bar{X})]^2$$

Assume inflation happens around the maximum $|X| \equiv \rho \simeq 0 \quad \Rightarrow$

Case 1: predictions

slow-roll parameters

$$\eta = \frac{1}{\kappa^2} \left(\frac{V''}{V} \right) = 2 \left(\frac{-4A + x^2}{2 + x^2} \right) + \mathcal{O}(\rho^2) \quad x = q/f$$

$$\epsilon = \frac{1}{2\kappa^2} \left(\frac{V'}{V} \right)^2 = 4 \left(\frac{-4A + x^2}{2 + x^2} \right)^2 \rho^2 + \mathcal{O}(\rho^4) \simeq \eta^2 \rho^2$$

η small: for instance $x \ll 1$ and $A \sim \mathcal{O}(10^{-1})$

inflation starts with an initial condition for $\phi = \phi_*$ near the maximum and ends when $|\eta| = 1$

$$\Rightarrow \text{number of e-folds } N = \int_{end}^{start} \frac{V}{V'} = \kappa \int \frac{1}{\sqrt{2\epsilon}} \simeq \frac{1}{|\eta_*|} \ln \left(\frac{\rho_{end}}{\rho_*} \right)$$

Case 1: predictions

amplitude of density perturbations $A_s = \frac{\kappa^4 V_*}{24\pi^2 \epsilon_*} = \frac{\kappa^2 H_*^2}{8\pi^2 \epsilon_*}$

spectral index $n_s = 1 + 2\eta_* - 6\epsilon_* \simeq 1 + 2\eta_*$

tensor – to – scalar ratio $r = 16\epsilon_*$

Planck '15 data : $\eta \simeq -0.02$, $A_s \simeq 2.2 \times 10^{-9}$, $N \gtrsim 50$

$\Rightarrow r \lesssim 10^{-4}$, $H_* \lesssim 10^{12}$ GeV assuming $\rho_{\text{end}} \lesssim 1/2$

Question: can a 'nearby' minimum exist with a tiny +ve vacuum energy?

Answer: Yes in a 'weaker' sense: perturbative expansion [15]

valid for the Kähler potential but not for the slow-roll parameters

generic V (not fine-tuned) $\Rightarrow 10^{-9} \lesssim r \lesssim 10^{-4}$, $10^{10} \lesssim H_* \lesssim 10^{12}$ GeV [34]

impose independent scales: proceed in 2 steps

- 1 SUSY breaking at $m_{SUSY} \sim \text{TeV}$

with an infinitesimal (tuneable) positive cosmological constant

Villadoro-Zwirner '05

I.A.-Knoops, I.A.-Ghilenca-Knoops '14, I.A.-Knoops '15

- 2 Inflation connected or independent? [7] [11] [28]

Toy model for SUSY breaking

Content (besides $N = 1$ SUGRA): one vector V and one chiral multiplet S
with a shift symmetry $S \rightarrow S - icw \leftarrow$ transformation parameter

String theory: compactification modulus or universal dilaton

$$s = 1/g^2 + ia \leftarrow \text{dual to antisymmetric tensor}$$

Kähler potential K : function of $S + \bar{S}$

$$\text{string theory: } K = -p \ln(S + \bar{S})$$

Superpotential: constant or single exponential if R-symmetry $W = ae^{bS}$

$$\int d^2\theta W \text{ invariant}$$

$$b < 0 \Rightarrow \text{non perturbative}$$

[24] [23]

Scalar potential

$$\mathcal{V}_F = a^2 e^{\frac{b}{l}} l^{p-2} \left\{ \frac{1}{p} (pl - b)^2 - 3l^2 \right\} \quad l = 1/(s + \bar{s})$$

Planck units

- $b > 0 \Rightarrow$ SUSY local minimum in AdS space with $l = b/p$
- $b \leq 0 \Rightarrow$ no minimum with $l > 0$ ($p \leq 3$)

but interesting metastable SUSY breaking vacuum when R-symmetry is gauged by V allowing a Fayet-Iliopoulos (FI) term:

$$\mathcal{V}_D = c^2 l (pl - b)^2 \quad \text{for gauge kinetic function } f(S) = S$$

- $b > 0$: $\mathcal{V} = \mathcal{V}_F + \mathcal{V}_D$ SUSY AdS minimum remains
- $b = 0$: SUSY breaking minimum in AdS ($p < 3$)
- $b < 0$: SUSY breaking minimum with tuneable cosmological constant Λ

minimisation and spectrum

Minimisation of the potential: $V' = 0$, $V = \Lambda$

In the limit $\Lambda \approx 0$ ($p = 2$) \Rightarrow [30]


$$b/l = \rho \approx -0.183268 \quad \Rightarrow \langle l \rangle = b/\rho$$

$$\frac{a^2}{bc^2} = 2 \frac{e^{-\rho}}{\rho} \frac{(2-\rho)^2}{2+4\rho-\rho^2} + \mathcal{O}(\Lambda) \approx -50.6602 \quad \Rightarrow c \propto a$$

Physical spectrum:

massive dilaton, $U(1)$ gauge field, Majorana fermion, gravitino

All masses of order $m_{3/2} \approx e^{\rho/2} l a \leftarrow$ TeV scale



[28] [26]

Properties and generalizations

- Metastability of the ground state: extremely long lived

$$l \simeq 0.02 \text{ (GUT value } \alpha_{GUT}/2) \quad m_{3/2} \sim \mathcal{O}(\text{TeV}) \Rightarrow$$

$$\text{decay rate } \Gamma \sim e^{-B} \text{ with } B \approx 10^{300}$$

- Add visible sector (MSSM) preserving the same vacuum

matter fields ϕ neutral under R-symmetry

$$K = -2 \ln(S + \bar{S}) + \phi^\dagger \phi \quad ; \quad W = (a + W_{MSSM}) e^{bS}$$

\Rightarrow soft scalar masses non-tachyonic of order $m_{3/2}$ (gravity mediation)

- Toy model classically equivalent to [20]


$$K = -p \ln(S + \bar{S}) + b(S + \bar{S}) \quad ; \quad W = a \quad \text{with } V \text{ ordinary } U(1)$$

- Dilaton shift can be identified with $B - L \supset$ matter parity $(-)^{B-L}$

Properties and generalizations

- R-charged fields needed for anomaly cancellation
- A simple (anomaly free) variation: $f = 1$ and $p = 1$
tuning still possible but scalar masses of neutral matter tachyonic
possible solution: add a new field Z in the 'hidden' SUSY sector
 \Rightarrow one extra parameter
- alternatively: add an S -dependent factor in Matter kinetic terms
$$K = -\ln(S + \bar{S}) + (S + \bar{S})^{-\nu} \sum \Phi \bar{\Phi} \quad \text{for } \nu \gtrsim 2.5$$
or the $B - L$ unit charge of SM particles \Rightarrow similar phenomenology
- distinct features from other models of SUSY breaking and mediation
- gaugino masses at the quantum level
 \Rightarrow suppressed compared to scalar masses and A-terms

Typical spectrum



The masses of sbottom squark (yellow), stop (black), gluino (red), lightest chargino (green) and lightest neutralino (blue) as a function of the gravitino mass. The mass of the lightest neutralino varies between ~ 40 and 150 GeV [19]

ATLAS SUSY SEARCHES SUMMARY

ATLAS SUSY Searches* - 95% CL Lower Limits

May 2017

ATLAS Preliminary

$\sqrt{s} = 7, 8, 13 \text{ TeV}$

\tilde{q}, \tilde{g}

$\tilde{\nu}_\tau, \tilde{b}$

$\tilde{\chi}^0, \tilde{\chi}^\pm, \tilde{t}, \tilde{b}$

Long-lived particles

RPV

Model	$\epsilon, \mu, \tau, \gamma$	Jets	$\frac{E_{T,miss}}{E_T}$	$[\mathcal{L} dt](\text{fb}^{-1})$	Mass limit	$\sqrt{s} = 7, 8 \text{ TeV}$	$\sqrt{s} = 13 \text{ TeV}$	Reference		
Include Searches	MSUGRA/CMSM	$0-3 e, \mu, \tau, \gamma$	2-10 jets/3 $\tilde{\chi}^0$	Yes	20.3	\tilde{g}	1.85 TeV	$m(\tilde{g})=m(\tilde{t}_1)$ $m(\tilde{t}_1)=200 \text{ GeV}, m(\tilde{t}_2)=m(\tilde{g})+2m(\tilde{g}, g)$	1507.05325 ATLAS-CONF-2017-022	
	$\tilde{g}, \tilde{q} \rightarrow q\tilde{g}$		0-2 jets	Yes	36.1	\tilde{q}	1.57 TeV	$m(\tilde{q})=m(\tilde{t}_1)+5 \text{ GeV}$	1604.07773 ATLAS-CONF-2017-022	
	$\tilde{g}, \tilde{q} \rightarrow q\tilde{g}$ (compressed)		mono-jet	1-3 jets	Yes	3.2	\tilde{g}	606 GeV	$m(\tilde{g})=200 \text{ GeV}$	ATLAS-CONF-2017-022
	$\tilde{g}, \tilde{q} \rightarrow q\tilde{g}$		0	2-6 jets	Yes	36.1	\tilde{g}	2.02 TeV	$m(\tilde{g})=200 \text{ GeV}$	ATLAS-CONF-2017-022
	$\tilde{g}, \tilde{q} \rightarrow q\tilde{g}$		0	2-6 jets	Yes	36.1	\tilde{q}	2.01 TeV	$m(\tilde{q})=200 \text{ GeV}, m(\tilde{t}_1)=0.5m(\tilde{t}_2)=m(\tilde{g})$	ATLAS-CONF-2017-022
	$\tilde{g}, \tilde{q} \rightarrow q\tilde{g}$		3 e, μ, τ	4 jets	-	36.1	\tilde{g}	1.825 TeV	$m(\tilde{g})=400 \text{ GeV}$	ATLAS-CONF-2017-030
	$\tilde{g}, \tilde{q} \rightarrow q\tilde{g}$		0-7 jets	Yes	36.1	\tilde{g}	1.8 TeV	$m(\tilde{g})=400 \text{ GeV}$	ATLAS-CONF-2017-033	
	GMSB (bino NLSP)		1.2 + 0.1 \tilde{t}	0-3 jets	Yes	3.2	\tilde{g}	2.8 TeV	$\tau(\text{NLSP}) > 0.1 \text{ ns}$	1607.05979 1606.09150
	GGM (bino NLSP)		2 γ	Yes	3.2	\tilde{g}	1.65 TeV	$m(\tilde{g})=200 \text{ GeV}, \tau(\text{NLSP}) > 0.1 \text{ ns}, \mu=0$	1505.02066	
	GGM (higgsino-bino NLSP)		7	1 \tilde{b}	Yes	20.3	\tilde{g}	1.37 TeV	$m(\tilde{g})=200 \text{ GeV}, \tau(\text{NLSP}) > 0.1 \text{ ns}, \mu=0$	1507.05493
GGM (higgsino-bino NLSP)		7	2 jets	Yes	13.5	\tilde{g}	1.8 TeV	$m(\tilde{g})=200 \text{ GeV}, \tau(\text{NLSP}) > 0.1 \text{ ns}, \mu=0$	ATLAS-CONF-2016-056	
GGM (higgsino NLSP)		2 e, μ (2)	2 jets	Yes	26.1	\tilde{g}	900 GeV	$m(\tilde{g})=1.8 \times 10^4 \text{ eV}, m(\tilde{g})=0.15 \text{ TeV}$	1502.01518	
Gravitino LSP		0	mono-jet	Yes	20.3	\tilde{g}	865 GeV			
$\tilde{\nu}_\tau, \tilde{b}$	$\tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		0	3 \tilde{b}	Yes	36.1	\tilde{b}	1.23 TeV	$m(\tilde{b})=200 \text{ GeV}$	ATLAS-CONF-2017-021
	$\tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		0-1 e, μ	3 \tilde{b}	Yes	36.1	\tilde{b}	1.97 TeV	$m(\tilde{b})=200 \text{ GeV}$	ATLAS-CONF-2017-021
	$\tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		0-1 e, μ	3 \tilde{b}	Yes	20.1	\tilde{b}	1.37 TeV	$m(\tilde{b})=200 \text{ GeV}$	1467.06000
$\tilde{\chi}^0, \tilde{\chi}^\pm, \tilde{t}, \tilde{b}$	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		0	2 \tilde{b}	Yes	36.1	\tilde{b}	590 GeV	$m(\tilde{b})=200 \text{ GeV}$	ATLAS-CONF-2017-038
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		2 e, μ (SS)	1 \tilde{b}	Yes	35.1	\tilde{b}	275-700 GeV	$m(\tilde{b})=200 \text{ GeV}, m(\tilde{t}_1)=m(\tilde{t}_2)+100 \text{ GeV}$	ATLAS-CONF-2017-030
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		0.2 e, μ	1.2 \tilde{b}	Yes	4.713.9	\tilde{b}	117-178 GeV	$m(\tilde{b})=1.2m(\tilde{t}_1), m(\tilde{t}_1)=35 \text{ GeV}$	1303.2102, ATLAS-CONF-2016-077
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		0.2 e, μ	0.2 jets/1-2 \tilde{b}	Yes	20.3, 36.1	\tilde{b}	30-198 GeV	$m(\tilde{b})=1 \text{ GeV}$	1508.08816, ATLAS-CONF-2017-020
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		0	mono-jet	Yes	3.2	\tilde{b}	90-323 GeV	$m(\tilde{b})=1 \text{ GeV}$	1604.07773
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		2 e, μ (2)	1 \tilde{b}	Yes	20.3	\tilde{b}	150-600 GeV	$m(\tilde{b})=150 \text{ GeV}$	1403.3202
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		3 e, μ (2)	1 \tilde{b}	Yes	36.1	\tilde{b}	296-739 GeV	$m(\tilde{b})=1 \text{ GeV}$	ATLAS-CONF-2017-019
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		1-2 e, μ	4 \tilde{b}	Yes	36.1	\tilde{b}	320-880 GeV	$m(\tilde{b})=1 \text{ GeV}$	ATLAS-CONF-2017-019
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		2 e, μ, τ	0	Yes	36.1	\tilde{b}	90-440 GeV	$m(\tilde{b})=0$	ATLAS-CONF-2017-039
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		2 e, μ, τ	0	Yes	36.1	\tilde{b}	710 GeV	$m(\tilde{b})=0, m(\tilde{t}_1)=0.5m(\tilde{t}_2)=m(\tilde{b})$	ATLAS-CONF-2017-039
EW direct	$\tilde{\chi}^0, \tilde{\chi}^\pm, \tilde{t}, \tilde{b}$		2 e, μ, τ	0	Yes	36.1	\tilde{b}	760 GeV	$m(\tilde{b})=0, m(\tilde{t}_1)=m(\tilde{t}_2)+0.5m(\tilde{t}_1)=m(\tilde{b})$	ATLAS-CONF-2017-030
	$\tilde{\chi}^0, \tilde{\chi}^\pm, \tilde{t}, \tilde{b}$		2 e, μ, τ	0	Yes	36.1	\tilde{b}	590 GeV	$m(\tilde{b})=0, m(\tilde{t}_1)=m(\tilde{t}_2)+0.5m(\tilde{t}_1)=m(\tilde{b})$	ATLAS-CONF-2017-030
	$\tilde{\chi}^0, \tilde{\chi}^\pm, \tilde{t}, \tilde{b}$		2.3 jets	0-2 jets	Yes	36.1	\tilde{b}	590 GeV	$m(\tilde{b})=0, m(\tilde{t}_1)=m(\tilde{t}_2)+0.5m(\tilde{t}_1)=m(\tilde{b})$	ATLAS-CONF-2017-030
	$\tilde{\chi}^0, \tilde{\chi}^\pm, \tilde{t}, \tilde{b}$		e, μ, τ	0-2 \tilde{b}	Yes	20.3	\tilde{b}	270 GeV	$m(\tilde{b})=0, m(\tilde{t}_1)=0, \tilde{t}$ decoupled	1501.07110
	$\tilde{\chi}^0, \tilde{\chi}^\pm, \tilde{t}, \tilde{b}$		e, μ, τ	0	Yes	20.3	\tilde{b}	635 GeV	$m(\tilde{b})=0, m(\tilde{t}_1)=0, \tilde{t}$ decoupled	1465.50086
	GGM (bino NLSP) weak prod., $\tilde{b} \rightarrow b\tilde{g}$		1 e, μ, τ	\tilde{b}	Yes	20.3	\tilde{b}	115-570 GeV	$m(\tilde{b})=0, m(\tilde{t}_1)=0, m(\tilde{t}_2)=0.5m(\tilde{t}_1)=0.2 \text{ m}$	1507.05493
	GGM (bino NLSP) weak prod., $\tilde{b} \rightarrow b\tilde{g}$		2 γ	\tilde{b}	Yes	20.3	\tilde{b}	900 GeV	$\tau < 1 \text{ ns}$	1507.05493
	Direct $\tilde{t}\tilde{t}^*$ prod., long-lived \tilde{t}_1^*		Disapp. trk	1 jet	Yes	36.1	\tilde{t}	430 GeV	$m(\tilde{t}_1)=m(\tilde{t}_2)=160 \text{ MeV}, m(\tilde{t}_1^*)=0.2 \text{ m}$	ATLAS-CONF-2017-017
	Direct $\tilde{t}\tilde{t}^*$ prod., long-lived \tilde{t}_1^*		dE/dx trk	Yes	18.4	\tilde{t}	495 GeV	$m(\tilde{t}_1)=m(\tilde{t}_2)=160 \text{ MeV}, m(\tilde{t}_1^*)=15 \text{ ns}$	1506.05332	
	Stable \tilde{t} R-hadron		0	1-5 jets	Yes	27.9	\tilde{t}	850 GeV	$m(\tilde{t})=100 \text{ GeV}, 10 \mu\text{s} < \tau < 1000 \text{ s}$	1310.65064
Metastable \tilde{t} R-hadron		0	1-5 jets	Yes	3.2	\tilde{t}	1.52 TeV	$m(\tilde{t})=100 \text{ GeV}, \tau > 10 \text{ ns}$	1606.05239	
GMSB, stable $\tilde{t}, \tilde{t}_1^* \rightarrow \tilde{t}, \tilde{b}, \tilde{g}, \mu, \tau, \nu$		1.2 μ	Yes	19.1	\tilde{t}	537 GeV	10-targ-50	1411.67956		
GMSB, $\tilde{t}_1^* \rightarrow \tilde{t}, \tilde{b}, \tilde{g}$, long-lived \tilde{t}_1^*		2 γ	Yes	20.3	\tilde{t}	440 GeV	1-coupled-3-ne, SPSS model	1469.3542		
GGM $\tilde{g}\tilde{g} \rightarrow \tilde{g}\tilde{g}$		displ. $\nu\tau/\mu\tau$	Yes	20.3	\tilde{g}	1.8 TeV	7 $\nu\tau/\mu\tau < 0.740 \text{ ms}, m(\tilde{g})=1.3 \text{ TeV}$	1504.05162		
GGM $\tilde{g}\tilde{g} \rightarrow \tilde{g}\tilde{g}$		displ. $\nu\tau + \text{jets}$	Yes	20.3	\tilde{g}	1.8 TeV	6 $\nu\tau/\mu\tau < 0.480 \text{ ms}, m(\tilde{g})=1.1 \text{ TeV}$	1504.05162		
Long-lived particles	LFV $\tilde{g}\tilde{g} \rightarrow X, \tilde{b}, \tilde{g} \rightarrow \nu\tau/\mu\tau$		e, μ, τ, ν	-	3.2	\tilde{g}	1.8 TeV	$A_{\tilde{g}\tilde{g}}=0.11, A_{\tilde{b}\tilde{g}}=0.07$	1607.05979	
	Bilinear RPV CMSM		2 e, μ (SS)	0-3 \tilde{b}	Yes	20.3	\tilde{b}	1.45 TeV	$m(\tilde{b})=m(\tilde{t}_1), A_{1121}=1$	1464.2500
	$\tilde{t}_1^* \tilde{t}_1^* \rightarrow W\tilde{t}_1^*, \tilde{t}_1^* \tilde{t}_1^* \rightarrow \nu\tau, \mu\tau, \nu\mu, \nu\tau, \nu\mu$		4 e, μ, τ	Yes	13.3	\tilde{t}	1.14 TeV	$m(\tilde{t}_1)=9000 \text{ GeV}, A_{1120}(\theta=1.2)$	ATLAS-CONF-2016-075	
	$\tilde{t}_1^* \tilde{t}_1^* \rightarrow W\tilde{t}_1^*, \tilde{t}_1^* \tilde{t}_1^* \rightarrow \nu\tau, \mu\tau, \nu\mu, \nu\tau, \nu\mu$		3 e, μ, τ	Yes	20.3	\tilde{t}	450 GeV	$\text{RPV} \rightarrow \text{RPV}, A_{1120}(\theta=1.2)$	1465.50086	
	$\tilde{b}, \tilde{b} \rightarrow \nu\tilde{b}$		0	4.5 large- β jets	Yes	14.8	\tilde{b}	1.08 TeV	$m(\tilde{b})=200 \text{ GeV}, A_{1120}(\theta=1.2)$	ATLAS-CONF-2016-057
	$\tilde{b}, \tilde{b} \rightarrow \nu\tilde{b}$		0	4.5 large- β jets	-	14.8	\tilde{b}	1.35 TeV	$m(\tilde{b})=200 \text{ GeV}$	ATLAS-CONF-2016-057
	$\tilde{b}, \tilde{b} \rightarrow \nu\tilde{b}, \tilde{b} \rightarrow \nu\tilde{b}$		1 e, μ	8-10 jets/0-4 \tilde{b}	-	36.1	\tilde{b}	2.1 TeV	$m(\tilde{b})=1 \text{ TeV}, A_{1120}$	ATLAS-CONF-2017-013
	$\tilde{b}, \tilde{b} \rightarrow \nu\tilde{b}, \tilde{b} \rightarrow \nu\tilde{b}$		1 e, μ	8-10 jets/0-4 \tilde{b}	-	36.1	\tilde{b}	1.65 TeV	$m(\tilde{b})=1 \text{ TeV}, A_{1120}$	ATLAS-CONF-2017-013
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		0	2 jets + 2 \tilde{b}	-	15.4	\tilde{b}	410 GeV	$m(\tilde{b})=1 \text{ TeV}, A_{1120}$	ATLAS-CONF-2016-022, ATLAS-CONF-2016-084
	$\tilde{b}, \tilde{b}, \tilde{b} \rightarrow b\tilde{b}$		2 e, μ, τ	2 \tilde{b}	-	36.1	\tilde{b}	450-510 GeV	$\text{RPV} \rightarrow \text{RPV}, A_{1120}$	ATLAS-CONF-2017-036
Other	Scalar charm, $\tilde{c} \rightarrow c\tilde{g}$		0	2 \tilde{c}	Yes	20.3	\tilde{c}	510 GeV	$m(\tilde{c})=200 \text{ GeV}$	1501.01325

*Only a selection of the available mass limits on new states or phenomena is shown. Many of the limits are based on simplified models, c.f. Refs. for the assumptions made.

Mass scale [TeV]

Direct, one-step and two-step decays

Simplified and pMSSM inspired models

Higher reach for electro-weakinos

Limits on the long lived chargino

Limits for RPV models

Case 2 example: toy model of SUSY breaking

I.A.-Chatrabhuti-Isono-Knoops '16

Can the dilaton be the inflaton in the simple model of SUSY breaking based on a gauged shift symmetry?

the only physical scalar left over, partner (partly) of the goldstino
partly because of a D-term auxiliary component

Same potential cannot satisfy the slow roll condition $|\eta| = |V''/V| \ll 1$
with the dilaton rolling towards the Standard Model minimum

\Rightarrow need to create an appropriate plateau around the maximum of V [23]
without destroying the properties of the SM minimum

\Rightarrow study possible corrections to the Kähler potential

only possibility compatible with the gauged shift symmetry [31]

Extensions of the SUSY breaking model

Parametrize the general **correction** to the Kähler potential:

$$K = -p\kappa^{-2} \log \left(s + \bar{s} + \frac{\xi}{b} F(s + \bar{s}) \right) + \kappa^{-2} b(s + \bar{s})$$

$$W = \kappa^{-3} a, \quad f(s) = \gamma + \beta s$$

$$\mathcal{P} = \kappa^{-2} c \left(b - p \frac{1 + \frac{\xi}{b} F'}{s + \bar{s} + \frac{\xi}{b} F} \right)$$

Three types of possible corrections:

- perturbative: $F \sim (s + \bar{s})^{-n}$, $n \geq 0$
- non-perturbative D-brane instantons: $F \sim e^{-\delta(s+\bar{s})}$, $\delta > 0$
- non-perturbative NS5-brane instantons: $F \sim e^{-\delta(s+\bar{s})^2}$, $\delta > 0$

Only the last can lead to slow-roll conditions with sufficient inflation

Slow-roll inflation

$F = \xi e^{\alpha b^2 \phi^2}$ with $\phi = s + \bar{s} = 1/l \Rightarrow$ two extra parameters $\alpha < 0$, ξ
they control the shape of the potential

slow-roll conditions: $\epsilon = 1/2(V'/V)^2 \ll 1$, $|\eta| = |V''/V| \ll 1$

\Rightarrow allowed regions of the parameter space with $|\xi|$ small

additional independent parameters: a, c, b

SM minimum with tuneable cosmological constant Λ : $V' = 0$, $V = \Lambda \approx 0$

$\xi = 0 \Rightarrow b\phi_{min} = \rho_0$, $\frac{a^2}{bc^2} = \lambda_0$ with ρ_0, λ_0 calculable constants [22]


b controls $\phi_{min} \sim 1/g_s$ choose it of order 10

tuning determines a in terms of c overall scale of the potential

$\xi \neq 0 \Rightarrow \rho_0, \lambda_0$ become functions $I(\xi, \alpha), \lambda(\xi, \alpha)$

numerical analysis \Rightarrow mild dependence

$\xi = 0.025, \alpha = -4.8, p = 2, b = -0.018$



Fit Planck '15 data and predictions

$p = 1$: similar analysis \Rightarrow

$$\phi_* = 64.53, \xi = 0.30, \alpha = -0.78, b = -0.023, c = 10^{-13}$$

N	n_s	r	A_s
889	0.959	4×10^{-22}	2.205×10^{-9}

SM minimum: $\langle \phi \rangle \approx 21.53$, $\langle m_{3/2} \rangle = 18.36$ TeV, $\langle M_{A_\mu} \rangle = 36.18$ TeV


During inflation:

$$H_* = \kappa \sqrt{\mathcal{V}_*/3} = 5.09 \text{ TeV}, m_{3/2}^* = 4.72 \text{ TeV}, M_{A_\mu}^* = 6.78 \text{ TeV}$$

Low energy spectrum essentially the same with $\xi = 0$:

$$m_0^2 = m_{3/2}^2 [-2 + \mathcal{C}], \quad A_0 = m_{3/2} \mathcal{C}, \quad B_0 = A_0 - m_{3/2}$$

$\mathcal{C} = 1.53$ vs at $\xi = 0$: $\mathcal{C}_0 = 1.52$, $m_{3/2}^0 = 17.27$, although $\langle \phi \rangle_0 \approx 9.96$ [15]



Conclusions

String pheno: consistent framework for particle physics and cosmology

Challenge of scales: at least three very different (besides M_{Planck})
electroweak, dark energy, inflation, SUSY?

their origins may be connected or independent

SUSY with infinitesimal (tuneable) +ve cosmological constant

- interesting framework for model building incorporating dark energy
- identify inflaton with goldstino superpartner
inflation at the SUSY breaking scale (TeV?)

General class of models with inflation from SUSY breaking:

(gauged) R-symmetry restored (case 1) or broken (case 2) during inflation
small field, avoids the η -problem, no (pseudo) scalar companion