

# One-loop exclusive diffractive processes in the CGC framework

Renaud Boussarie

Institute of Nuclear Physics PAN

Excited QCD 2017


RB, A.V.Grabovsky, L.Szymanowski, S.Wallon  
JHEP 409 (2014) 026 and JHEP 1611 (2016) 149

RB, A.V.Grabovsky, D.Yu.Ivanov, L.Szymanowski, S.Wallon  
arXiv:1612.08026 [hep-ph]


## Diffractive DIS

Rapidity gap events at HERA

Experiments at HERA : about 10% of scattering events reveal a **rapidity gap**



DIS events




DDIS events

**Rapidity gap  $\equiv$  Pomeron exchange**

DIS : Deep Inelastic Scattering, DDIS : Diffractive DIS

## Diffractive DIS

Theoretical approaches for DDIS using perturbative QCD

## • Collinear factorization approach


- Moderate- $x$  approach  $Q^2 \sim s$
- Relies on a QCD factorization theorem
- One needs to introduce a **diffractive distribution function**

•  $k_T$  factorization approach for two exchanged gluons

- low- $x$  QCD approach :  $s \gg Q^2 \gg \Lambda_{QCD}$
- The pomeron is described as a **two-gluon color singlet state**



A recent analysis of diffractive dijet production in DIS at HERA seems to **favor  $k_t$  factorization in the small diffractive mass regime** [ZEUS collaboration, 2015]

## Diffractive DIS




- Shockwave (CGC) approach

- low-x QCD approach :  $s \gg Q^2 \gg \Lambda_{QCD}$
- The pomeron exchange is described as the action of a **color singlet** Wilson line operator on the target states



[Bartels, Wüsthoff]

## Kinematics



$$p_1 = p^+ n_1 - \frac{Q^2}{2s} n_2$$

$$p_2 = \frac{m_t^2}{2p_2^-} n_1 + p_2^- n_2$$

$$p^+ \sim p_2^- \sim \sqrt{\frac{s}{2}}$$

$$(p_1 + p_2)^2 = s \gg Q_H^2 \gg \Lambda_{QCD}^2$$

Lightcone (Sudakov) vectors


$$n_1 = \sqrt{\frac{1}{2}}(1, 0_\perp, 1), \quad n_2 = \sqrt{\frac{1}{2}}(1, 0_\perp, -1), \quad (n_1 \cdot n_2) = 1$$

Lightcone coordinates:

$$x = (x^0, x^1, x^2, x^3) \rightarrow (x^+, x^-, \vec{x})$$

$$x^+ = x_- = (x \cdot n_2) \quad x^- = x_+ = (x \cdot n_1)$$



# Rapidity separation



Let us split the gluonic field between "fast" and "slow" gluons

$$\begin{aligned}
 \mathcal{A}^{\mu a}(k^+, k^-, \vec{k}) &= \mathcal{A}_\eta^{\mu a}(|k^+| < e^\eta p^+, k^-, \vec{k}) \\
 &+ \mathcal{B}_\eta^{\mu a}(|k^+| > e^\eta p^+, k^-, \vec{k})
 \end{aligned}$$

# Large longitudinal boost to the projectile frame



$$b^+(x^+, x^-, \vec{x})$$

$$b^-(x^+, x^-, \vec{x})$$

$$b^k(x^+, x^-, \vec{x})$$

 $\longrightarrow$ 

$$\frac{1}{\Lambda} b^+(\Lambda x^+, \frac{x^-}{\Lambda}, \vec{x})$$

$$\Lambda b^-(\Lambda x^+, \frac{x^-}{\Lambda}, \vec{x})$$

$$b^k(\Lambda x^+, \frac{x^-}{\Lambda}, \vec{x})$$

$$\Lambda \sim \sqrt{\frac{s}{m_t^2}}$$

$$b^\mu(x) \rightarrow b^-(x) n_2^\mu = \delta(x^+) \mathbf{B}(\vec{x}) n_2^\mu + \left( \sqrt{\frac{m_t^2}{s}} \right)$$

# QCD Lagrangian

Rewrite the QCD Lagrangian in terms of the  
"fast" **internal** field and the "slow" **external** field

$$\begin{aligned}
 \mathcal{L}_{free} &= gf_{abc}[(A_\eta^b \cdot \partial)A_\eta^a]A_\eta^c - \frac{1}{4}g^2 f_{abc} f_{ade}(A_\eta^a \cdot A_\eta^d)(A_\eta^b \cdot A_\eta^e) \\
 &+ i\bar{\psi}[-igt^a \hat{A}_\eta^a]\psi + i\bar{\psi}[-igt^a \hat{b}_\eta^a]\psi - gf_{abc}((b_\eta^b \cdot \partial)A_\eta^a) \cdot A_\eta^c \\
 &- gf_{abc}(((A_\eta^b \cdot \partial)A_\eta^a) \cdot b_\eta^c + ((b_\eta^b \cdot \partial)A_\eta^a) \cdot b_\eta^c \\
 &+ ((A_\eta^b \cdot \partial)b_\eta^a) \cdot A_\eta^c + ((b_\eta^b \cdot \partial)b_\eta^a) \cdot A_\eta^c] \\
 &- \frac{1}{4}g^2 f_{abc} f_{ade}[(A_\eta^a \cdot A_\eta^d)((b_\eta^e \cdot A_\eta^b) + (b_\eta^b \cdot A_\eta^e)) \\
 &+ (b_\eta^d \cdot A_\eta^a)((A_\eta^b \cdot A_\eta^e) + (b_\eta^e \cdot A_\eta^b) + (b_\eta^b \cdot A_\eta^e)) \\
 &+ (b_\eta^a \cdot A_\eta^d)((A_\eta^b \cdot A_\eta^e) + (b_\eta^e \cdot A_\eta^b) + (b_\eta^b \cdot A_\eta^e))]
 \end{aligned}$$

Gray terms cancel in **lightcone gauge**  $(n_2 \cdot \mathcal{A}) = 0$



# Propagator through the external shockwave field


$$G(z_2, z_0) = - \int d^4 z_1 \theta(z_2^+) \delta(z_1^+) \theta(-z_0^+) G(z_2 - z_1) \gamma^+ G(z_1 - z_0) U_1$$

Wilson lines :


$$U_i^\eta = U_{\bar{z}_i}^\eta = P \exp \left[ ig \int_{-\infty}^{+\infty} b_\eta^-(z_i^+, \bar{z}_i) dz_i^+ \right]$$

$$U_i^\eta = 1 + ig \int_{-\infty}^{+\infty} b_\eta^-(z_i^+, \bar{z}_i) dz_i^+ + (ig)^2 \int_{-\infty}^{+\infty} b_\eta^-(z_i^+, \bar{z}_i) b_\eta^-(z_j^+, \bar{z}_j) \theta(z_{ji}^+) dz_i^+ dz_j^+$$

...



## Factorized picture



Factorized amplitude


$$\mathcal{A}^\eta = \int d^{D-2} \vec{z}_1 d^{D-2} \vec{z}_2 \Phi^\eta(\vec{z}_1, \vec{z}_2) \langle P' | [\text{Tr}(U_{\vec{z}_1}^\eta U_{\vec{z}_2}^{\eta\dagger}) - N_c] | P \rangle$$

Dipole operator  $U_{ij}^\eta = \frac{1}{N_c} \text{Tr}(U_{\vec{z}_i}^\eta U_{\vec{z}_j}^{\eta\dagger}) - 1$

Written similarly for any number of Wilson lines in any color representation!

# Evolution for the dipole operator

$$\mathcal{U}_{12}^{\eta+\delta\eta} - \mathcal{U}_{12}^{\eta}$$



## B-JIMWLK hierarchy of equations

[Balitsky, Jalilian-Marian, Iancu, McLerran, Weigert, Leonidov, Kovner]


$$\frac{\partial \mathcal{U}_{12}^{\eta}}{\partial \eta} = \frac{\alpha_s N_c}{2\pi^2} \int d\vec{z}_3 \frac{\vec{z}_{12}^2}{\vec{z}_{13}^2 \vec{z}_{23}^2} [\mathcal{U}_{13}^{\eta} + \mathcal{U}_{32}^{\eta} - \mathcal{U}_{12}^{\eta} + \mathcal{U}_{13}^{\eta} \mathcal{U}_{32}^{\eta}]$$

$$\frac{\partial \mathcal{U}_{13}^{\eta} \mathcal{U}_{32}^{\eta}}{\partial \eta} = \dots$$

Evolves a dipole into a double dipole

# The BK equation

Mean field approximation, or 't Hooft planar limit  $N_c \rightarrow \infty$  in Balitsky's equation



⇒ **BK equation** [Balitsky, 1995] [Kovchegov, 1999]

$$\frac{\partial \langle \mathcal{U}_{12}^n \rangle}{\partial \eta} = \frac{\alpha_s N_c}{2\pi^2} \int d\vec{z}_3 \frac{\vec{z}_{12}^2}{\vec{z}_{13}^2 \vec{z}_{23}^2} [\langle \mathcal{U}_{13}^n \rangle + \langle \mathcal{U}_{32}^n \rangle - \langle \mathcal{U}_{12}^n \rangle + \langle \mathcal{U}_{13}^n \rangle \langle \mathcal{U}_{32}^n \rangle]$$

BFKL/BKP part

Triple pomeron vertex

Non-linear term : **saturation**

# The JIMWLK Hamiltonian

## Hamiltonian formulation of the hierarchy of equations

For an operator built from  $n$  Wilson lines, the JIMWLK evolution is given at LO accuracy by

$$\frac{\partial}{\partial \eta} \left[ U_{\vec{z}_1}^\eta \dots U_{\vec{z}_n}^\eta \right] = \sum_{i,j=1}^n H_{ij} \cdot \left[ U_{\vec{z}_1}^\eta \dots U_{\vec{z}_n}^\eta \right],$$

### JIMWLK Hamiltonian


$$H_{ij} = \frac{\alpha_s}{2\pi^2} \int d\vec{z}_k \frac{\vec{z}_{ik} \cdot \vec{z}_{kj}}{\vec{z}_{ik}^2 \vec{z}_{kj}^2} \left[ T_{i,L}^a T_{j,L}^a + T_{i,R}^a T_{j,R}^a - U_{\vec{z}_k}^{ab} (T_{i,L}^a T_{j,R}^b + T_{j,L}^a T_{i,R}^b) \right]$$

Known at NLO accuracy

# Known NLO impact factors

Very few NLO **CGC** impact factors are known

- $\gamma^* \rightarrow \gamma^*$  [Balitsky, Chirilli; Beuf]?
- Single inclusive particle production [Chirilli, Xiao, Yuan]
- Exclusive diffractive electro- and photo-production of a forward dijet [RB, Grabovsky, Szymanowski, Wallon]
- $\gamma_{L,T}^{(*)} \rightarrow V_L$  [RB, Grabovsky, Ivanov, Szymanowski, Wallon]




# Assumptions

- Regge-Gribov limit :  $s \gg Q^2 \gg \Lambda_{QCD}$
- Otherwise **completely general kinematics**
- **Shockwave (CGC) Wilson line approach**
- **Transverse dimensional regularization  $d = 2 + 2\epsilon$ , longitudinal cutoff**

$$|p_g^+| > \alpha p_\gamma^+$$

## LO diagram



Color factor

$$\frac{\delta^{ik}}{\sqrt{N_c}} [(\tilde{U}_{\vec{p}_1}^\alpha)_{ij} (\tilde{U}_{-\vec{p}_2}^\alpha)_{jk} - \delta_{ij} \delta_{jk}]$$

$$= \sqrt{N_c} \tilde{U}^\alpha(\vec{p}_1, \vec{p}_2)$$

$$\mathcal{A} = \frac{\delta^{ik}}{\sqrt{N_c}} \int d^D z_0 [\bar{u}(p_q, z_0)]_{ij} (-ie_q) \hat{\varepsilon}_\gamma e^{-i(p_\gamma \cdot z_0)} [v(p_{\bar{q}}, z_0)]_{jk} \theta(-z_0^+)$$

Impact factor


$$\mathcal{A}_{LO} \propto \delta(p_q^+ + p_{\bar{q}}^+ - p_\gamma^+) \int d^d \vec{p}_1 d^d \vec{p}_2 \delta(\vec{p}_q + \vec{p}_{\bar{q}} - \vec{p}_\gamma - \vec{p}_1 - \vec{p}_2) \Phi_{v1}(\vec{p}_1, \vec{p}_2)$$

$$\times \langle P' | \tilde{U}^\alpha(\vec{p}_1, \vec{p}_2) | P \rangle$$

$$\tilde{U}^\alpha(\vec{p}_1, \vec{p}_2) = \int d^d \vec{z}_1 d^d \vec{z}_2 e^{-i(\vec{p}_1 \cdot \vec{z}_1) - i(\vec{p}_2 \cdot \vec{z}_2)} \left[ \frac{1}{N_c} \text{Tr}(U_{\vec{z}_1}^\alpha U_{\vec{z}_2}^{\alpha\dagger}) - 1 \right]$$






# NLO open $q\bar{q}$ production



Diagrams contributing to the NLO correction

# First kind of virtual corrections





Color factor

$$\frac{C_F}{\sqrt{N_c}} \tilde{U}^\alpha(\vec{p}_1, \vec{p}_2)$$

Impact factor

$$A_{NLO}^{(1)} \propto \delta(p_q^+ + p_{\bar{q}} - p_\gamma^+) \int d^d \vec{p}_1 d^d \vec{p}_2 \delta(\vec{p}_q + \vec{p}_{\bar{q}} - \vec{p}_\gamma - \vec{p}_1 - \vec{p}_2) \Phi_{V1}(\vec{p}_1, \vec{p}_2) \\ \times C_F \langle P' | \tilde{U}^\alpha(\vec{p}_1, \vec{p}_2) | P \rangle$$

## Second kind of virtual corrections




Color factor

$$\frac{\delta^{ik}}{\sqrt{N_c}} (t^a U_1 t^b U_2^\dagger)_{ik} (U_3)^{ab}$$

$$C_F \mathcal{U}_{12} + \frac{1}{2} [\mathcal{U}_{13} + \mathcal{U}_{32} - \mathcal{U}_{12} + \mathcal{U}_{13} \mathcal{U}_{32}] = C_F \mathcal{U}_{12} + \mathcal{W}_{123}$$

Impact factor

$$\begin{aligned} \mathcal{A}_{NLO}^{(2)} &\propto \delta(p_q^+ + p_{\bar{q}} - p_\gamma^+) \int d^d \vec{p}_1 d^d \vec{p}_2 d^d \vec{p}_3 \delta(\vec{p}_q + \vec{p}_{\bar{q}} - \vec{p}_\gamma - \vec{p}_1 - \vec{p}_2 - \vec{p}_3) \\ &\times [\Phi'_{V1}(\vec{p}_1, \vec{p}_2) C_F \langle P' | \tilde{\mathcal{U}}^\alpha(\vec{p}_1, \vec{p}_2) | P \rangle \\ &+ \Phi_{V2}(\vec{p}_1, \vec{p}_2, \vec{p}_3) \langle P' | \tilde{\mathcal{W}}(\vec{p}_1, \vec{p}_2, \vec{p}_3) | P \rangle] \end{aligned}$$

LO open  $q\bar{q}g$  production

$$\mathcal{A}_R^{(2)} \propto \delta(p_q^+ + p_{\bar{q}}^+ + p_g^+ - p_\gamma^+) \int d^d \vec{p}_1 d^d \vec{p}_2 d^d \vec{p}_3 \delta(\vec{p}_q + \vec{p}_{\bar{q}} + \vec{p}_g - \vec{p}_\gamma - \vec{p}_1 - \vec{p}_2 - \vec{p}_3) \\ \times [\Phi'_{R1}(\vec{p}_1, \vec{p}_2) C_F \langle P' | \tilde{U}^\alpha(\vec{p}_1, \vec{p}_2) | P \rangle \\ + \Phi_{R2}(\vec{p}_1, \vec{p}_2, \vec{p}_3) \langle P' | \tilde{W}(\vec{p}_1, \vec{p}_2, \vec{p}_3) | P \rangle]$$



$$\mathcal{A}_R^{(1)} \propto \delta(p_q^+ + p_{\bar{q}}^+ + p_g^+ - p_\gamma^+) \int d^d \vec{p}_1 d^d \vec{p}_2 \delta(\vec{p}_q + \vec{p}_{\bar{q}} + \vec{p}_g - \vec{p}_\gamma - \vec{p}_1 - \vec{p}_2) \\ \times \Phi_{R1}(\vec{p}_1, \vec{p}_2) C_F \langle P' | \tilde{U}^\alpha(\vec{p}_1, \vec{p}_2) | P \rangle$$

# Divergences


## Divergences

- Rapidity divergence  $p_g^+ \rightarrow 0$   $\Phi_{V2}\Phi_0^* + \Phi_0\Phi_{V2}^*$
- UV divergence  $\vec{p}_g^2 \rightarrow +\infty$   $\Phi_{V1}\Phi_0^* + \Phi_0\Phi_{V1}^*$
- Soft divergence  $p_g \rightarrow 0$   $\Phi_{V1}\Phi_0^* + \Phi_0\Phi_{V1}^*, \Phi_{R1}\Phi_{R1}^*$
- Collinear divergence  $p_g \propto p_q$  or  $p_{\bar{q}}$   $\Phi_{R1}\Phi_{R1}^*$
- Soft and collinear divergence  $p_g = \frac{p_g^+}{p_q^+} p_q$  or  $\frac{p_g^+}{p_{\bar{q}}^+} p_{\bar{q}}, p_g^+ \rightarrow 0$   $\Phi_{R1}\Phi_{R1}^*$

# Rapidity divergence



Double dipole virtual correction  $\Phi_{V2}$



**B-JIMWLK evolution** of the LO term :  $\Phi_0 \otimes \mathcal{K}_{BK}$

# Rapidity divergence

## B-JIMWLK equation

$$\frac{\partial \tilde{U}_{12}^\alpha}{\partial \log \alpha} = 2\alpha_s N_c \mu^{2-d} \int \frac{d^d \vec{k}_1 d^d \vec{k}_2 d^d \vec{k}_3}{(2\pi)^{2d}} \delta(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 - \vec{p}_1 - \vec{p}_2) \left( \tilde{U}_{13}^\alpha \tilde{U}_{32}^\alpha + \tilde{U}_{13}^\alpha + \tilde{U}_{32}^\alpha - \tilde{U}_{12}^\alpha \right) \\ \times \left[ 2 \frac{(\vec{k}_1 - \vec{p}_1) \cdot (\vec{k}_2 - \vec{p}_2)}{(\vec{k}_1 - \vec{p}_1)^2 (\vec{k}_2 - \vec{p}_2)^2} + \frac{\pi^{\frac{d}{2}} \Gamma(1 - \frac{d}{2}) \Gamma^2(\frac{d}{2})}{\Gamma(d-1)} \left( \frac{\delta(\vec{k}_2 - \vec{p}_2)}{[(\vec{k}_1 - \vec{p}_1)^2]^{1-\frac{d}{2}}} + \frac{\delta(\vec{k}_1 - \vec{p}_1)}{[(\vec{k}_2 - \vec{p}_2)^2]^{1-\frac{d}{2}}} \right) \right]$$

$\eta$  **rapidity divide**, which separates the upper and the lower impact factors

$$\tilde{U}_{12}^\alpha \Phi_0 \rightarrow \Phi_0 \tilde{U}_{12}^\eta + 2 \log \left( \frac{e^\eta}{\alpha} \right) \mathcal{K}_{BK} \Phi_0 \tilde{W}_{123}$$

# Rapidity divergence

Virtual contribution

$$(\Phi_{V2}^\mu)_{div} \propto \Phi_0^\mu \left\{ 4 \ln \left( \frac{x\bar{x}}{\alpha^2} \right) \left[ \frac{1}{\epsilon} + \ln \left( \frac{\vec{p}_3^2}{\mu^2} \right) \right] - \frac{6}{\epsilon} \right\}$$

BK contribution

$$(\Phi_{BK}^\mu)_{div} \propto \Phi_0^\mu \left\{ 4 \ln \left( \frac{\alpha^2}{e^{2\eta}} \right) \left[ \frac{1}{\epsilon} + \ln \left( \frac{\vec{p}_3^2}{\mu^2} \right) \right] \right\}$$

Sum : the  $\alpha$  dependence cancels

$$(\Phi'_{V2}^\mu)_{div} \propto \Phi_0^\mu \left\{ 4 \ln \left( \frac{x\bar{x}}{e^{2\eta}} \right) \left[ \frac{1}{\epsilon} + \ln \left( \frac{\vec{p}_3^2}{\mu^2} \right) \right] - \frac{6}{\epsilon} \right\}$$



# Rapidity divergence

Cancellation of the remaining  $1/\epsilon$  divergence

Convolution

$$\begin{aligned}
 (\Phi'_{V_2}{}^\mu \otimes \mathcal{W}) &= 2 \int d^d \vec{p}_1 d^d \vec{p}_2 d^d \vec{p}_3 \left\{ 4 \ln \left( \frac{x\bar{x}}{e^{2\eta}} \right) \left[ \frac{1}{\epsilon} + \ln \left( \frac{\vec{p}_3^2}{\mu^2} \right) \right] - \frac{6}{\epsilon} \right\} \\
 &\times \delta(\vec{p}_{q1} + \vec{p}_{\bar{q}2} - \vec{p}_3) \left[ \tilde{\mathcal{U}}_{13} + \tilde{\mathcal{U}}_{32} - \tilde{\mathcal{U}}_{12} - \tilde{\mathcal{U}}_{13}\tilde{\mathcal{U}}_{32} \right] \Phi_0^\mu(\vec{p}_1, \vec{p}_2)
 \end{aligned}$$

Rq :

- $\Phi_0(\vec{p}_1, \vec{p}_2)$  only depends on one of the  $t$ -channel momenta.
- The double-dipole operators **cancel** when  $\vec{z}_3 = \vec{z}_1$  or  $\vec{z}_3 = \vec{z}_2$ .

This permits one to show that the convolution **cancel the remaining  $\frac{1}{\epsilon}$  divergence**.

Then  $\tilde{\mathcal{U}}_{12}^\alpha \Phi_0 + \Phi_{V_2}$  is **finite**

# Divergences

- Rapidity divergence

- UV divergence  $\vec{p}_g^2 \rightarrow +\infty$

$$\Phi_{V1} \Phi_0^* + \Phi_0 \Phi_{V1}^*$$

- Soft divergence  $p_g \rightarrow 0$

$$\Phi_{V1} \Phi_0^* + \Phi_0 \Phi_{V1}^*, \Phi_{R1} \Phi_{R1}^*$$

- Collinear divergence  $p_g \propto p_q$  or  $p_{\bar{q}}$


$$\Phi_{R1} \Phi_{R1}^*$$

- Soft and collinear divergence  $p_g = \frac{p_g^+}{p_q^+} p_q$  or  $\frac{p_g^+}{p_{\bar{q}}^+} p_{\bar{q}}$ ,  $p_g^+ \rightarrow 0$

$$\Phi_{R1} \Phi_{R1}^*$$

## UV divergence

## Tadpole diagrams



Some null diagrams just contribute to turning **UV divergences** into **IR divergences**

$$\Phi \propto \int \frac{d^D k}{(k^2 + i0)^2} \propto \left( \frac{1}{2\epsilon_{IR}} - \frac{1}{2\epsilon_{UV}} \right)$$

# Divergences

- Rapidity divergence

- UV divergence

- Soft divergence  $p_g \rightarrow 0$

$$\Phi_{V1} \Phi_0^* + \Phi_0 \Phi_{V1}^*, \Phi_{R1} \Phi_{R1}^*$$

- Collinear divergence  $p_g \propto p_q$  or  $p_{\bar{q}}$

$$\Phi_{R1} \Phi_{R1}^*$$

- Soft and collinear divergence  $p_g = \frac{p_g^+}{p_q^+} p_q$  or  $\frac{p_g^+}{p_{\bar{q}}^+} p_{\bar{q}}$ ,  $p_g^+ \rightarrow 0$

$$\Phi_{R1} \Phi_{R1}^*$$

## Constructing a finite cross section

Exclusive diffractive production of a forward dijet

From partons to jets


# Soft and collinear divergence

## Jet cone algorithm

We define a **cone** width for each pair of particles with momenta  $p_i$  and  $p_k$ , rapidity difference  $\Delta Y_{ik}$  and relative azimuthal angle  $\Delta\varphi_{ik}$


$$(\Delta Y_{ik})^2 + (\Delta\varphi_{ik})^2 = R_{ik}^2$$

If  $R_{ik}^2 < R^2$ , then the two particles together define a **single jet** of momentum  $p_i + p_k$ .



Applying this in the small  $R^2$  limit cancels our **soft and collinear** divergence.

# Remaining divergence



- Soft divergence  $p_g \rightarrow 0$

$$\Phi_{V1} \Phi_0^* + \Phi_0 \Phi_{V1}^*, \Phi_{R1} \Phi_{R1}^*$$

- Collinear divergence  $p_g \propto p_q$  or  $p_{\bar{q}}$

$$\Phi_{V1} \Phi_0^* + \Phi_0 \Phi_{V1}^* + \Phi_{R1} \Phi_{R1}^*$$

# Remaining divergence

## Soft real emission

$$(\Phi_{R1} \Phi_{R1}^*)_{soft} \propto (\Phi_0 \Phi_0^*) \int_{\text{outside the cones}} \left| \frac{p_q^\mu}{(p_q \cdot p_g)} - \frac{p_{\bar{q}}^\mu}{(p_{\bar{q}} \cdot p_g)} \right|^2 \frac{dp_g^+}{p_g^+} \frac{d^d p_g}{(2\pi)^d}$$

## Collinear real emission

$$(\Phi_{R1} \Phi_{R1}^*)_{col} \propto (\Phi_0 \Phi_0^*) (\mathcal{N}_q + \mathcal{N}_{\bar{q}})$$

Where  $\mathcal{N}$  is the number of jets in the quark or the antiquark

$$\mathcal{N}_k = \frac{(4\pi)^{\frac{d}{2}}}{\Gamma(2 - \frac{d}{2})} \int_{\alpha p_\gamma^+}^{p_{jet}^+} \frac{dp_g^+ dp_k^+}{2p_g^+ 2p_k^+} \int_{\text{in cone } k} \frac{d^d \vec{p}_g d^d \vec{p}_k}{(2\pi)^d \mu^{d-2}} \frac{\text{Tr}(\hat{p}_k \gamma^\mu \hat{p}_{jet} \gamma^\nu) d_{\mu\nu}(p_g)}{2p_{jet}^+ (p_k^- + p_g^- - p_{jet}^-)^2}$$

Those two contributions **cancel exactly the virtual divergences**



## Cancellation of divergences

## Total divergence

$$(d\sigma_1)_{div} = \alpha_s \frac{\Gamma(1-\epsilon)}{(4\pi)^{1+\epsilon}} \left( \frac{N_c^2 - 1}{2N_c} \right) (S_V + S_V^* + S_R + \mathcal{N}_{jet1} + \mathcal{N}_{jet2}) d\sigma_0$$

## Virtual contribution

$$S_V = \left[ 2 \ln \left( \frac{x_j x_{\bar{j}}}{\alpha^2} \right) - 3 \right] \left[ \ln \left( \frac{x_j x_{\bar{j}} \mu^2}{(x_j \vec{p}_j - x_{\bar{j}} \vec{p}_{\bar{j}})^2} \right) - \frac{1}{\epsilon} \right]$$

$$+ 2i\pi \ln \left( \frac{x_j x_{\bar{j}}}{\alpha^2} \right) + \ln^2 \left( \frac{x_j x_{\bar{j}}}{\alpha^2} \right) - \frac{\pi^2}{3} + 6$$

## Real contribution

$$S_R + \mathcal{N}_{jet1} + \mathcal{N}_{jet2} = 2 \left[ \ln \left( \frac{(x_{\bar{j}} \vec{p}_j - x_j \vec{p}_{\bar{j}})^4}{x_{\bar{j}}^2 x_j^2 R^4 \vec{p}_j^2 \vec{p}_{\bar{j}}^2} \right) \ln \left( \frac{4E^2}{x_{\bar{j}} x_j (\rho_{\gamma}^+)^2} \right) \right.$$

$$+ 2 \ln \left( \frac{x_{\bar{j}} x_j}{\alpha^2} \right) \left( \frac{1}{\epsilon} - \ln \left( \frac{x_{\bar{j}} x_j \mu^2}{(x_{\bar{j}} \vec{p}_j - x_j \vec{p}_{\bar{j}})^2} \right) \right) - \ln^2 \left( \frac{x_{\bar{j}} x_j}{\alpha^2} \right)$$

$$\left. + \frac{3}{2} \ln \left( \frac{16\mu^4}{R^4 \vec{p}_j^2 \vec{p}_{\bar{j}}^2} \right) - \ln \left( \frac{x_j}{x_{\bar{j}}} \right) \ln \left( \frac{x_j \vec{p}_j^2}{x_{\bar{j}} \vec{p}_{\bar{j}}^2} \right) - \frac{3}{\epsilon} - \frac{2\pi^2}{3} + 7 \right]$$

# Cancellation of divergences

## Total "divergence"


$$\begin{aligned}
 div &= S_V + S_V^* + S_R + \mathcal{N}_{jet1} + \mathcal{N}_{jet2} \\
 &= 4 \left[ \frac{1}{2} \ln \left( \frac{(x_{\bar{j}} \vec{p}_{\bar{j}} - x_j \vec{p}_{\bar{j}})^4}{x_{\bar{j}}^2 x_j^2 R^4 \vec{p}_{\bar{j}}^2 \vec{p}_j^2} \right) \left( \ln \left( \frac{4E^2}{x_{\bar{j}} x_j (\rho_{\gamma}^+)^2} \right) + \frac{3}{2} \right) \right. \\
 &\quad \left. + \ln(8) - \frac{1}{2} \ln \left( \frac{x_j}{x_{\bar{j}}} \right) \ln \left( \frac{x_j \vec{p}_{\bar{j}}^2}{x_{\bar{j}} \vec{p}_j^2} \right) + \frac{13 - \pi^2}{2} \right]
 \end{aligned}$$

Our cross section is thus **finite**

## Constructing a finite amplitude

Exclusive diffractive production of a light neutral vector meson

## Additional factorization




Once the amplitude is factorized in terms of **impact factors**, we perform an additional **twist expansion** in powers of a hard Björken scale (photon virtuality, Madelstam  $t..$ ).

Then we can factorize, in terms of **collinear factorization**, the bilocal matrix element

$$\langle V(p) | \bar{\psi}(z_{12}) \gamma^\mu \psi(0) | 0 \rangle_{z_{12}^2 \rightarrow 0} = p_\mu f_V \int_0^1 dx e^{ix(p \cdot z_{12})} \varphi_{\parallel}(x)$$

$\phi_{\parallel}(x) =$  meson **Distribution Amplitude (DA)**

# Exclusive diffractive production of a light neutral vector meson



$$\begin{aligned}
 \mathcal{A}_0 &= -\frac{e_V f_V \varepsilon_\beta}{N_c} \int_0^1 dx \varphi_{\parallel}(x) \int \frac{d^d \vec{p}_1}{(2\pi)^d} \frac{d^d \vec{p}_2}{(2\pi)^d} \\
 &\times (2\pi)^{d+1} \delta(p_V^+ - p_\gamma^+) \delta(\vec{p}_V - \vec{p}_\gamma - \vec{p}_1 - \vec{p}_2) \\
 &\times \Phi_0^\beta(x, \vec{p}_1, \vec{p}_2) \tilde{U}_{12}^n.
 \end{aligned}$$

Leading twist for a longitudinally polarized meson

Otherwise **general kinematics**, including transverse virtual photon (twist 3) contributions, and the photoproduction limit (for large  $t$ -channel momentum transfer)

## ERBL evolution equation

Efremov, Radyushkin, Brodsky, Lepage evolution equation for a DA

Renormalization of the bilocal operator

$$\bar{\psi}(z_{12})\gamma^\mu\psi(0)$$

⇒ Evolution equation for the distribution amplitude in the  $\overline{MS}$  scheme

$$\frac{\partial\varphi(x,\mu_F^2)}{\partial\ln\mu_F^2} = \frac{\alpha_s C_F}{2\pi} \frac{\Gamma(1-\epsilon)}{(4\pi)^\epsilon} \left(\frac{\mu_F^2}{\mu^2}\right)^\epsilon \int_0^1 dz\varphi(z,\mu_F^2)\mathcal{K}(x,z),$$

$\mathcal{K}$  = ERBL kernel

## ERBL evolution equation

Evolution equation for the distribution amplitude in the  $\overline{MS}$  scheme

$$\frac{\partial \varphi(x, \mu_F^2)}{\partial \ln \mu_F^2} = \frac{\alpha_s C_F}{2\pi} \frac{\Gamma(1-\epsilon)}{(4\pi)^\epsilon} \left(\frac{\mu_F^2}{\mu^2}\right)^\epsilon \int_0^1 dz \varphi(z, \mu_F^2) \mathcal{K}(x, z),$$

where we parameterize the **ERBL kernel** for consistency as

$$\begin{aligned} \mathcal{K}(x, z) &= \frac{x}{z} \left[ 1 + \frac{1}{z-x} \right] \theta(z-x-\alpha) \\ &+ \frac{1-x}{1-z} \left[ 1 + \frac{1}{x-z} \right] \theta(x-z-\alpha) \\ &+ \left[ \frac{3}{2} - \ln \left( \frac{x(1-x)}{\alpha^2} \right) \right] \delta(z-x). \end{aligned}$$

It is equivalent to the usual ERBL kernel

# Infrared finiteness

The amplitude we obtain is finite. For example the dipole  $\gamma_L^* \rightarrow V_L$  contribution reads

$$\begin{aligned}
 \Phi_1^+(x) &= \int_0^x dz \left( \frac{x-z}{x} \right) \Phi_0^+(x-z) \\
 &\times \left[ 1 + \left( 1 + \left[ \frac{1}{z} \right]_+ \right) \ln \left( \frac{\left( ((\bar{x}+z)\vec{p}_1 - (x-z)\vec{p}_2)^2 + (x-z)(\bar{x}+z)Q^2 \right)^2}{\mu_F^2(x-z)(\bar{x}+z)Q^2} \right) \right] \\
 &+ \frac{1}{2} \Phi_0^+(x) \left[ \frac{1}{2} \ln^2 \left( \frac{\bar{x}}{x} \right) + 3 - \frac{\pi^2}{6} - \frac{3}{2} \ln \left( \frac{\left( (\bar{x}\vec{p}_1 - x\vec{p}_2)^2 + x\bar{x}Q^2 \right)^2}{x\bar{x}\mu_F^2Q^2} \right) \right] \\
 &+ \frac{(\rho_\gamma^+)^2}{2x\bar{x}} \int_0^x dz \left[ (\phi_5)_{LL} |_{\vec{p}_3=\vec{0}} + (\phi_6)_{LL} |_{\vec{p}_3=\vec{0}} \right]_+ + (x \leftrightarrow \bar{x}, \vec{p}_1 \leftrightarrow \vec{p}_2).
 \end{aligned}$$


**No end point singularity**, even for a transverse photon and even in the photoproduction limit.



## Practical use of such results for phenomenology

## Practical use of such results

- Compute the upper impact factor using the effective Feynman rules ( $\sim$  BFKL gluon exchange!)
- Build non-perturbative models for the matrix elements of the Wilson line operators acting on the target states
- Solve the B-JIMWLK evolution for these matrix elements with such non-perturbative initial conditions at a typical target rapidity  $\eta = Y_0$
- Evaluate the solution at a typical projectile rapidity  $\eta = Y$
- Convolute the solution and the impact factor



$$\mathcal{A} = \int d\vec{z}_1 \dots d\vec{z}_n \Phi(\vec{z}_1, \dots, \vec{z}_n) \times \langle P' | U_{\vec{z}_1} \dots U_{\vec{z}_n} | P \rangle$$

## Residual parameter dependence

### Required parameters

- Renormalization scale  $\mu_R$
- Factorization scale  $\mu_F$  in the case of meson production (if assumed that  $\mu_F \neq \mu_R$ )
- Typical target rapidity  $Y_0$
- Typical projectile rapidity  $Y$


In the linear BFKL limit, the cross section only depends on  $Y - Y_0$ , so one only needs one arbitrary parameter  $s_0$  defined by

$$Y - Y_0 = \ln \left( \frac{s}{s_0} \right).$$

Modifying any of these parameter results in a higher order (NNLO) contribution

# General amplitude


- Most general kinematics
- The hard scale can be  $Q^2$ ,  $t$ ,  $M_X^2$ ...
- The target can be either a **proton** or an **ion**, or another impact factor.
- **Finite results for  $Q^2 = 0$**
- One can study **ultraperipheral collision** by tagging the particle which emitted the photon, in the limit  $Q^2 \rightarrow 0$ .



The general amplitude

# Phenomenological applications : exclusive dijet production at NLO accuracy


- HERA data for exclusive dijet production in diffractive DIS can be fitted with our results
- We can also give predictions for the same process in a future electron-ion or electron-proton collider (EIC, LHeC...)
- For  $Q^2 = 0$  we can give predictions for ultraperipheral  $pp$  and  $pA$  collisions at the LHC



Amplitude for diffractive dijet production

# Phenomenological applications : exclusive trijet production at LO accuracy

- HERA data for exclusive trijet production in diffractive DIS can be fitted with our results
- We can also give predictions for the same process in a future electron-ion collider
- For  $Q^2 = 0$  we can give predictions for ultraperipheral collisions at the LHC




Amplitude for diffractive trijet production

[Ayala, Hentschinski, Jalilian-Marian, Tejeda-Yeomans]

# Phenomenological applications

- Most general kinematics
- The hard scale can be  $Q^2$  or  $t$ .
- The target can be either a proton or an ion, or another impact factor.
- Finite results for  $Q^2 = 0$
- One can study ultraperipheral collision by tagging the particle which emitted the photon, in the limit  $Q^2 \rightarrow 0$ .



Amplitude for diffractive  $V$  production

## Conclusion

- We provided the **full computation** of the impact factor for the exclusive diffractive production of a forward dijet and of a light neutral vector meson with **NLO accuracy** in the **shockwave approach**
- It leads to an enormous number of possible phenomenological applications to test QCD in its Regge limit and towards saturation **in past, present and future  $ep$ ,  $eA$ ,  $pp$  and  $pA$  colliders**
- The linear limit of our result would provide interesting insight on the **linearized CGC/BFKL equivalence** at NLO accuracy