


Phenomenology 2017 Symposium



Contribution ID: 228

Type: parallel talk

Higgs to light jets decay at the LHC

Monday, 8 May 2017 14:45 (15 minutes)

We study the Higgs boson (h) decay to two light jets at the 14 TeV High-Luminosity-LHC (HL-LHC), where a light jet (j) represents any non-flavor tagged jet from the observational point of view.

% On the theory side, j is a gluon as expected in the Standard Model}.

The decay mode $h \rightarrow gg$ is chosen as the benchmark since it is the dominant channel in the Standard Model (SM), but the bound obtained is also applicable to the light quarks ($j = u, d, s$). We estimate the achievable bounds on the decay branching fractions through the associated production Vh ($V = W^\pm, Z$). Events of the Higgs boson decaying into heavy (tagged) or light (un-tagged) jets are correlatively analyzed.

%

We find that with 3000 fb^{-1} data at the HL-LHC, we should expect approximately 1σ statistical significance on the SM $Vh(gg)$ signal in this channel. This corresponds to a reachable upper bound $\text{BR}(h \rightarrow jj) \leq 4 \text{ BR}^{SM}(h \rightarrow gg)$ at 95% confidence level.

A consistency fit also leads to an upper bound

%

$\text{BR}(h \rightarrow cc) < 15 \text{ BR}^{SM}(h \rightarrow cc)$ at 95% confidence level.

%

The estimated bound may be further strengthened by adopting multiple variable analyses, or adding other production channels.

Summary

Primary authors: Ms HENDRICKS, Khalida (Ohio State University); CARPENTER, Linda (Ohio State University); ZHOU, Ning (Tsinghua University (CN)); HAN, Tao (University of Pittsburgh); QIAN, Zhuoni (University of Pittsburgh)

Presenter: QIAN, Zhuoni (University of Pittsburgh)

Session Classification: Higgs