

# Simplified model for Dark Matter

---

BAKUL AGARWAL

MICHIGAN STATE UNIVERSITY

In collaboration with Joshua Isaacson(MSU), Kirtimaan Mohan(MSU), Tim Tait(UC Irvine), Jan Winter(MSU)  
& C.-P. Yuan(MSU)

# Customary Dark Matter slide



# Motivation

---

- Too many dark matter models
- UV Complete theories have many free parameters
- Easier to constrain and make predictions



# Outline

---

1. The Model
2. Wilson Coefficients
3. Renormalisation Group Evolution
4. Preliminary results from Direct detection constraints

# The Simplified Model

---

$$\bullet \mathcal{L} = i\bar{\chi}\not{\partial}\chi + (D_\mu\tilde{\phi}_i)^*(D^\mu\tilde{\phi}_i) - M_\phi^2\tilde{\phi}_i^*\tilde{\phi}_i + \mathcal{L}_{int}$$

Majorana Dark Matter

Coloured scalar(s)

$$\text{uR model : } \mathcal{L}_{int} = g_{DM}(\tilde{\phi}_i^* \bar{\chi} P_R q_i + h.c.) \quad q_i = \{u, c, t\}$$

$$\text{dR model : } \mathcal{L}_{int} = g_{DM}(\tilde{\phi}_i^* \bar{\chi} P_R q_i + h.c.) \quad q_i = \{d, s, b\}$$

$$\text{qL model : } \mathcal{L}_{int} = g_{DM}(\tilde{\phi}_i^* \bar{\chi} P_L q_i + h.c.) \quad q_i = \{u, d, s, c, b, t\}$$

$$D_\mu = \partial_\mu - ig_s T^a G_\mu^a$$

*Simplified models for Dark Matter interacting with quarks : <https://arxiv.org/pdf/1308.2679.pdf>*

# The Simplified Model

- $\mathcal{L} = i\bar{\chi}\not{\partial}\chi + (D_\mu\tilde{\phi}_i)^*(D^\mu\tilde{\phi}_i) - M_\phi^2\tilde{\phi}_i^*\tilde{\phi}_i + \mathcal{L}_{int}$
- Effective lagrangian for interaction with quarks and gluons :

$$\mathcal{L}_q^{eff} = f_q(\bar{\chi}\chi)(m_q\bar{q}q) + \frac{g1_q}{m_\chi}(\bar{\chi}i\partial^\mu\gamma^\nu\chi)(O_{\mu\nu}^q) + \frac{g2_q}{m_\chi}(\bar{\chi}i\partial^\mu i\partial^\nu\chi)(O_{\mu\nu}^q)$$

$$\mathcal{L}_g^{eff} = f_g(\bar{\chi}\chi)(G^{a,\mu\nu}G_{\mu\nu}^a) + \frac{g1_g}{m_\chi}(\bar{\chi}i\partial^\mu\gamma^\nu\chi)(O_{\mu\nu}^g) + \frac{g2_g}{m_\chi}(\bar{\chi}i\partial^\mu i\partial^\nu\chi)(O_{\mu\nu}^g)$$

Scalar operators

Twist-2 operators


$$O_{\mu\nu}^q = \frac{1}{2}\bar{q}i\left(D_\mu\gamma_\nu + D_\nu\gamma_\mu - \frac{1}{2}g_{\mu\nu}\not{\partial}\right)q$$

$$O_{\mu\nu}^g = G^{a,\rho}_{\mu}G^a_{\rho\nu} + \frac{1}{4}g_{\mu\nu}G^{a,\alpha\beta}G^a_{\alpha\beta}$$


# The Simplified Model

---


$\chi q \rightarrow \chi q$  :



$\chi g \rightarrow \chi g$  :



# Wilson Coefficients



In Momentum Space :

$$f_g (\bar{\chi} \chi) (G^{a, \mu\nu} G_{\mu\nu}^a) \longrightarrow 8i f_g (p_2^\mu k_2^\nu - g^{\mu\nu} p_2 \cdot k_2) \epsilon_\mu(p_2) \epsilon_\nu(k_2) (\bar{\chi} \chi)$$

$$\frac{g 2_G}{m_\chi^2} (\bar{\chi} i \partial^\mu \partial^\nu \chi) (O_{\mu\nu}^g) \longrightarrow g 2_G (\dots)^{\mu\nu} \epsilon_\mu(p_2) \epsilon_\nu(k_2) (\bar{\chi} \chi)$$

**Orthogonal**

Neutralino nucleon scattering reexamined : <https://journals.aps.org/prd/pdf/10.1103/PhysRevD.48.3483>

# RG Evolution

---

- Need effective coupling to nucleons for direct detection constraints

$$\mathcal{L}_N = f_N(\bar{N}N)(\bar{\chi}\chi)$$

- Nuclear matrix elements of  $m_q\bar{q}q$ ,  $G^{a,\mu\nu}G^a_{\mu\nu}$ , etc. are known
- Evolve the parton level Wilson coefficients to nuclear scale

- RG evolution through QCD running and matching at heavy quark thresholds

$$c_i(\mu_{low}) = R_{ij}(\mu_{low}, \mu_{high}) c_j(\mu_{high})$$



$$c'_i(\mu_Q) = M_{ij}(\mu_Q) c_j(\mu_Q)$$

- Corresponds to mixing of operators of same spin and integrating out heavy quarks
- Evaluate the Wilson coefficients at the high scale and evolve them to the nuclear scale

*Standard Model anatomy of WIMP dark matter direct detection*



*II: QCD analysis and hadronic matrix elements :- <https://arxiv.org/pdf/1409.8290.pdf>*

# RG Evolution





Effect of RGE on nuclear effective coupling for different models

# RG Evolution





Effect of RGE on individual operators for varying masses

# RG Evolution





Effect of RGE on individual operators for varying masses

# RG Evolution





Contribution of each operator with and without RGE

# Constraints from direct detection





SI constraints on  $q_L$  model from LUX - <https://journals.aps.org/prl/pdf/10.1103/PhysRevLett.118.021303>

# Constraints from direct detection



SI constraints on  $u_R$  model from LUX - <https://journals.aps.org/prl/pdf/10.1103/PhysRevLett.118.021303>

# Constraints from direct detection



SI constraints on dR model from LUX - <https://journals.aps.org/prl/pdf/10.1103/PhysRevLett.118.021303>

# Conclusions

---

- Study of simplified models useful for constraining BSM physics
- RGE extremely important for direct detection constraints
- Higher-dimensional operators can be relevant, even dominant
  
- Next: Collider constraints