



Pheno 2017 Symposium



JET OBSERVABLES AND STOPS AT 100 TEV COLLIDER

[JJI FAN, PRERIT JAISWAL, SHING CHAU LEUNG, ARXIV: 1704.03014]


John Shing Chau Leung
Brown University
May 9, 2017

MOTIVATION

Once LHC@14 TeV is done. What's next?

- Pin down the naturalness of SM.
- An exploratory machine. For generic physics involving high mass states, a large centre of mass energy provides the most direct access to new physics!

[Arkani-Hamed, Han, Mangano, Wang, 16]






OUR PAPER

- Studies the mass reach for three simplified SUSY models.
- Addresses challenges involving hyper-boosted SM objects from generic heavy states in 100 TeV collider
- Implements jet substructure techniques to improve the analysis sensitivity



SUSY MODELS

SIMPLIFIED MODELS



- Model 1 – LSP Bino (decay channel: $t + \text{MET}$)
- Model 2 – LSP Higgsino ($t + \text{MET} / b + \text{MET}$)
- Model 3 – LSP Bino, a gluino between LSP and stop ($3t + \text{MET}$)


LHC LIMITS

[CMS-PAS-SUS-16-051, CMS-PAS-SUS-16-036, CMS COLLABORATION]

$\sqrt{s} = 13$ TeV LHC constraint


- Stop ~ 1 TeV
- Weak LSP ~ 1 TeV
- Gluino ~ 2 TeV

Single lepton search



- In a $\sqrt{s} = 100$ TeV collider, more jets at higher energy.
- SM objects are boosted and collimated.
- Jet substructure techniques are needed!

MT2 search





JET SUBSTRUCTURE TOOLS:



MASS DROP x_{μ}

[THALER AND WANG, 08; REHERMANN AND TWEEDIE, 11]



A variable to distinguish between boosted leptonic top jet and QCD jets

$$X_u = 1 - \frac{m_{j_{t\bar{t}}}^2}{m_j^2}$$

which is the one minus ratio between jet mass without muons ($m_{j_{t\bar{t}}}$) and whole jet mass (m_j).



- for leptonic top jet 1
 - m_j = mass of top
 - $m_{j_{t\bar{t}}}$ = mass of b
- X_u for b-jet has a broad spectrum since $m_j \approx m_{j_{t\bar{t}}}$.






SUBSTRUCTURE TOOLS: N-SUBJETINESS τ_N

[J. THALER AND K. VAN TILBURG, 11]


How much does jet “looks like” from a N-body decay?

N-subjetiness (τ_N) \rightarrow how close are the jet constituents are from N sub-jet axes in the jet cone.




$$\tau_N^{(\beta)} = \frac{1}{R_0} \sum_i p_{T,i} \min \{ \Delta R_{1,i}^\beta, \Delta R_{2,i}^\beta, \dots, \Delta R_{N,i}^\beta \}$$

- When a jet looks like a 3-prong object, τ_3 will be small but τ_2 will be large.
e.g. a hadronic top jet ($t \rightarrow Wb \rightarrow qqb$)
- A spread-out QCD jet can have a small τ_N , but τ_{N-1} is small as well!
- τ_3/τ_2 is a good variable to distinguish a hadronic top jet from QCD jet.



QCD vs hadronic top






ANALYSIS STRATEGY AND RESULT




SEARCH STRATEGY STOP-BINO/HIGGSINO

- Two jets with $p_T > 1$ TeV;
- No isolated lepton with $p_T > 1$ TeV;
- MET > 3 TeV
- Muon-in-jet (> 200 GeV) requirement
 - high p_T muons are either from leptonic W decay or from B-meson decay.
 - muon is in jet due to the boosted t / b quarks.




STOP-HIGGSINO MASS REACH

The discovery/exclusion contours for stop-higgsino model with an integrated luminosity $L = 3 \text{ ab}^{-1}$.



(a) Discovery



(b) Exclusion

The solid lines are 5σ discovery contour (left) and exclusion at 95% C.L. (right). The dashed lines are the $\pm 1\sigma$ boundaries.

The stop-bino and stop-higgsino model signature differs in the b-jet signature.

We define the variable r_-


$$r_- = \frac{N_b - N_t}{N_b + N_t}$$

N_b = # of B-jet defined by $m_j < 120$ GeV and $x_u < 0.5$;


N_t = # of top-jet defined by $m_j > 120$ GeV or $x_u > 0.5$.

Model-distinguishability Reach:

$$\{m_{\tilde{\tau}} \sim 6 \text{ TeV}, m_{\tilde{\chi}} \sim 2 \text{ TeV}\}$$




r_- distribution




95% confident level

SEARCH STRATEGY STOP-GLUINO-LSP



Pair-production



Associated-production

- Kinematic cuts
 - 3 Top-tagging. Definition of top tagging for jets in events passing kinematic cuts
 - Leptonic top jets (jets with $p_T > 200$ GeV muon): $x_u > 0.7$
 - Hardronic top jets (jets without tagged muon): $0.1 < \tau_3/\tau_2 < 0.45$
- (jet mass window: $140 \text{ GeV} < m_j < 240 \text{ GeV}$)

STOP MASS REACH

$m_{\tilde{t}}$ (TeV)	S	B	σ
5.5	10.7	1.7	6.3
6.0	10.0	6.7	3.5

3σ can be reached for the benchmark point of 6 TeV stop and 2.75 TeV of gluino at $L = 30 \text{ ab}^{-1}$.

An increased L is needed due to high the high susy background.

GLUINO MASS REACH

Using the same set of jet substructure techniques:

A gluino mass reach to 11 TeV at $L = 3 \text{ ab}^{-1}$ (pair-production channel).

~2 TeV improvement compared to the same sign Di-lepton gluino search in [Cohen, et al., 14].

$m_{\tilde{g}}$ (TeV)	Top tags	S	B	σ
10.0	2	12.4	0.8	8.1
11.0	1	13.8	9.5	3.9

SUMMARY

- A 100 TeV collider will extend our understanding to the naturalness of Standard Model and explore new physics in general.
- We need jet substructure technique to effectively extract physics from a 100 TeV collider.
- Stop-Bino/Higgsino discovery mass reach at 7 TeV, model identification reach at 6 TeV ($L = 3 \text{ ab}^{-1}$)
- Stop-Gluino-LSP model mass reach 6 TeV ($L = 30 \text{ ab}^{-1}$). Gluino mass reach 11 TeV ($L = 3 \text{ ab}^{-1}$).



BACK-UP SLIDES

REASON FOR MASS SPECTRUM

Why do we have light stop?

- The need for cancelation of Higgs renormalization.
- Large renormalization and t_R - t_L mixing pull the t down to the foot of the scalar spectrum. [Pierce, Bagger, Matchev, Zhang, 97]

Why the Scalar-Gaugino mass splitting? [Arkani-Hamed, Dimopoulos, Giudice, Romanino, 04]

Suppose there is no gauge singlet in an underlying SUSY theory. SUSY breaking from a hidden sector is mediated by a messenger with a group index .

- Scalar mass is generated at tree-level in the Kähler potential

$$\int d^4\theta \frac{S_a^\dagger S_a}{M_*^2} \Phi^\dagger \Phi = m_{soft}^2 \Phi^\dagger \Phi$$

- Tree-level gaugino mass is forbidden by symmetry mass suppression!


← Uncontracted group index

~~$$\int d^2\theta \frac{S_a}{M_*^2} WW$$~~


Mass term

An extra factor of suppression


STOP-BINO MODEL



Final state: $t\bar{t}$ + MET




STOP-HIGGSINO MODEL



- \tilde{t} 50% decay to $t + \text{MET}$
- \tilde{t} 50% decay to $b + \text{MET}$

Final states: combinatoric of both



Stop

STOP-GLUINO-LSP MODEL

+t



Gluino

+2t

Bino

- Two production channels (stop-pair and stop-gluino associated)
- Final states: 6t + MET


Stop-gluino-bino Model




BOOSTED OBJECT CHALLENGES

[COHEN, D'AGNOLO, HANCE, LOU, AND WACKER, 14]

- At high luminosity, the high Standard Model event rate can conceal new physics.
- At high energy, SM objects are boosted. Different objects are collimated along the boost direction, to a point they are smaller than a calorimeter cell.
- New jet substructure techniques are needed.




A top quark decaying in the LHC





A boosted top in a 100 TeV collider

DETECTOR DESIGN




EFFICIENCY




Signal/background efficiencies for stop-higgsino cut

EFFICIENCY



(a)



(b)

QCD mistag vs signal efficiency Signal/background efficiencies top-tagging