Electroweak Splitting Functions and High Energy Showering

Junmou Chen

University of Pittsburgh

arXiv:1611.00788 J.C. T. Han B. Tweedie

May 9, 2017

Collisions in Colliders

Proton-proton scattering: sum of free quarks and gluons scatterings.



QCD Parton Shower and Electroweak Shower

- For Strong Interactions, when hard scattering scales $\sim$ 50 GeV

$$\frac{50\,GeV}{\Lambda_{QCD} \sim 500\,MeV} >> 1$$

  Parton shower emerges

- In a 100 TeV Collider, hard scattering could scale $\sim$ 10 TeV

$$\frac{10\,TeV}{m_W \sim 100\,GeV} \sim \frac{50\,GeV}{\Lambda_{QCD} \sim 500\,MeV} >> 1$$

- We are facing a new phenomenon: electroweak parton shower.

Novelties of Electroweak Showering

Novelties of electroweak showering:

- Spontaneous symmetry breaking:
  1. massive particle: perturbative shut-off
  2. additional degrees of freedom: longitudinal and higgs
- Chirality: Left-handed fermions and right-handed fermions couple differently.
- Neutral bosons interference: γ/Z_T; h/Z_L
- Weak isospin self-averaging.

Our focus today: VEV corrections, i.e. the effects of EW symmetry breaking.

Collinear Splitting Functions



$$E_B \simeq zE_A \quad E_C \simeq (1-z)E_A = \bar{z}E_A \quad k_T \simeq z\bar{z}E_A\theta$$

$$d\sigma_{X,B+C} \simeq d\sigma_{X,A} \times d\mathcal{P}_{A\to B+C}$$

Assumption: Interferences are small

Splitting functions serve as kernels of parton shower.

Splitting Functions for Unbroken $SU(2)_L \times U(1)_Y$

Splittings analogue to QCD



$$P_{f_L \to f_L + W_T/Z_T} \sim \alpha_W \frac{1+(1+z)^2}{z} \cdot \frac{1}{p_T^2}$$



$$P_{W_T \to W_T Z_T/\gamma} \sim \alpha_W \frac{(1-z\bar{z})^2}{z(1-z)} \cdot \frac{1}{p_T^2}$$



$$P_{W_T/Z_T \to f_L \bar{f}_R} \sim \alpha_W \frac{1}{p_T^2}$$

Splitting Functions for Unbroken $SU(2)_L \times U(1)_Y$

Splittings with higgs and longitudinal, with $W_L^\pm \sim \phi_L^\pm$ and $Z_L \sim \phi_0$



$P_{f_L \to f_R^{(\prime)} + h/W_L^\pm} \sim y^2_{f^{(\prime)}} \frac{1}{p_T^2}$
(helicity flipped)



$P_{W_L^\pm \to W_T^\pm h/Z_L} \sim \alpha_W \frac{1}{z} \frac{1}{p_T^2}$



$P_{h/W_L^\pm \to f_L \bar{f}_L^{(\prime)}} \sim y^2_{f^{(\prime)}} \frac{1}{p_T^2}$



$P_{V_T \to V_L(/h) V_L/h} \sim \alpha_W \frac{1}{p_T^2}$

Longitudinal in Broken(Real) Theory: Huge Intereference

Longitudinal vector boson $\epsilon_L \sim \frac{k^\mu}{m_W}$, bad energy behavior and huge interference!





Figure: higgs contribution $\propto \frac{m_t m_h}{v^2}$ fixes the problem.

Goldstone Equivalence Gauge

- Hint to the solution: $\epsilon_L \sim \frac{k^\mu}{m_W} \to \phi_W$ in high energy — goldstone equivalence theorem.
- Write $\epsilon_L^\mu(k) = \frac{k^\mu}{m_W} - \frac{m_W}{n\cdot k} n^\mu$, with $n^\mu = (-1, \hat{k})$. So impose gauge-fixing $\frac{1}{2\xi}(n^\mu W_\mu)^2$.
- Consequences:
  1. $n \cdot k \neq 0$: k^μ terms are eliminated – power counting is good
  2. $n \cdot n = 0$, $\epsilon_n \simeq \frac{m_W}{2E}$, VEV corrections to unbroken limit.
  3. $|W_L> = |W_n> + i|\phi>$ —- manifest goldstone equivalence theorem
  4. Broken theory continuously goes to unbroken theory when $v \to 0$



$\epsilon^\mu_{s=\pm,n}$ are "stripped " out of propagators, and inserted into vertices.

Systematic VEV(mass) corrections

"Scalarizing" k^μ/m_W enables systematic incorporation of all splittings (up to log) proportional to VEV(masses)

- Massless $f \to f + V_L$ corresponds to $f \to f\phi$ according to goldstone equivalence

$$y_f = 0 \to \quad \text{gives } 0$$

- Contribution comes from VEV correction $V_n \sim \frac{m_V}{E}$: "Effective W approximation".

$$\frac{d\mathcal{P}_{f\to f+V_L}}{dzdk_T^2} \sim \frac{\alpha_2}{2\pi}\frac{1-z}{z}\frac{m_V^2}{k_T^4} \tag{1}$$

- Has been known for years: vector boson fusion observed in LHC

  S. Dawson (1985); G. Kane et al. (1984); Chanowitz & Gailard (1984)

Ultra-collinear splitting

There are characteristically new channels in the broken phase:

$$\frac{v^2}{k_T^2}\frac{dk_T^2}{k_T^2} \sim (1-\frac{v^2}{Q^2})$$

"Ultra collinear":

$k_T^2 > m_W^2$, it shuts off;

$k_T^2 < m_W^2$, flattens out!

The DPFs for W_L thus don't run at leading log:

"Bjorken scaling" restored (higher-twist effects)!





New Broken Splittings: ultra-collinear behavior (11 new types)

A whole new set of broken splittings

Splitting in the Broken Gauge

New fermion splitting: $\frac{v^2}{k_T^2}\frac{dk_T^2}{k_T^2} \sim (1-\frac{v^2}{Q^2})$

V_L is of IR, h no IR

	$\frac{1}{16\pi^2}\frac{v^2}{\bar{k}_T^4}\left(\frac{1}{z}\right)$	$\frac{1}{16\pi^2}\frac{v^2}{\bar{k}_T^4}$	$\frac{1}{16\pi^2}\frac{v^2}{\bar{k}_T^4}$
	$\to V_L\, f_s^{(\prime)}$ $(V=\gamma)$	$h\ f_s$	$V_T\, f_{-s}^{(\prime)}$
$f_{s=L}$	$\left(I_f^V (y_f^2\bar{z} - y_{f^{(\prime)}}^2)z - Q_{f_L}^V g_V^2 \bar{z}\right)^2$	$\frac{1}{4}y_f^4 z(1+\bar{z})^2$	$g_V^2 z\left(Q_{f_R}^V y_f \bar{z} - Q_{f_L}^V y_{f^{(\prime)}}\right)^2$
$f_{s=R}$	$\left(I_f^V y_f y_{f^{(\prime)}} z^2 - Q_{f_R}^V g_V^2 \bar{z}\right)^2$	$\frac{1}{4}y_f^4 z(1+\bar{z})^2$	$g_V^2 z\left(Q_{f_L}^V y_f \bar{z} - Q_{f_R}^V y_{f^{(\prime)}}\right)^2$

Chirality flipping: $\sim m_f$

Chirality conserving:
Non-zero for massless f

New Broken Splittings: ultra-collinear behavior (11 new types)

A whole new set of broken splittings

Splitting in the Broken Gauge

New gauge boson splitting in 3-W_L

Vector boson V_L is of IR.



$$\frac{v^2}{k_T^2}\frac{dk_T^2}{k_T^2} \sim (1-\frac{v^2}{Q^2}$$

$\frac{1}{16\pi^2}\frac{v^2}{\tilde{k}_T^4}\left(\frac{1}{z\bar{z}}\right)$

	$\to W_L^+ W_L^-$	$Z_L W_L^\pm / Z_L$
$W_L^\pm$	0	$\frac{1}{16}g_2^4\left((\bar{z}-z)(2+z\bar{z})-t_W^2\bar{z}(1+\bar{z})\right)^2$
h	$\frac{1}{4}\left(g_2^2(1-z\bar{z})-\lambda_h z\bar{z}\right)^2$	$\frac{1}{8}\left(g_Z^2(1-z\bar{z})-\lambda_h z\bar{z}\right)^2$
Z_L	$\frac{1}{16}g_2^4\left((\bar{z}-z)(2+z\bar{z}-t_W^2 z\bar{z})\right)^2$	0
$[hZ_L]$	$\frac{i}{8}g_2^2\left(g_2^2(1-z\bar{z})-\lambda_h z\bar{z}\right)(\bar{z}-z)\left(2+z\bar{z}-t_W^2 z\bar{z}\right)$	0

h has no IR.



	$\frac{1}{16\pi^2}\frac{v^2}{\tilde{k}_T^4}\left(\frac{1}{\bar{z}}\right)$ $\to h\, W_L^\pm / Z_L$	$\frac{1}{16\pi^2}\frac{v^2}{\tilde{k}_T^4}$ $h\,h$
$W_L^\pm$	$\frac{1}{4}z\left(g_2^2(1-z\bar{z})+\lambda_h\bar{z}\right)^2$	0

Top decay & shower





Figure: a) Splitting corrections to "tail" of top decay; b) interplay of multiple elements of top showering: yukawa (ϕ) & gauge coupling (W_n); helicity conserving & flipping.

Summary

- With the increase of energy scale in colliders, we are facing a new phenomenon of EW shower.
- EW sector presents rich physics
- With the help of a new gauge (GEG), we are able to calculate all the EW (collinear log) splitting functions.
  1. Unbroken limit: analogue to QCD
  2. Broken: systematic VEV corrections – ultra-collinear splittings
- We incorporate EW showering into a Monte Carlo program
  1. More complete
  2. More accurate in implementation
- Looking forward to seeing them in real world!

Thank you!

New Broken Splittings: ultra-collinear behavior

A whole new set of broken splittings

Splitting in the Broken Gauge

New gauge boson splitting to $W_L\, W_T$

Vector boson V_L is of IR.



$$\frac{v^2}{k_T^2}\,\frac{dk_T^2}{k_T^2} \sim (1-\frac{v^2}{Q^2})$$

$\frac{1}{16\pi^2}\frac{v^2}{\tilde{k}_T^4}\left(\frac{1}{z}\right)$

	$\to W_L^\pm\,\gamma_T$	$W_L^\pm\,Z_T$	$Z_L\,W_T^\pm$	$W_L^+\,W_L^-$ or $W_L^-\,W_L^+$
$W_T^\pm$	$e^2 g_2^2 \bar{z}^3$	$\frac{1}{4}c_W^2 g_2^4 \bar{z}\left((1+\bar{z})+t_W^2 z\right)^2$	$\frac{1}{4}g_2^4\bar{z}(1+\bar{z})^2$	0
γ_T	0	0	0	$e^2 g_2^2 \bar{z}$
Z_T	0	0	0	$\frac{1}{4}c_W^2 g_2^4 \bar{z}\left((1+\bar{z})-t_W^2 z\right)^2$
$[\gamma Z]_T$	0	0	0	$\frac{1}{2}c_W e g_2^3 \bar{z}\left((1+\bar{z})-t_W^2 z\right)$

h & f have no IR.



	$\frac{1}{16\pi^2}\frac{v^2}{\tilde{k}_T^4}$ $\to h\,V_T\ (V\neq\gamma)$	$\frac{1}{16\pi^2}\frac{v^2}{\tilde{k}_T^4}$ $f_s\,\bar{f}_s^{(\prime)}$
V_T	$\frac{1}{4}z\bar{z}g_V^4$	$\frac{1}{2}g_V^2\left(Q_{f_s}^V y_{f^{(\prime)}} z + Q_{f_{-s}}^V y_f \bar{z}\right)^2$
$[\gamma Z]_T$	0	$\frac{1}{2}e g_Z y_f^2 Q_f^\gamma \left(Q_{f_s}^Z z + Q_{f_{-s}}^Z \bar{z}\right)$

Estimations of Splitting Rates at 1 TeV and 10 TeV

Process	$\approx \mathcal{P}(E)$ (leading-log term)	$\mathcal{P}(1\text{ TeV})$	$\mathcal{P}(10\text{ TeV})$
$q \to V_T q^{(\prime)}$ (CL+IR)	$(3\times10^{-3})\left[\log\frac{E}{m_W}\right]^2$	1.6%	7%
$q \to V_L q^{(\prime)}$ (UC+IR)	$(2\times10^{-3})\log\frac{E}{m_W}$	0.4%	1.1%
$t_R \to W_L^+ b_L$ (CL)	$(8\times10^{-3})\log\frac{E}{m_W}$	2.5%	4%
$t_R \to W_T^+ b_L$ (UC)	(6×10^{-3})	0.6%	0.6%
$V_T \to V_T V_T$ (CL+IR)	$(0.015)\left[\log\frac{E}{m_W}\right]^2$	7%	34%
$V_T \to V_L V_T$ (UC+IR)	$(0.014)\log\frac{E}{m_W}$	2.7%	7%
$V_T \to f\bar{f}$ (CL)	$(0.02)\log\frac{E}{m_W}$	5%	10%
$V_L \to V_T h$ (CL+IR)	$(2\times10^{-3})\left[\log\frac{E}{m_W}\right]^2$	0.8%	4%
$V_L \to V_L h$ (UC+IR)	$(2\times10^{-3})\log\frac{E}{m_W}$	0.5%	1%

- Double log $>$ single log.
- "Color" factor $C_A(adjoint) > C_F(fundamental)$: biggest from gauge-triple splitting

"Weak jets": EW Showering with Multiple Gauge Bosons



EW Showering with Multiple Gauge Bosons





Figure: $O(10\%)$ suppression for every additional boson radiation;
Back reaction & angular veto: more precisely (comparing with Pythia 8) taking care of phase space.

Neutral boson interference: γ/Z_T initiated splittings





Figure: Left: LH electron source, constructive interference;
Right: RH electron source, mainly B_0 intermediated, destructive. More transparent in B_0/W_0 basis.

Neutral Interference

- The splitting rate is

$$\frac{d\mathcal{P}}{dk_T^2 dz} = \frac{tr(\hat{\rho}\frac{d\hat{P}}{dk_T^2 dz})}{tr(\hat{\rho})} \tag{2}$$

  Both $\hat{\rho}$ and $\frac{d\hat{P}}{dk_T^2 dz}$ are 2×2 matrices.

- $\hat{\rho}$ is the production density matrix from hard scattering,

$$\rho_{ij} = M_i^*(hard)M_j(hard) \tag{3}$$

Collimated higgses



Figure: $h \to hh$, rare but interesting.

Implementation and New Phenomenon: EW PDF

- At high energy, EW particles "show up" in the initial beams, i.e. EW PDFs.
- A useful intermediate concept is luminosities, with which the total cross section could be written as,

$$\sigma_{PP}(V_1 V_2 \to X) = \int_{\tau_{\rm low}}^{\tau_{\rm high}} d\tau \, \frac{d\mathcal{L}_{V_1 V_2}}{d\tau} \, \hat{\sigma}(V_1 V_2 \to \hat{X}_\tau) \,, \qquad (4)$$

-

$$\frac{d\mathcal{L}_{V_1 V_2}}{d\tau} \simeq \frac{2}{(\delta_{V_1 V_2}+1)} \int_\tau^1 \frac{d\xi}{\xi} \int_{\tau/\xi}^1 \frac{dz_1}{z_1} \int_{\tau/\xi/z_1}^1 \frac{dz_2}{z_2} \times \sum_{q_1, q_2} f_{V_1 \in q_1}(z_1) f_{V_2 \in q_2}(z_2) \; f_{q_1 \in P}(\xi) f_{q_2 \in P}\left(\frac{\tau}{\xi z_1 z_2}\right) \qquad (5)$$


√s (TeV)
0.4 0.6 0.8 1 1.2 1.4 1.6 1.8
14 TeV
qq̄
qγ
qW±T
W±Tγ
W+TW-T
qW±L
W+LW-L
dL/dτ
0.05 0.1
√τ= √s/S


√s (TeV)
2 4 6 8 10 12
100 TeV
qq̄
qγ
qW±T
W±Tγ
W+TW-T
qW±L
W+LW-L
dL/dτ
0.05 0.1
√τ = √s/S

:

:


W+LW-L
W+TW-T
W±Tγ
qq̄
qγ
qW±T
qW±L
dL100TeV/ dL14TeV
√s (TeV)

More on neutral boson interference





Implementation and New Phenomenon: h/Z_L and higgs



Figure: (a) h/Z_L interference, usually maximal. More transparent in H_0/H_0^* basis (b) $h \to hh$, rare but interesting.

EW Showering Effects on New Physics



Figure: Showered events from 20 TeV W'^{+} decays. Invariant mass more sharply peaked for multiple splittings.

Isospin self-averaging & Combining QCD effects





Figure: Showered events from 20 TeV W'^{+} decays. (a) $W'^{+} \to t_L \bar{b}_R$, full EW shower, and (b) combining EW and QCD showering.