

Searches for New Resonances Decaying to Third Generation Quarks at CMS

Petar Maksimovic
Johns Hopkins

PHENO 2017

The layout of this talk


- Many BSM models predict such resonances
 - composite Higgs, warped extra dimensions, etc.
- Many results to cover, focus on recent results with top quark(s) in final state
- Channels with large top BR: $t \rightarrow b\nu\ell$ and $t \rightarrow b(q\bar{q}')$
- $Z' \rightarrow t\bar{t}$ (semileptonic + hadronic) B2G-16-015
- $W' \rightarrow tb$ (semileptonic + hadronic) B2G-17-010 B2G-16-019
- $Z' \rightarrow tT' \rightarrow t(bW)$ (hadronic) B2G-16-013

$Z' \rightarrow t\bar{t}$

- Three boosted channels:

- Dileptons


lepton
+ b jet



lepton
+ b jet

- Semileptonic


lepton
+ E_T^{miss} +
b jet



merged
into a
single
top jet

- All-hadronic

top jet



top jet



- Isolation specially handled

- Top tagging tools are by now quite mature

$Z' \rightarrow t\bar{t}$: top tagging variables



B2G-16-015

Semileptonic

N-subjettiness τ_{32} 

Top Jet Soft Drop mass [GeV]

Fully hadronic



Jet τ_{32} 

Jet soft-drop mass [GeV]

$Z' \rightarrow t\bar{t}$: signal shapes

- At high Z' masses, large widths result in very long tail
 - (may need to rethink strategy for higher masses)



B2G-16-015



$Z' \rightarrow t\bar{t}$: hadronic

- Divided according to # of subjet b-tags (0,1,2), and $\Delta y_{t\bar{t}}$
- Main bkg: QCD (not-top multijet), derived from data


B2G-16-015



$Z' \rightarrow t\bar{t}$: semileptonic



- Divided into e and μ , and according to # of b-tags and top tags
- Main bkg: $t\bar{t}$:
 - template from simulation floating in final fit

B2G-16-015




$Z' \rightarrow t\bar{t}$: limits

Narrow




Wider

Very Wide




B2G-16-015




RS gluon

$W' \rightarrow tb$

- Semileptonic channel



- All-hadronic channel




- Similar to $Z' \rightarrow t\bar{t}$, except only one boosted top

- trickier (larger QCD background)



- Hadronic: b-tagged jet + top-tagged jet

- QCD est. from data



$W' \rightarrow tb$

- Hadronic:
 - limits on 2015 data similar to Run 1




- Semileptonic:
 - full 2016 dataset
 - improved analysis strategy
 - limits at ~ 3.3 -3.5 TeV !

W' – generalized couplings

B2G-17-010

$$\mathcal{L} = \frac{V_{q_i q_j}}{2\sqrt{2}} g_w \bar{q}_i \gamma_\mu \left(a_{q_i q_j}^R (1 + \gamma^5) + a_{q_i q_j}^L (1 - \gamma^5) \right) W' q_j + \text{H.c.}$$

- Both W'_R and W'_L could contribute
- Mass limits can be interpreted as a function of both couplings: a_{tb}^R , a_{tb}^L




$Z' \rightarrow tT'$



- But what if there are 3rd generation VLQ?
 - i.e. if something like this were possible:

- e.g.,
 - $Z' \rightarrow tT' \rightarrow t(bW)$
 - $\rightarrow t(tZ)$
 - $\rightarrow t(tH)$



- If kinematically allowed, $Z' \rightarrow tT'$ would dominate $Z' \rightarrow t\bar{t}$
- There's a loophole in $Z' \rightarrow t\bar{t}$ limits!

$Z' \rightarrow tT' \rightarrow t(bW)$ reconstruction

B2G-16-013

- “Mercedes sign” topology:

top jet:

- large R
- high pt
- top mass window
- \mathcal{T}_{32} (3-prong)



b-tagged
jet



W jet:

- large R
- W mass window
- \mathcal{T}_{21} (2-prong)

Large
T' mass
(kills $t\bar{t}$)

- Divide events by # of subject b-tags in top jet = total 1 or 2


$Z' \rightarrow tT'$: also capture ttH and ttZ

B2G-16-013

- “Mercedes sign” topology:


top jet:

- large R
- high pt
- top mass window
- \mathcal{T}_{32} (3-prong)





- Divide events by # of subject b-tags in top jet = total 1 or 2

- Also sensitive to ttH , ttZ :



$Z' \rightarrow tT'$: data in signal region

B2G-16-013




- 2b-tag category dominates: \sim same signal efficiency, but much less background

$Z' \rightarrow tT'$: limits for various Z' and T' masses

- Limits in 4D! Each plane is $\mathcal{B}(T \rightarrow bW)$ vs. $\mathcal{B}(T \rightarrow tH)$
 - $\mathcal{B}(T' \rightarrow tZ) = 1 - \mathcal{B}(T \rightarrow bW) - \mathcal{B}(T \rightarrow tH)$

B2G-16-013



- Low Z' masses limits worse (higher): lower top jet boost
- Worst for small $m_{Z'} - m_{T'}$ (top barely boosted!)
- bW corner more sensitive

$Z' \rightarrow tT'$: limits for fixed T' mass (1.2 TeV)

B2G-16-013

- Model 1: Composite Higgs

- $Z' \equiv \rho_L^0$

- T' branching fractions:



$$bW : tH : tZ = 0\% : 50\% : 50\%$$

- Model 2: Warped Extra Dim.

- $Z' \equiv G^*$ (KK gluon)

- T' branching fractions:

$$bW : tH : tZ = 50\% : 25\% : 25\%$$



Conclusions

- CMS has a broad program of searches for $t\bar{t}$, and tb resonances
- Traditional analyses performing well
- Improvements (e.g. $W' \rightarrow tb$) still possible and could yield large benefits
- New searches!
 - $Z' \rightarrow tT'$
 - many similar channels interesting

BACKUP MATERIAL

$Z' \rightarrow tT'$: background estimate

B2G-16-013

- Two categories: 1b-tag + 2b-tags
- Simulation:

