



The Standard Model Higgs and beyond

Michele Gallinaro

LIP Lisbon

April 17, 2017



- ✓ The Higgs boson and beyond
- ✓ (Charged Higgs in top quark decays)
- ✓ BSM Higgs: light pseudo-scalar, non-SM Higgs decay
- ✓ Higgs boson and Dark Matter




$$H \rightarrow ZZ \rightarrow 4e, 4\mu, 2e2\mu$$

- Signal: 4 isolated leptons from same vertex
 - Small background
 - Fully reconstructed, mass resolution $\sim 1\%$

The golden channel










2012: A new boson discovery



M. Gallinaro - "The Higgs boson and beyond" - April 17, 2017

July 4th, 2012: A Higgs boson








Higgs boson

PRD 89 (2014) 092007, PLB726(2013)088

- Progress since Higgs discovery (July 2012)
 - Observation in boson channels
 - Evidence for fermion couplings
 - Precision mass measurement (~ 125 GeV)
 - Spin determined
- It looks more like SM Higgs boson



Mass in the individual channels




- Most accurate measurement in the $\gamma\gamma$ and $4l$ channels
- Some “tension” between the four measurements (p-value $\sim 10\%$)

Couplings: individual channels





EPJC 75(2015)212, arXiv:1507.04548

Results based on the full
Run 1 data samples




Search for rare decays

PLB 726(2013)587, arXiv:1507.03031, JPLB 03(2015)048, CMS-HIG-15-012, PRL 114(2015)121801, ATLAS-CONF-2016-041



Standard Model theory of everything?

- Discovery of the Higgs boson marks the triumph of the SM
- However, even with the inclusion of the Higgs boson, SM is an incomplete theory



Beyond the Standard Model


The SM answers many of the questions about the structure of matter. But SM is not complete; still many unanswered questions:

- a) Why do we observe matter and almost no antimatter if we believe there is a symmetry between the two in the universe?
- b) What is this "dark matter" that we can't see that has visible gravitational effects in the cosmos?
- c) Are quarks and leptons actually fundamental, or made up of even more fundamental particles?
- d) Why are there three generations of quarks and leptons? What is the explanation for the observed pattern for particle masses?
- e) How does gravity fit into all of this?

Higgs and BSM

ATLAS-CONF-2015-044, CMS-HIG-15-002

- Is there BSM physics **hidden** in the “Higgs sector”?




$$(\sigma \cdot \text{BR})(gg \rightarrow H \rightarrow \gamma\gamma) = \sigma_{\text{SM}}(gg \rightarrow H) \cdot \text{BR}_{\text{SM}}(H \rightarrow \gamma\gamma) \cdot \frac{\kappa_g^2 \cdot \kappa_\gamma^2}{\kappa_H^2}$$

Strategy: parametrize deviations wrt SM in production and decay
 ⇒ loops are sensitive to BSM physics

Experimental approach


- Measure H(125) properties
- Search for additional Higgs bosons
- Search for BSM in signatures with Higgs bosons
- Search for BSM Higgs decays



Looking for new particles

JHEP08(2016)045


- Constrain BR_{BSM} in a scenario with free parameters
- $\Gamma_{tot} = \Gamma_{WW} + \Gamma_{ZZ} + \Gamma_{bb} + \dots + \Gamma_{BSM}$
- Likelihood scan vs BR_{BSM}
- Assuming couplings bound by SM expectations ($k_v < 1$)
- $0 \leq BR_{BSM} \leq 0.34$ at 95%CL




Couplings: decays

ATLAS-CONF-2015-044, CMS-HIG-15-002, JHEP08(2016)045

BSM physics in the loop



Vector and fermion couplings



BR_{BSM} can be measured

$BR_{\text{BSM}} < 0.34$ at 95% C.L. (assuming $\kappa_V \leq 1$)

BR_{BSM} includes non standard decays, visible or invisible


⇒ Results in agreement with SM ($\kappa_V = \kappa_F = 1$) within 1σ

High mass: $H \rightarrow WW/ZZ$

JHEP 10(2015)144, HIG-15-004

- Search for a heavy Higgs boson
 - $H \rightarrow ZZ \rightarrow 4\ell, 2\ell 2\nu, 2\ell qq$
 - $H \rightarrow WW \rightarrow 2\ell 2\nu, 2\ell qq$
- optimized separately for VBF and gluon fusion production processes
- SM-like Higgs boson excluded in 4ℓ and $2\ell 2\nu/\ell\nu qq$ channels at 95%CL in mass ranges up to 1000 GeV
- Search interpreted in BSM scenario (heavy Higgs, heavy EWK singlet state)
 - evolution of signal strength of the singlet state with modified couplings/width wrt SM.
 - assume new scalar does not decay to any new particle

No significant excess
 \Rightarrow set limits: $\sigma \cdot B(X \rightarrow ZZ \rightarrow 4\ell)$
 \sim a few fb @ $m_X = 400$ GeV
 \sim 1 fb @ $m_X = 1$ TeV



high-mass searches improve at 13TeV

Extending searches

- Minimal Supersymmetric SM (MSSM)
 - Neutral Higgs: $\phi \rightarrow \tau\tau/bb/\mu\mu$
 - Charged Higgs
- Next-to-MSSM
 - Light pseudoscalar: $h \rightarrow aa$
 - Non-SM decays: $h \rightarrow 2a \rightarrow 4\tau/4\mu$
 - Heavy Higgs: $H \rightarrow h_{125} h_{125}$ or $A \rightarrow Zh_{125}$
- FCNC: $t \rightarrow cH$

Higgs sector in the MSSM

Higgs sector in SUSY contains two scalar doublets:



- 5 physical Higgs bosons
 - 3 neutral: CP-even $\phi=h, H$ CP-odd A
 - 2 charged H^\pm
- SM-like Higgs boson: h

Neutral Higgs ϕ decay modes:

- $BR(\phi \rightarrow b\bar{b}) \sim 90\%$
- $BR(\phi \rightarrow \tau\tau) \sim 10\%$
- $BR(\phi \rightarrow \mu\mu) \sim 0.1\%$

Two main production modes:



- $gg \rightarrow H$
- $b\bar{b}H$



Neutral MSSM Higgs

JHEP 10(2014)212, arXiv:1409.6064

- Enhanced couplings of MSSM Higgs to down-type fermions (large $\tan\beta$)
 \Rightarrow increased BR to τ leptons and b-quarks
- Search for neutral MSSM Higgs boson
- 5 final states used: $\mu\tau_h$, $e\tau_h$, $\tau_h\tau_h$, $e\mu$, $\mu\mu$
 - Reconstruct tau-pair invariant mass
 - Split in b-tag/no b-tag categories to enhance sensitivity
- Main backgrounds: $Z \rightarrow \tau\tau$, QCD/W+jets, DY, ttbar, dibosons




Neutral MSSM Higgs: $\phi \rightarrow \tau\tau$


JHEP 10(2014)212, arXiv:1409.6064, CMS-HIG-16-037

- Direct search: inclusive and b-tagged
- τ in both leptonic and hadronic decays

$\tan\beta$ vs m_A window becoming smaller



Model-independent limits by separating production modes







No significant excess over bkg expectations

Neutral MSSM Higgs: $\phi \rightarrow \mu\mu$


arXiv:1508.01437, ATLAS-CONF-2012-094

- Search for a $\mu\mu$ mass resonance
- Good mass resolution
 - full and clean reconstructed final state
- Split in b-tagged and non b-tagged categories to be sensitive to $gg \rightarrow \phi$ and $bb\phi$ production modes
- Main backgrounds: $Z(b\bar{b})$, $t\bar{t}$, WW





Charged Higgs

- If found, a clear indication of BSM
- Study non-SM Higgs in **two mass regimes**:
- $m_H < m_{\text{top}}$
 - Mostly produced in top quark decays
 - Large $\tan\beta$: $H^\pm \rightarrow \tau^+\nu$
 - Small $\tan\beta$ (<1): $H^\pm \rightarrow c\bar{s}$
- $m_H > m_{\text{top}}$
 - Produced in gluon-gluon fusion
 - Main decays: $H^\pm \rightarrow tb$, $H^\pm \rightarrow \tau^+\nu$
- Main backgrounds: $t\bar{t}$, W +jets




Charged Higgs (cont.)

- Different strategies for low- and high-mass searches
- tau+lepton, lep+jets, and $e\mu$ final states
- b-tagged jet categorization
- limited by statistics at high-mass



$$m_H < m_{top}$$




$$m_H > m_{top}$$


Charged Higgs and top quark decays

JHEP 07(2012)143, arXiv:1508.07774, HIG-16-031


- Look for charged Higgs in four final states:
 - Tau+lepton (electron or muon)
 - Dilepton (tau decays leptonically)
 - lepton+jets
 - Fully hadronic: tau+jets




τ_h +jets



τ_h +lepton (e/μ)



di-lepton (eμ)




Looking at tau decays

CMS-HIG-12-052

Low H^+ mass:



- Use R variable in the limit extraction: binned maximum-likelihood fit
- Tau fake component is data-driven, includes uncertainties

SM






VS

BSM



Number of b-tagged jets




High-mass H^\pm search: look at b-tag multiplicity



Is there a charged Higgs?

JHEP 07(2012)143, CMS-HIG-12-052, arXiv:1508.07774

- If anomalous tau/lepton production in $t\bar{t}$ decays there may be contribution from H^\pm
- Yields in agreement with expectations \Rightarrow set limits
- m_{H^\pm} : 80-160 GeV $\mathcal{B}(t \rightarrow bH^\pm) < 1.2-0.3\%$
 200-600 GeV $\sigma(pp \rightarrow \bar{t}(b)H^\pm) < 2.0-0.2 \text{ pb}$



At 13TeV, expect improvement with 5-10/fb for $m_{H^\pm} > 300 \text{ GeV}$




- $t\bar{t}$ bar xsection increases x3.3
- signal increases x6(x7) for $m_{H^\pm} = 500(600) \text{ GeV}$

Still hope for MSSM?

JHEP 07(2012)143, CMS-HIG-12-052, arXiv:1508.07774

- A new modified MSSM scenario: m_h^{mod} (arXiv:1302.7033)
- Reduce amount of mixing in the stop sector (X_t/M_{SUSY})
- A/H decays to chargino/neutralinos allowed (arXiv:0709.1029)
- Allows for reduction of decays into $\tau\tau$ and bb



Cross section ratios

PRD 80(2009) 071102




Many systematic unc. cancel in the ratio

Study of cross section ratios

⇒ sensitive to BSM

1. $BR(\ell+jets)/BR(\ell\ell)$



2. $BR(\ell+\tau)/BR(\ell\ell)$



Combination of more channels

JHEP 03(2013)076

- Search for charged Higgs boson
- Use $\tau_{\text{had}} + \text{lep}$ and $\tau_{\text{had}} + \text{jets}$ final states
 - compare to $e\mu$ yields
- Search for anomalous decays






Set limits on:

$$B(t \rightarrow bH^+)$$

Light charged Higgs: $c\bar{s}$

JHEP 12(2015)1, arXiv:1510.04252

- $H \rightarrow c\bar{s}$ decay
 - dominant in low $\tan\beta$ region
- Lepton+jet final states
- Dominant bkg from $t\bar{t}$
- Kinematic fit to reconstruct W/H mass
- Set model-independent limits on $BR(t \rightarrow H^+b) \sim 2-7\%$



Doubly charged Higgs

EPJC 72 (2012) 2189, CMS-HIG-14-039, HIG-16-036




• Model

- SM extended with scalar triplet (Φ^{++} , Φ^+ , Φ^0)
- Triplet responsible for neutrino masses
- Search for doubly- and singly-charged
- DY pair production is most common
- SS lepton pair of any flavor combination

• Search with ≥ 3 leptons of any flavor



- Search for excess of events in one or more flavor combinations of SS lepton pairs

• Dilepton invariant mass as discriminant




non-SM Higgs decay: $h \rightarrow aa \rightarrow 4X$

- Standard search for light (pseudo)- scalar Higgs with $m_a < m_h/2$
 - generic prediction of BSM theories (extended Higgs sector, NMSSM, etc)
 - Final states go to fermions (b, τ, μ, \dots)
 - BR depends on boson mass, model parameters




non-SM Higgs decay: $h \rightarrow aa \rightarrow 4\mu$

PLB 726(2013)564, arXiv:1506.00424, HIG-16-035



- Explore non-SM decays of a Higgs boson (h)
 - Higgs boson (h) can be SM or not
 - include production of two new light boson (a^0)
- Search for generic Higgs decays: $h \rightarrow 2a + X \rightarrow 4\mu + X$
 - Require two dimuon pairs with consistent masses
 - Observe 9 events in off-diagonal region
 - Signal region: **1 event** (2.2 ± 0.7 bkg)
 - Limits on production rates, benchmark models




NMSSM and Dark SUSY Limits

PLB 726(2013)564, arXiv:1506.00424



Results interpreted in NMSSM and dark SUSY

- Dark SUSY: h decay to pair of neutralinos (n_1): LSP

$n_1 \rightarrow n_D \gamma_D$ decays
 $\rightarrow \mu\mu$
 \rightarrow invisible






- NMSSM: Extend MSSM by adding a complex singlet field (1 CP-even+1 CP-odd boson)
- NMSSM: $h_{1,2} \rightarrow 2a_1$; $a_1 \rightarrow 2\mu$
- Compare to SM Higgs cross section




non-SM Higgs decay: $H_{125} \rightarrow 2h(a) \rightarrow 4\tau$

JHEP01(2016)079

- Search for **very light Higgs** in NMSSM
 - $h_{1,2}$ (CP-even), $a_{1,2}$ (CP-odd) to a pair of τ leptons
 - $H(125) \rightarrow h_1 h_2$ ($a_1 a_2$) $\rightarrow 4\tau$
- Reconstruct μ -track invar. mass (m_1, m_2)
 - SS dimuon sample (removes DY)
 - bin in 2-dim distribution, fit signal and bkg
 - QCD bkg from control region
- No excess over SM backgrounds








Summary for Higgs exotic decays



Low mass Higgs: $a(\rightarrow\tau\tau)bb$

arXiv:1511.03610

- Low mass Higgs in the NMSSM
- Low mass pseudo-scalar ($a_1 \rightarrow \tau\tau$) in association with $b\bar{b}$: $a_1 b\bar{b} \rightarrow \tau\tau b\bar{b}$
- Similar strategy to $H \rightarrow \tau\tau$
- Search for a_1 masses below Z mass
- No evidence for signal
- Set limits: $\sigma \times B \sim 9-39$ pb




di-Higgs searches


- Destructive interference in SM
- Could be altered in BSM
- If constructive, it could be large enhancement
- In SM, only $\sigma=33\text{fb}$ at 13 TeV
- Study different final states

	BR	Mass scale
$(X \rightarrow) hh \rightarrow$	$bbbb$ 34%	High
	$bb\tau\tau$ 7.3%	
	$bbWW$ 27%	
	$bb\gamma\gamma$ 0.26%	Low

non-resonant production




resonant production




Heavy Higgs to $h_{125}h_{125} \rightarrow \tau\tau bb$

CMS-EXO-15-008, CMS-HIG-16-012, CMS-HIG-17-002




- **Resonant and non-resonant production**
 - Double Higgs production to determine λ_{hhh}
 - Check couplings: $\kappa_\lambda = \lambda_{hh}/\lambda_{hhh}^{SM}$; $\kappa_t = y_t/y_t^{SM}$
 - BSM could enhance non-resonant hh production
 - $H \rightarrow h_{125}h_{125} \rightarrow bb\tau\tau$
- **h_{125} decay products nearly collinear**
 - boosted “single” merged jet ($\rightarrow bb$)
- **use $\tau_e\tau_h$, $\tau_\mu\tau_h$, and $\tau_h\tau_h$ final states**
 - sidebands/inverted isolation to estimate bkg
- **set limits as function of mass**





Heavy Higgs: $H \rightarrow h_{125}h_{125}$, $A \rightarrow Zh_{125}$

PRD90(2014)112013, PLB755(2016)217

- MSSM: Heavy Higgs searches
 - Search for $A \rightarrow Zh_{125}$ and $H \rightarrow hh$
- Exclusive search in **multilepton** and **+lepton** channels
- Search for FCNC decays
- Search for $tt \rightarrow (bW)(ch)$
 - Not forbidden but **highly suppressed**
 - enhanced w/some parameter models
- SM Higgs now a background
 - ATLAS: $H \rightarrow \gamma\gamma$
 - CMS: $H \rightarrow \gamma\gamma$ and multileptons
- b-tag provides bkg suppression



- $H \rightarrow WW^* \rightarrow l\nu l\nu$,
- $H \rightarrow \tau\tau$,
- $H \rightarrow ZZ^* \rightarrow jjll, \nu\nu ll, llll$,
- $H \rightarrow \gamma\gamma$.



Process	SM	QS	2HDM-III	FC-2HDM	MSSM
$t \rightarrow u\gamma$	$3.7 \cdot 10^{-16}$	$7.5 \cdot 10^{-9}$	—	—	$2 \cdot 10^{-6}$
$t \rightarrow uZ$	$8 \cdot 10^{-17}$	$1.1 \cdot 10^{-4}$	—	—	$2 \cdot 10^{-6}$
$t \rightarrow uH$	$2 \cdot 10^{-17}$	$4.1 \cdot 10^{-5}$	$5.5 \cdot 10^{-6}$	—	10^{-5}
$t \rightarrow c\gamma$	$4.6 \cdot 10^{-14}$	$7.5 \cdot 10^{-9}$	$\sim 10^{-6}$	$\sim 10^{-9}$	$2 \cdot 10^{-6}$
$t \rightarrow cZ$	$1 \cdot 10^{-14}$	$1.1 \cdot 10^{-4}$	$\sim 10^{-7}$	$\sim 10^{-10}$	$2 \cdot 10^{-6}$
$t \rightarrow cH$	$3 \cdot 10^{-15}$	$4.1 \cdot 10^{-5}$	$1.5 \cdot 10^{-3}$	$\sim 10^{-5}$	10^{-5}

FCNC decays

BR($t \rightarrow cH$) (95%CL)

ATLAS obs(exp)
<0.79% (0.51%)




CMS
<0.56% (0.65%)

LFV in Higgs decays

PLB 763(2016)472, CMS-HIG-16-005

- Some BSM models allow for LFV Higgs decays
- Search for $H \rightarrow e\tau, e\mu, \mu\tau$ final states
- Categories: N_{jet} , lepton kinematics
 - N_{jet} to target ggH and VBF production
- Main background from DY, ttbar, WW


	95%CL (obs/exp)	Best fit
$h \rightarrow \mu\tau$ (run1)	<1.51/0.75%	$0.84^{+0.39}_{-0.37}\%$
$h \rightarrow \mu\tau$ (run2)	<1.20/1.62%	$-0.76^{+0.81}_{-0.84}\%$



Dark Matter+Higgs



arXiv:1510.06218, arXiv:1506.01081

- Generic search: $pp \rightarrow X + \text{MET}$
- Search for DM + $h(\rightarrow bb)$
- Model-independent search
 - Signature: $h(\rightarrow ZZ/bb/\gamma\gamma) + \text{MET}$
 - Simplified model with Z' or pseudo-scalar Higgs $A(\rightarrow \chi\chi)$



DM particle (χ): can be scalar or fermion
Pseudo-scalar Higgs A

Signal events at large MET



Summary

- Excellent consistency of SM but **SM is incomplete**
- Extensions foresee existence of additional bosons
- Searches for BSM bosons natural companion to precision SM Higgs boson measurements
 - Charged Higgs searches with top quark decays
 - Other BSM searches show no indication of deviations
- Searches provide **no hints for BSM yet**

