### Double parton distributions in QCD

Krzysztof Golec-Biernat

Institute of Nuclear Physics in Kraków and Rzeszow University

WE-Heraeus Physics School, Bad Honnef, September 24-30, 2017

**∃** ⊳

- Double parton scattering
- Double parton distributions
- Evolution equations
- DPS cross section computation
- Transverse momentum dependent double distributions

э

ヨト イヨト

### Multiparton interactions

► Hard processes with the scale  $Q \gg \Lambda \sim 1 \text{ GeV}$  due to collisions of quarks and gluons.

- At the LHC multiparton interactions (MPI) become increasingly important.
- ▶ If no hard scale is involved, MPI are crucial for modeling of underlying event.
- ► For rising center-of-mass energy MPI lead to more hard scatterings.

(E) < E)</p>

For example, two vector bosons with transverse momenta  $q_1$  and  $q_2$ 



Single parton scattering (SPS)

Double parton scattering (DPS)

► Introduction to DPS: M. Diehl, D. Ostermeier, A. Schafer, JHEP 1203 (2012) 089

## DPS versus SPS cross sections



• Inclusive DPS is enhanced due to rising parton densities for  $x \rightarrow 0$ 

 $d\sigma^{SPS} \sim x^{-2\lambda}$   $d\sigma^{DPS} \sim x^{-4\lambda}$ 

▶ SPS suppressed in same sign vector boson production  $W^{\pm}W^{\pm}$ .

Pocket formula

$$\sigma_{AB}^{\rm DPS} = \frac{1}{1 + \delta_{AB}} \frac{\sigma_A^{\rm SPS} \sigma_B^{\rm SPS}}{\sigma_{\rm eff}}$$

• Effective cross section:  $\sigma_{\rm eff} \approx 15 \, {\rm mb}$ .



.⊒ →

#### Recent update

Experiment (energy, final state, year) ATLAS ATLAS ( $\sqrt{s} = 8$  TeV,  $J/\psi + J/\psi$ , 2016) Hel DØ  $(\sqrt{s} = 1.96 \text{ TeV}, J/\psi + J/\psi, 2014)$ нон DØ ( $\sqrt{s} = 1.96$  TeV, J/ $\psi + \Upsilon$ , 2016) HAH LHCb ( $\sqrt{s} = 7\&8$  TeV,  $\Upsilon(1S) + D^{0,+}$ , 2015) HV4 LHCb ( $\sqrt{s} = 7$  TeV,  $J/\psi + \Lambda_c^+$ , 2012) LHCb ( $\sqrt{s} = 7$  TeV,  $J/\psi + D_{*}^{+}$ , 2012) ----LHCb ( $\sqrt{s} = 7$  TeV,  $J/\psi + D^+$ , 2012) \_\_\_\_ LHCb ( $\sqrt{s} = 7$  TeV, J/ $\psi$  + D<sup>0</sup>, 2012) ATLAS ( $\sqrt{s} = 7$  TeV, 4 jets, 2016) CDF ( $\sqrt{s} = 1.8$  TeV, 4 jets, 1993) UA2 ( $\sqrt{s} = 630$  GeV, 4 jets, 1991) AFS ( $\sqrt{s} = 63$  GeV, 4 jets, 1986) н DØ ( $\sqrt{s} = 1.96$  TeV,  $2\gamma + 2$  jets, 2016) DØ ( $\sqrt{s} = 1.96$  TeV,  $\gamma + 3$  jets, 2014) Ed.  $DO(\sqrt{s} = 1.96 \text{ TeV}, \gamma + b/c + 2 \text{ jets}, 2014)$ DØ ( $\sqrt{s} = 1.96$  TeV,  $\gamma + 3$  jets, 2010) CDF ( $\sqrt{s} = 1.8$  TeV,  $\gamma + 3$  jets, 1997) HOH ATLAS ( $\sqrt{s} = 8$  TeV,  $Z + J/\psi$ , 2015) CMS ( $\sqrt{s} = 7$  TeV, W + 2 jets, 2014) ATLAS ( $\sqrt{s} = 7$  TeV, W + 2 jets, 2013) 10 15 25 0 5 20 30  $\sigma_{eff}$  [mb]

**∃** ⊳

## LHC strategy for DPS



## SPS and DPS cross sections

▶ Partons with transverse momenta  $k_{\perp} \sim \Lambda \ll Q$  (naive parton model)



• For SPS single parton distributions (PDF) with 0 < x < 1

$$\frac{d\sigma_{AB}^{SPS}}{dx \, d\overline{x}} = \sum_{i\overline{i}} D_i(x) \, \sigma_{i\overline{i}}^{AB} \, D_{\overline{i}}(\overline{x})$$

▶ For DPS double parton distributions (DPDF) with  $0 < x_1 + x_2 < 1$ 

$$\frac{d\sigma_{AB}^{DPS}}{dx_1 dx_2 d\overline{x}_1 d\overline{x}_2} = \sum_{i\overline{i}j\overline{j}} \int \frac{d^2\mathbf{q}}{(2\pi)^2} D_{ij}(x_1, x_2, \mathbf{q}) \ \sigma_{j\overline{i}}^A \sigma_{j\overline{j}}^B \ D_{\overline{i}\overline{j}}(\overline{x}_1, \overline{x}_2; -\mathbf{q})$$

▲글▶ ▲글▶

Single PDF

$$D_i(x) = \int dz^- e^{i(xP^+)z^-} \langle P| \underbrace{\bar{\Psi}_i(0,0,\mathbf{0})\gamma^+\Psi_i(0,z^-,\mathbf{0})}_{\mathcal{O}_i(0,z)} |P\rangle$$

- $\mathcal{O}_i(0, z)$  connects two points on the light cone
- Double PDF

$$D_{ij}(x_1, x_2, \mathbf{q}) = \int dz_1^- dz_2^- e^{i(x_1 z_1^- + x_2 z_2^-)P^+} \int dy^- d^2 \mathbf{y} e^{-i\mathbf{q}\cdot\mathbf{y}}$$
$$\times \langle P | \mathcal{O}_j(0, z_2) \mathcal{O}_i(y, z_1) | P \rangle$$

- ▶ O<sub>j</sub>(y, z<sub>2</sub>) does not connect light cone points since y = (0, y<sup>-</sup>, y) has transverse component y.
- ▶ Transverse momentum **q** is Fourier conjugate to transverse separation **y**.

3

< 🗇 🕨

•  $\langle P | \mathcal{O}_i(0, z_2) \mathcal{O}_i(y, z_1) | P \rangle$  has 4 fermionic operators. In momentum space



- Momentum conservation leads to 3 independent vectors: k<sub>1</sub>, k<sub>2</sub>, q.
- Integrated transverse momentum dependent distributions

$$D^{(ab)}_{(lphaeta)(ij)}(x_1,x_2,{f q}) = \int d^2{f k}_1\, d^2{f k}_2\, F^{(ab)}_{(lphaeta)(ij)}(x_1,x_2,{f k}_1,{f k}_2,{f q})$$

▶ *D<sub>ij</sub>* is color singlet, spin averaged transverse momentum integrated distribution.

#### Geometric interpretation of DPDF

> y is the transverse separation between the two partons in a hadron



> Partons with the same transverse position interact with each other

$$\frac{d\sigma_{AB}^{DPS}}{dx_1 dx_2 d\overline{x}_1 d\overline{x}_2} = \sum_{i\overline{i}j\overline{j}} \int d^2 \mathbf{y} \, d^2 \mathbf{b} \, d^2 \mathbf{b}' \, \tilde{D}_{ij}(x_1, x_2, \mathbf{y}, \mathbf{b}) \, \sigma_{i\overline{i}}^A \, \sigma_{j\overline{j}}^B \, \tilde{D}_{\overline{i}\overline{j}}(\overline{x}_1, \overline{x}_2, \mathbf{y}, \mathbf{b}')$$

## Evolution equations for single PDF

▶ Partons with transverse momenta  $\Lambda \ll k_{\perp} \ll Q$  (improved parton model)



- Collinear divergences must be subtracted.
- ▶ PDF become factorization scale dependent,  $D_f(x, Q)$ .
- DGLAP evolution equations

$$\frac{\partial}{\partial \ln Q^2} D_f(x, Q) = \frac{\alpha_s}{2\pi} \sum_{f'} \int_x^1 du \, \mathcal{P}_{ff'}\left(\frac{x}{u}\right) D_{f'}(u, Q)$$

+ <u>initial conditions</u>  $D_f(x, Q_0)$ , determined from global fits to data.



Can we obtain the same precision for DPDF ?

-

### Evolution equations for DPDF

• For equal scales,  $Q_1 = Q_2 \equiv Q$ , and  $\mathbf{q} = \mathbf{0}$ 

$$D_{f_1f_2}(x_1,x_2,Q,\mathbf{q}=0) = \int d^2\mathbf{y} \, D_{f_1f_2}(x_1,x_2,Q,\mathbf{y})$$

Evolution equations (Snigirev hep-ph/0304172)



Coupling with DGLAP equations for PDF in splitting term.

#### Sum rules obeyed by evolution equations

- How to determine initial conditions  $D_{f_1 f_2}(x_1, x_2, Q_0)$ ?
- Momentum and quark number sum rules obeyed by single PDF

$$\sum_{f} \int_{0}^{1} dx \times D_{f}(x) = 1$$
$$\int_{0}^{1} dx \left( D_{q}(x) - D_{\overline{q}}(x) \right) = N_{q} \qquad (=2,1,0)$$

Momentum and quark number sum rules obeyed DPDF

$$\sum_{f_1} \int_0^{1-x_2} dx_1 x_1 D_{f_1 f_2}(x_1, x_2) = (1-x_2) D_{f_2}(x_2)$$
$$\int_0^{1-x_2} dx_1 \left\{ D_{qf_2}(x_1, x_2) - D_{\bar{q}f_2}(x_1, x_2) \right\} = (N_q - \delta_{qf_2} + \delta_{\bar{q}f_2}) D_{f_2}(x_2)$$

▶ Sum rules for DPDF constrain  $D_{f_1f_2}(x_1, x_2, Q_0)$ . Try to impose them.

E > < E >

#### Initial conditions for DPDF

 Most popular initial condition at Q<sub>0</sub> are built from the known single PDF (J.R. Gaunt, W.J. Stirling, arXiv:0910.4347)

$$D_{f_1f_2}(x_1, x_2) = D_{f_1}(x_1) D_{f_2}(x_2) \frac{(1 - x_1 - x_2)^2}{(1 - x_1)^{2 + n_1} (1 - x_2)^{2 + n_2}}$$

- They obey sum rules for DPDF only approximately.
- Factorizable form for  $x_1, x_2 \ll 1$

ratio = 
$$\frac{D_{f_1 f_2}(x_1, x_2)}{D_{f_1}(x_1) D_{f_2}(x_2)} = 1$$

- Study this ratio as a function of  $Q^2$  during evolution  $\rightarrow$  factorization test.
- Can we go beyond the factorized ansatz?

E > < E >

#### Pure gluonic case

(GB, Lewandowska, Serino, Snyder, Stasto, arXiv:1606.01679)

▶ MSTW08 gluon distr. at  $Q_0 = 1$  GeV with known parameters  $A_k, \alpha_k, \eta$ 

$$D_g(x) = \sum_{k=1}^{3} A_k x^{\alpha_k} (1-x)^{\gamma_k}$$

Solve momentum sum rule

$$\int_0^{1-x_2} dx_1 x_1 D_{gg}(x_1, x_2) = (1-x_2) D_g(x_2)$$

for the ansatz

$$\int_{0}^{1-x_{2}} dx_{1} x_{1} \underbrace{\sum_{k=1}^{3} \bar{N}_{k} (x_{1} x_{2})^{a_{k}} (1-x_{1}-x_{2})^{b_{k}}}_{D_{gg}(x_{1},x_{2})} = (1-x_{2}) \underbrace{\sum_{k=1}^{3} \bar{A}_{k} x_{2}^{a_{k}} (1-x_{2})^{a_{k}+b_{k}+1}}_{D_{gg}(x_{2})}$$

From the comparison with MSTW08 gluon

 $ar{A}_k = A_k$ ,  $a_k = lpha_k$ ,  $a_k + b_k + 1 = \eta$ 

Non-factorizable initial double gluon distribution.

3

Numerical results for  $x_2 = 10^{-2}$ 



Krzysztof Golec-Biernat Double parton distributions in QCD

## Numerical results for $x_2 = 0.5$



Krzysztof Golec-Biernat Double parton distributions in QCD

## Adding quarks

 Light-cone Fock expansion of the proton state in terms of partonic states is the origin of the sum rules

$$|P\rangle = \sum_{N} \sum_{f_1...f_N} \int dx_1 \dots dx_N \,\delta\Big(1 - \sum_{k=1}^N x_k\Big) \,\Psi_N\big(x_1 \dots x_n; f_1 \dots f_N\big) \,|x_1 \dots x_N; f_1 \dots f_N\rangle$$

• Try to model wave functions  $\Psi_N$ 

$$D_f(x) = \sum_N \sum_{f_1...f_N} \int d\Pi_N \left| \Psi_N \right|^2 \left\{ \sum_{i=1}^N \delta(x - x_i) \, \delta_{ff_i} \right\}$$

$$D_{fh}(x,y) = \sum_{N} \sum_{f_1...f_N} \int d\Pi_N |\Psi_N|^2 \left\{ \sum_{i=1}^N \sum_{j\neq i}^N \delta(x-x_i) \delta(y-x_j) \, \delta_{ff_i} \delta_{hf_j} \right\}$$

• What  $\Psi_N$  leads to the MSTW08 parameterization?

э

E > < E >

## A model

Assume (Broniowski, Ruiz Ariola, GB, arXiv:1602.00254)

$$|\Psi_{N}(x_{1}...x_{N};f_{1}...f_{N})|^{2} = A_{f_{1}...f_{N}}^{N} x_{1}^{\alpha_{f_{1}}^{N}-1} x_{2}^{\alpha_{f_{2}}^{N}-1} ...x_{N}^{\alpha_{f_{N}}^{N}-1}$$

Single PDF

$$D_f(x) = \sum_{N} \sum_{(f_1...f_N)'} \bar{A}_{f_1...f_N}^N x^{\alpha_f^N - 1} (1 - x)^{\alpha_{(f_1}^N + ... + \alpha_{f_N}^N)'} = 0$$

• Small x powers  $\alpha_{f_i}^N$  determine the large x powers

$$\eta_f^N = \alpha_{(f_1}^N + \ldots + \alpha_{f_N}^N)'$$

Such a relation is not valid for MSTW08 parameterization of PDF.

э

▲ 臣 ▶ | ▲ 臣 ▶ |

- Non-factorizable initial condition which fulfill momentum sum rule in the pure gluon case was constructed.
- ► Factorization of double gluon distribution at small x sets rather quickly

 $D_{gg}(x_1, x_2, Q) \approx D_g(x_1, Q) \cdot D_g(x_2, Q)$ 

- Extension to the case with quarks has not been successful.
- ► Factorizable Gaunt-Stirling prescription is still mostly used.

3

医下 不正下

### DPS cross section computation

Solution to the DPDF evolution equations

 $D_{f_1f_2}(x_1, x_2, Q, \mathbf{q} = \mathbf{0}) = D_{f_1f_2}^{(1)}(x_1, x_2, Q) + D_{f_1f_2}^{(2)}(x_1, x_2, Q)$ 



Sum of hadronic and point-like contributions

$$D(\mathbf{x}_{1}, \mathbf{x}_{2}, \mathbf{Q}, \mathbf{q}) = U_{1}(\mathbf{Q}, \mathbf{Q}_{0}) \otimes \underbrace{D(\mathbf{x}_{1}, \mathbf{x}_{2}, \mathbf{Q}_{0})}_{initial DPDF} \otimes U_{2}^{T}(\mathbf{Q}, \mathbf{Q}_{0}) \times F^{2}(\mathbf{q})$$

$$+ \int_{max\{|\mathbf{q}|, \mathbf{Q}_{0}\}}^{Q} d\mathbf{Q}' U_{1}(\mathbf{Q}, \mathbf{Q}') \otimes \underbrace{D(\mathbf{x}_{1} + \mathbf{x}_{2}, \mathbf{Q}')}_{single PDF} \otimes P(\mathbf{Q}') \otimes U_{2}^{T}(\mathbf{Q}, \mathbf{Q}')$$

q-prescription from M.G. Ryskin, A.M. Snigirev, arXiv:1103.3495

Partonic form factor (Frankfurt, Strikman)

$$F^2(q^2) = rac{1}{\left(1+q^2/m^2
ight)^4} \qquad m_g pprox 1.5 \ {
m GeV}$$

DPS cross section written in terms of the two components

$$\sigma_{AB} = \int d^2 \mathbf{q} \, (D^{(1)} + D^{(2)}) \, \sigma_A \, \sigma_B \, (D^{(1)} + D^{(2)})$$
$$= \sigma_{AB}^{(11)} + \, \sigma_{AB}^{(12+21)} + \, \sigma_{AB}^{(22)}$$

• Pocket formula from  $\sigma^{(11)}_{AB}$  with factorized  $D^{(1)}$  and

$$rac{1}{\sigma_{
m eff}} = \int d^2 \mathbf{q} \, F^4(\mathbf{q}) \qquad pprox \qquad (15\,{
m mb})^{-1}$$

• How important are splitting contributions  $\sigma_{AB}^{(12+21)}$  and  $\sigma_{AB}^{(22)}$  ?

< ≣ >

## Splitting contributions



► W<sup>+</sup>W<sup>-</sup> production in DPS from GB, Lewandowska, arXiv:1407.4038



•  $\sigma_{AB}^{(12+21)}$  and  $\sigma_{AB}^{(22)}$  are important. (also Gaunt, Maciuła, Szczurek, arXiv:1407.5821)

∃⇒

## How to avoid double counting ?

Are the splitting graphs part of SPS or DPDF? (Diehl, Gaunt, Schönwald, arXiv:1702.06486)



 $\blacktriangleright$  Procedure based on removing  $1/\mathbf{y}^2$  divergence in the point-like DPDF

$$D_{f_1f_2}^{(2)}(x_1, x_2, \mathbf{y}) = \frac{1}{\mathbf{y}^2} \frac{\alpha_s}{2\pi} \sum_{f'} D_{f'}(x_1 + x_2) P_{f' \to f_1f_2}^R \left(\frac{x_1}{x_1 + x_2}\right)$$

by introducing monotonic cut-off function  $\Phi(u) \in [0, 1]$  to

$$\sigma_{
m DPS} \sim \int d^2 {f y} \, |\Phi(Q{f y})|^2 \, D_{f_1 f_2}(x_1,x_2,{f y}) \, D_{f_3 f_4}(x_1,x_2,{f y})$$

and subtraction term to the cross section

$$\sigma = \sigma_{\rm DPS} - \sigma_{\rm sub} + \sigma_{\rm SPS}$$

• For  $y \ll 1/Q$   $\sigma_{\rm DPS} \approx \sigma_{\rm sub}$  while for  $y \gg 1/Q$   $\sigma_{\rm SPS} \approx \sigma_{\rm sub}$ 

Need of TMDD

$$D_{f_1f_2}(x_1, x_2, \mathbf{y}) = \int d^2 \mathbf{k}_1 d^2 \mathbf{k}_2 \underbrace{F_{f_1f_2}(x_1, x_2, \mathbf{k}_1, \mathbf{k}_2, \mathbf{y})}_{\text{TMDD}}$$

be able to compute more exclusive cross sections

$$\frac{d\sigma_{AB}^{DPS}}{dx_i d\bar{x}_i d^2 \mathbf{q}_i} = \sigma_A \sigma_B \int d^2 \mathbf{k}_i \, d^2 \bar{\mathbf{k}}_i \, d^2 \mathbf{y} \, \delta^2(\mathbf{q}_i - \mathbf{k}_i - \bar{\mathbf{k}}_i) \, F(\mathbf{x}_i, \mathbf{k}_i, \mathbf{y}) \, F(\bar{\mathbf{x}}_i, \bar{\mathbf{k}}_i, \mathbf{y})$$

- Unfolding evolution equations for DPDF (for TMDD integrated over y or at q = 0) (GB, Stasto, arXiv:1611.02033).
- ▶ Analysis with y-dependent TMDD, following Collins-Soper method. TMDD factorization proved for  $q_1 \ll Q_i$  by extending proof for TMD. (Buffing, Diehl, Kasemets, arXiv:1708.03528)

B 🕨 🖌 B 🕨

Unfolding DGLAP evolution in the last step (Kimber, Martin, Ryskin). For TMD



•  $T_a(Q, \mathbf{k})$  is Sudakov form factor. For TMDD (GB, Stasto, arXiv:1611.02033)

$$F_{a_{1}a_{2}}(x_{i},\mathbf{k}_{i},Q_{i}) = T_{a_{1}}(Q_{1},\mathbf{k}_{1}) T_{a_{2}}(Q_{2},\mathbf{k}_{2}) \times \\ \times \sum_{b,c} \int_{\frac{x_{1}}{1-x_{2}}}^{1-\Delta(\mathbf{k}_{1})} \frac{dz_{1}}{z_{1}} \int_{\frac{x_{2}}{1-x_{1}/z_{1}}}^{1-\Delta(\mathbf{k}_{2})} \frac{dz_{2}}{z_{2}} P_{a_{1}b}(z_{1},\mathbf{k}_{1}) P_{a_{2}c}(z_{2},\mathbf{k}_{2}) D_{bc}\left(\frac{x_{1}}{z_{1}},\frac{x_{2}}{z_{2}},\mathbf{k}_{1},\mathbf{k}_{2}\right)$$

TMD and TMDD built out of known PDF and DPDF.

-∢ ≣ ▶

# $F_{gg}(\mathbf{k}_1, \mathbf{k}_2)$ in pure gluon case



 $Q^2 = 100 x_1 = x_2 = 0.01$ 

- Double parton scattering processes are being measured.
- > Double parton distributions are well defined object with known properties.
- Significant progress in theoretical understanding of DPS and DPDF:
  - evolution equations
  - spin, color and momentum correlations
  - hadron and point-like contribution to DPDF
  - tranverse momentum double distributions
  - factorization theorems to compute DPS cross sections
- Still much to be done.

3

프 + + 프 +

# Thank you!

æ

∢ ≣ ≯

≣ ▶