

CP violating Dark Matter

Venus Keus

University of Helsinki & Helsinki Institute of Physics



In collaboration with

S. F. King & S. Moretti & D. Sokolowska
J. Hernandez & D. Rojas & A. Cordero

JHEP 1612, 014 (2016) & work in progress

RISE - March 2017

- 1 Introduction and motivation
- 2 CPV DM in 3HDMs
- 3 LHC bounds
- 4 LHC signals
- 5 Summary

Dark Matter (DM)

- Cold (non-relativistic at the onset of galaxy formation)
- Non-baryonic
- Neutral and weakly interacting
 - ⇒ **Weakly Interacting Massive Particle (WIMP)**
- Stable due to a discrete symmetry

$$\underbrace{\text{DM DM} \rightarrow \text{SM SM}}_{\text{pair annihilation}}, \quad \underbrace{\text{DM} \not\rightarrow \text{SM}, \dots}_{\text{stable}}$$

- Freeze-out (drop out of thermal equilibrium)
- Agree with the observed relic density

$$\Omega_{DM} h^2 = 0.1199 \pm 0.0027$$

BSMs to the rescue

Scalar extensions with and without a Z_2 symmetry:

- Higgs portal models: SM + scalar singlet

- $\phi_{SM}, S \Rightarrow$ CPV, DM

- $\phi_{SM}, S \Rightarrow$ DM, CPV

- 2HDM: SM + scalar doublet

- Types I, II, III, IV: $\phi_1, \phi_2 \Rightarrow$ CPV, DM

- IDM - I(1+1)HDM: $\phi_1, \phi_2 \Rightarrow$ DM, CPV

- 3HDM: SM + 2 scalar doublets

- Weinberg model: $\phi_1, \phi_2, \phi_3 \Rightarrow$ CPV, DM

- I(1+2)HDM: $\phi_1, \phi_2, \phi_3 \Rightarrow$ DM, CPV

- I(2+1)HDM: $\phi_1, \phi_2, \phi_3 \Rightarrow$ CPV, DM

CP-violating DM in 3HDMs

$$\phi_1, \phi_2, \phi_3$$

$$g_{Z_2} = \text{diag}(-1, -1, +1)$$

$$VEV = (0, 0, v)$$

The scalar potential with explicit CPV

$$V_{3HDM} = V_0 + V_{Z_2}$$

$$V_0 = \sum_i^3 \left[-\mu_i^2 (\phi_i^\dagger \phi_i) + \lambda_{ii} (\phi_i^\dagger \phi_i)^2 \right]$$

$$+ \sum_{i,j}^3 \left[\lambda_{ij} (\phi_i^\dagger \phi_i) (\phi_j^\dagger \phi_j) + \lambda'_{ij} (\phi_i^\dagger \phi_j) (\phi_j^\dagger \phi_i) \right]$$

$$V_{Z_2} = -\mu_{12}^2 (\phi_1^\dagger \phi_2) + \lambda_1 (\phi_1^\dagger \phi_2)^2 + \lambda_2 (\phi_2^\dagger \phi_3)^2 + \lambda_3 (\phi_3^\dagger \phi_1)^2 + h.c.$$

$$+ \lambda_4 (\phi_3^\dagger \phi_1) (\phi_2^\dagger \phi_3) + \lambda_5 (\phi_1^\dagger \phi_2) (\phi_3^\dagger \phi_3) + \lambda_6 (\phi_1^\dagger \phi_2) (\phi_1^\dagger \phi_1)$$

$$+ \lambda_7 (\phi_1^\dagger \phi_2) (\phi_2^\dagger \phi_2) + \lambda_8 (\phi_3^\dagger \phi_1) (\phi_3^\dagger \phi_2) + h.c.$$

The Z_2 symmetry

$$\phi_1 \rightarrow -\phi_1, \quad \phi_2 \rightarrow -\phi_2, \quad \phi_3 \rightarrow \phi_3, \quad \text{SM fields} \rightarrow \text{SM fields}$$

The CP-mixed mass eigenstates

The doublet compositions

$$\phi_1 = \begin{pmatrix} H_1^+ \\ \frac{H_1^0 + iA_1^0}{\sqrt{2}} \end{pmatrix}, \quad \phi_2 = \begin{pmatrix} H_2^+ \\ \frac{H_2^0 + iA_2^0}{\sqrt{2}} \end{pmatrix}, \quad \phi_3 = \begin{pmatrix} G^+ \\ \frac{v + h + iG^0}{\sqrt{2}} \end{pmatrix}$$

The mass eigenstates

$$S_1 = \frac{\alpha H_1^0 + \alpha H_2^0 - A_1^0 + A_2^0}{\sqrt{2\alpha^2 + 2}}, \quad S_2 = \frac{-H_1^0 - H_2^0 - \alpha A_1^0 + \alpha A_2^0}{\sqrt{2\alpha^2 + 2}}$$

$$S_3 = \frac{\beta H_1^0 - \beta H_2^0 + A_1^0 + A_2^0}{\sqrt{2\beta^2 + 2}}, \quad S_4 = \frac{-H_1^0 + H_2^0 + \beta A_1^0 + \beta A_2^0}{\sqrt{2\beta^2 + 2}}$$

$$S_1^\pm = \frac{e^{\mp i\theta_{12}/2}}{\sqrt{2}} (H_2^\pm + H_1^\pm), \quad S_2^\pm = \frac{e^{\mp i\theta_{12}/2}}{\sqrt{2}} (H_2^\pm - H_1^\pm)$$

S_1 is assumed to be the DM candidate

Input parameters and constraints

DM mass m_{S_1} , Mass splittings $\delta_{S_2-S_1}$, $\delta_{S_1^\pm-S_1}$, $\delta_{S_2^\pm-S_1^\pm}$,
Higgs-DM coupling $g_{S_1 S_1 h}$, CPV phases θ_2 , θ_{12}

Constraints taken into account include:

- Stability of the potential
- Positive-definiteness of the Hessian
- Limits from gauge bosons width:
- Limits on charged scalar mass and lifetime:
- Null DM collider searches excluding simultaneously:
- S,T,U parameters

Relevant DM scenarios

In the low mass region ($m_{S_1} < m_Z$):

- **Scenario A:** no coannihilation

$$m_{S_1} \ll m_{S_2}, m_{S_3}, m_{S_4}, m_{S_1^\pm}, m_{S_2^\pm}$$

- **Scenario B:** coannihilation with S_3



$$m_{S_1} \sim m_{S_3} \ll m_{S_2}, m_{S_4}, m_{S_1^\pm}, m_{S_2^\pm}$$

- **Scenario C:** coannihilation with all neutral particles

$$m_{S_1} \sim m_{S_3} \sim m_{S_2} \sim m_{S_4} \ll m_{S_1^\pm}, m_{S_2^\pm}$$



Low DM mass region

Higgs-mediated and Z-mediated (co)annihilation




Medium DM mass region

Higgs-mediated and quartic (co)annihilation



Filling the plot

C-type scenarios are the winners!



Relevant DM scenarios

In the heavy mass region ($m_{S_1} > 400$ GeV):



- **Scenario G:** coannihilation within the "family"

$$m_{S_1} \sim m_{S_3} \sim m_{S_1^\pm} \lll m_{S_2} \sim m_{S_4} \sim m_{S_2^\pm}$$

- **Scenario H:** coannihilation with all inert particles

$$m_{S_1} \sim m_{S_3} \sim m_{S_2} \sim m_{S_4} \sim m_{S_1^\pm} \sim m_{S_2^\pm}$$

Heavy DM mass region



LHC bounds on CPV DM

Higgs invisible branching ratio and total decay

From ATLAS and CMS

$$\text{BR}(h \rightarrow \text{inv}) < 0.23 - 0.36$$

for $m_{i,j} < m_h/2$ if long lived


$$\text{BR}(h \rightarrow \text{inv}) = \frac{\sum_{i,j} \Gamma(h \rightarrow S_i S_j)}{\Gamma_h^{\text{SM}} + \sum_i \Gamma(h \rightarrow S_i S_j)}$$

The **total decay** signal strength

$$\mu_{\text{tot}} = \frac{\text{BR}(h \rightarrow \text{XX})}{\text{BR}(h_{\text{SM}} \rightarrow \text{XX})} = \frac{\Gamma_{\text{tot}}^{\text{SM}}(h)}{\Gamma_{\text{tot}}^{\text{SM}}(h) + \Gamma^{\text{inert}}(h)}$$

We use $\mu_{\text{tot}} = 1.17 \pm 0.17$ at 3σ level.

Relic density vs. Higgs decay bounds

 $g_{S_1 S_1 h}$ 

— case C1

- - - $\mu_{\min}^{\text{tot}}(h)=0.66$ — $\text{Br}(h \rightarrow \text{inv})=0.20$



$h \rightarrow \gamma\gamma$ signal strength bounds



From ATLAS and CMS: $\mu_{\gamma\gamma} = 1.16^{+0.20}_{-0.18}$

$$\mu_{\gamma\gamma} = \frac{\Gamma(h \rightarrow \gamma\gamma)^{3\text{HDM}} \Gamma(h)^{\text{SM}}}{\Gamma(h \rightarrow \gamma\gamma)^{\text{SM}} \Gamma(h)^{3\text{HDM}}}$$

Modified by

- charged scalars contribution to $\Gamma(h \rightarrow \gamma\gamma)^{3\text{HDM}}$
- light neutral scalars contribution to $\Gamma(h)^{3\text{HDM}}$


Relic density vs. $\mu_{\gamma\gamma}$ - scenario C

Relic density vs. $\mu_{\gamma\gamma}$ - scenarios G & H


LHC signatures of CPV DM

Inert cascade decays at the LHC

When there is a **large mass splitting** between DM and other inert particles:



When there is a **small mass splitting** between DM and other inert particles (winning scenarios):



$E_{miss}^T + e^+e^-$ cross section at the LHC

- Higgs-strahlung at **tree level**:

$$q\bar{q} \rightarrow Z \rightarrow S_1 S_{2,3,4} \rightarrow S_1 S_1 Z^* \rightarrow S_1 S_1 e^+ e^- \text{ with } \sigma \sim 10^{-2} \text{ pb}$$

- Higgs-strahlung at **loop level**:

$$q\bar{q} \rightarrow Z \rightarrow S_1 S_{2,3,4} \rightarrow S_1 S_1 \gamma^* \rightarrow S_1 S_1 e^+ e^- \text{ with } \sigma \sim 10^{-3} \text{ pb}$$


- Gluon-fusion at **tree level**:

$$pp \rightarrow h \rightarrow S_1 S_{2,3,4} \rightarrow S_1 S_1 Z^* \rightarrow S_1 S_1 e^+ e^- \text{ with } \sigma \sim 10^{-5} \text{ pb}$$

Other CPV observables

(JHEP1605,025(2016))

ZZZ and ZWW vertices



Summary

- CP-Violation in **I(1+2)HDM**
 - IDM-like inert sector: **CPC DM**
 - CPV in the active sector: $\tilde{H}_1, \tilde{H}_2, \tilde{H}_3$
 - Interesting LHC phenomenology, however, very limited CPV
- CP-Violation in **I(2+1)HDM**
 - SM-like active sector: $H_3 \equiv h^{SM}$
 - Unbounded CPV in the inert sector: $H_{1,2}, A_{1,2} \rightarrow S_{1,2,3,4}$ **CPV DM**
 - opens up new regions of parameter space
 - New observables at the LHC: $S_i S_j Z$ vertices

BACKUP SLIDES

Parameters of the model

- no new phenomenology from $\lambda_4, \dots, \lambda_8$ terms $\rightarrow \lambda_{4-8} = 0$
- “dark” parameters $\lambda_1, \lambda_{11}, \lambda_{22}, \lambda_{12}, \lambda'_{12}$
- “dark democracy” limit
 $\mu_1^2 = \mu_2^2, \quad \lambda_3 = \lambda_2, \quad \lambda_{31} = \lambda_{23}, \quad \lambda'_{31} = \lambda'_{23}$
- fixed by the Higgs mass $\mu_3^2 = v^2 \lambda_{33} = m_h^2/2$

7 important parameters

- CPV and mass splittings $\mu_{12}^2 = |\mu_{12}^2| e^{i\theta_{12}}, \quad \lambda_2 = |\lambda_2| e^{i\theta_2}$
- Higgs-DM coupling $\lambda_2, \lambda_{23}, \lambda'_{23}$
- Mass scale of inert particles μ_2^2


In the CPC limit

$$\alpha = \frac{-|\mu_{12}^2| \cos \theta_{12} + v^2 |\lambda_2| \cos \theta_2 - \Lambda}{|\mu_{12}^2| \sin \theta_{12} + v^2 |\lambda_2| \sin \theta_2} \rightarrow \infty$$
$$\beta = \frac{|\mu_{12}^2| \cos \theta_{12} + v^2 |\lambda_2| \cos \theta_2 - \Lambda'}{|\mu_{12}^2| \sin \theta_{12} - v^2 |\lambda_2| \sin \theta_2} \rightarrow \infty$$

where

$$\Lambda = \sqrt{v^4 |\lambda_2|^2 + |\mu_{12}^2|^2 - 2v^2 |\lambda_2| |\mu_{12}^2| \cos(\theta_{12} + \theta_2)},$$
$$\Lambda' = \sqrt{v^4 |\lambda_2|^2 + |\mu_{12}^2|^2 + 2v^2 |\lambda_2| |\mu_{12}^2| \cos(\theta_{12} + \theta_2)}.$$

Relevant DM scenarios and sum of the CPV phases



Benchmark scenarios

$$A1 : \delta_{12} = 125 \text{ GeV}, \delta_{1c} = 50 \text{ GeV}, \delta_c = 50 \text{ GeV}, \theta_2 = \theta_{12} = 1.5$$

$$B1 : \delta_{12} = 125 \text{ GeV}, \delta_{1c} = 50 \text{ GeV}, \delta_c = 50 \text{ GeV}, \theta_2 = \theta_{12} = 0.82$$

$$C1 : \delta_{12} = 12 \text{ GeV}, \delta_{1c} = 100 \text{ GeV}, \delta_c = 1 \text{ GeV}, \theta_2 = \theta_{12} = 1.57$$

$$G1 : \delta_{12} = 2 \text{ GeV}, \delta_{1c} = 1 \text{ GeV}, \delta_c = 1 \text{ GeV}, \theta_2 = \theta_{12} = 0.82$$

$$H1 : \delta_{12} = 50 \text{ GeV}, \delta_{1c} = 1 \text{ GeV}, \delta_c = 50 \text{ GeV}, \theta_2 = \theta_{12} = 0.82$$