




# Solar modulation with AMS



**Veronica Bindi, AMS Collaboration**



Physics and Astronomy Department  
University of Hawaii at Manoa  
Honolulu, Hawaii, US





# AMS on the ISS

May 19, 2011 and for the duration of the ISS.






# Particle Identification with AMS




AMS is a general purpose detector which measures particles in the GV-TV rigidity range


**TRD**  
Identify  $e^+$ ,  $e^-$




Particles and nuclei are defined by their charge ( $Z$ ) and energy ( $E \sim P$ )




**TOF**  
 $Z, V$



**Silicon Tracker**  
 $Z, P$



**Magnet**  
 $\pm Z$



**ECAL**  
 $E$  of  $e^+$ ,  $e^-$ ,  $\gamma$




**RICH**  
 $Z, V$





# Particle identification with AMS

In 6 years of operation, AMS has measured over 100 billion events.







# AMS Scientific Goals



Indirect Dark Matter search

Galactic Cosmic Rays study




M. Aguilar (AMS collaboration) PRL110,141102 (2013)

Interesting features have been measured at high energies and more has to come in the near future.


**New Goal: Study of the time variation of the low energy part of the spectrum.**



# Solar modulation: Dark Matter search



Uncertainties at low energies due to the effects of Solar modulation, increase the errors on theoretical models used for Dark Matter interpretation of the excess in the antimatter channels and in evaluation of the secondary background.



Charge sign effects affect the DM signatures expected at low energies (anti-protons, anti-deuteron).



# Solar Modulation: GCR in heliosphere



Precise measurements of the time-dependent GCRs spectra are important:

- to understand the propagation of GCRs in the heliosphere.
- to test theories of particles diffusion (charge and mass) and drift (charge-sign).
- to study the effect on cosmic rays due to the reversal in the solar polarity.



$$\frac{\partial f}{\partial t} = \nabla \cdot [K \cdot \nabla f] - V \cdot \nabla f - \langle \nu_D \rangle \cdot \nabla f + \frac{1}{3} (\nabla \cdot V) \frac{\partial f}{\partial \ln p} + Q(r, p, t)$$

CR phase space density, averaged over momentum directions

Diffusion

Convection SW

Particle Drift

Adiabatic energy losses


Local sources

Parker transport equation (1965)



# Solar activity measured by AMS


The Sun goes through an 11-year activity cycle shown by sunspots number. At each solar maximum the Sun flips its magnetic field polarity (A>0, A<0) showing a periodicity of 22 years.



The flux of galactic cosmic rays is anti-correlated with the intensity of the solar activity.



# AMS Monthly Proton Flux five years of data






# Monthly Proton & Helium Fluxes




Minimum of the flux around Feb 2014 for most of rigidities



Monthly fluxes


- 41.90-45.10 GV
- 10.10-11.00 GV
- 5.37-5.90 GV
- 3.29-3.64 GV
- 2.15-2.40 GV
- 1.51-1.71 GV
- 1.16-1.33 GV



- 41.90-45.10 GV
- 10.10-11.00 GV
- 5.37-5.90 GV
- 3.29-3.64 GV
- 2.15-2.40 GV
- 1.51-1.71 GV



# Monthly Proton & Helium Fluxes



AMS will study how solar modulation affects all different cosmic ray species.



# Electron & Positron Fluxes





Each color represents a 27 days integration flux.

## Electrons


## Positrons

Preliminary Data  
Please refer to the AMS  
forthcoming publication in PRL





# Monthly $e^+$ and $e^-$ flux Time profile







# Change of Solar Polarity and Particle Drift



[1.0 -1.2]GV



Electrons A < 0      Electrons A > 0




AMS e+/e- ratio clearly shows the charge-sign dependence of solar modulation.

Repeat the study with pbar/p.


AMS will measure the drift of positive vs negative particles during a solar minimum with positive polarity A>0.

[2.0 -2.3]GV



Polarity inversion period from: X. Sun et al., *Astroph. J.*, 798, 114 (2015)

# AMS data taking till 2024 - Agenda



**DRIFT :**  
e-, e+ , p, pbar

Comparing data collected before, during and after the solar maximum AMS can study in details the effects due to the solar polarity reversal.

AMS will measure the drift of positive vs negative particles during the solar minimum with positive polarity  $A > 0$ .

AMS will compare solar maximum of cycles 25 ( $A > 0$ ) and 24 ( $A < 0$ ) and the solar polarity reversal.

**DIFFUSION :**  
All species of GCR

AMS is measuring the modulation of different nuclei during a solar cycle.

**FD and SEP :**

Measured 27 SEPs and identified so far 33 FDs.

Not expected major events during minimum

Lots of new SEPs and FDs.



# Summary & Conclusions

A new era in galactic cosmic rays understanding has started, not only at high energy, but also at low energy in the region affected by the solar modulation thanks to the precise and continuous observations from space by PAMELA in solar cycle 23 and now AMS-02 in solar cycle 24 and 25.

New and precise measurements are increasing our knowledge of important effects such as diffusion and drift in the heliosphere, allowing the detailed study of propagation.

These measurements will serve as a high-precision baseline for continued studies of GCR solar modulation, SEPs, space radiation hazards, magnetospheric effects, trapped particles and in many other fields.

**Near future forthcoming AMS publications on solar modulation: time evolution of proton and helium fluxes, electron and positron fluxes, and antiproton/proton ratio.**