

Contribution ID: 35

Type: not specified

D meson nuclear modification factor and v_n harmonics in PbPb collisions at 5.02 TeV with CMS

Wednesday 1 November 2017 11:45 (35 minutes)

The measurement of heavy flavour production is a powerful tool to study the properties of the high-density QCD medium created in heavy-ion collisions as heavy quarks are sensitive to the transport properties of the medium and may interact with the QCD matter differently from light quarks. In particular, the comparison between the nuclear modification factors (RAA) of light- and heavy-flavour particles provides insights into the expected flavour dependence of in-medium parton energy loss. Furthermore, azimuthal anisotropy coefficient (vn) of heavy-flavor particles provide insights into the degree of the thermalization of the bulk medium at low pT, and unique information about the path length dependence of heavy quark energy loss at high pT. Using the large statistics proton-proton and PbPb samples collected at 5.02 TeV during the 2015 LHC run, high precision open charm measurements are performed with the CMS detector in a wide transverse momentum range. This allows us to set an important milestone in our understanding of the interactions between heavy quarks and the medium. In this talk, the most recent results of RAA, v2 and v3 of D0 mesons in PbPb collisions at 5.02 TeV are presented and compared to the same results for charged hadrons at the same energy.

Primary author: WANG, Jing (Massachusetts Inst. of Technology (US))Presenter: WANG, Jing (Massachusetts Inst. of Technology (US))Session Classification: Heavy flavor and Quarkonia