



Contribution ID: 64

Type: **not specified**

## Heavy Photon Search

Compelling motivations, including promising classes of dark matter models compatible with various observed astrophysical anomalies, motivate the existence of a massive gauge boson carrier of a  $U'(1)$  gauge symmetry beyond the Standard Model. This hypothesized “heavy photon” (abbreviated  $A'$ ) would interact feebly with the SM, through kinetic mixing with the SM photon. Our experiment seeks to produce the  $A'$  in a laboratory setting by striking a tungsten foil target with a continuous electron beam. An  $A'$  may be produced in a process analogous to bremsstrahlung, and then decay into an  $e^+e^-$  pair. Our detector setup consists of a silicon vertex tracker surrounded by a large magnet for tracking the pair, and an electronic calorimeter for triggering. To find the  $A'$  signal amongst the much larger QED background, we use resonance search (bump-hunt) and displaced-vertex search techniques. We have taken data with two beam energies: 1.05 GeV during our 2015 engineering run, and 2.3 GeV in our 2016 production run. This poster describes the experimental setup and presents preliminary results for the two datasets, as well as projected reach for future running.

**Authors:** PAUL, Sebouh; DIAMOND, Miriam Deborah Joy (University of Toronto (CA))

**Presenters:** PAUL, Sebouh; DIAMOND, Miriam Deborah Joy (University of Toronto (CA))

**Session Classification:** Poster social