Physics Motivation For Future Colliders

Hep 2018, UTFSM, Jan. 12, 2018

Tao Han, University of Pittsburgh

TLEP Report 1308.6176; CEPS pre-CDR; Snowmass Reports;
N. Arkani-Hamed, T. Han, M. Mangano, L.T. Wang, Phys. Rep. 2017;
More refs: CERN Yellow Reports, 2017.

High Energy Physics IS at an extremely interesting time:

The completion of the SM:

First time ever, we have a consistent relativistic/quantum mechanical theory: weakly coupled, unitary, renormalizable, vacuum-(quasi?)stable.

Valid up to an exponentially high scale, perhaps to the Planck scale M_{Pl}!

“... most of the grand underlying principles have been firmly established. (An eminent physicist remarked that) the future truths of physical science are to be looked for in the sixth place of decimals. ”

"... most of the grand underlying principles have been firmly established. (An eminent physicist remarked that) the future truths of physical science are to be looked for in the sixth place of decimals. "

--- Albert Michelson (1894)

Michelson–Morley experiments (1887):
"the moving-off point for the theoretical aspects of the second scientific revolution"

Will History repeat itself (soon)?

New Era:
Under the Higgs lamp post

Question 1: The Nature of EWSB ?

In the SM, $m_H^2 = 2\mu^2 = 2\lambda v^2 \quad \Rightarrow \quad \mu \approx 89 \text{ GeV}, \quad \lambda \approx \frac{1}{8}.$

underwent a 2nd order phase transition (?)



Question 1: The Nature of EWSB ?

In the SM, $m_H^2 = 2\mu^2 = 2\lambda v^2 \quad \Rightarrow \quad \mu \approx 89 \text{ GeV}, \quad \lambda \approx \frac{1}{8}.$

underwent a 2nd order phase transition (?)



With new physics near the EW scale:

$$V(h) \to m_h^2(h^\dagger h) + \frac{1}{2}\lambda(h^\dagger h)^2 + \frac{1}{3!\Lambda^2}(h^\dagger h)^3, \rightarrow \lambda_{hhh} = (7/3)\lambda_{hhh}^{SM}$$

$$\to \frac{1}{2}\lambda(h^\dagger h)^2 \log\left[\frac{(h^\dagger h)}{m^2}\right] \rightarrow \lambda_{hhh} = (5/3)\lambda_{hhh}^{SM}$$

Question 1: The Nature of EWSB ?

In the SM, $m_H^2 = 2\mu^2 = 2\lambda v^2 \quad \Rightarrow \quad \mu \approx 89\ \text{GeV}, \quad \lambda \approx \frac{1}{8}.$

underwent a 2nd order phase transition (?)



With new physics near the EW scale:

$$V(h) \to m_h^2(h^\dagger h) + \frac{1}{2}\lambda(h^\dagger h)^2 + \frac{1}{3!\Lambda^2}(h^\dagger h)^3, \rightarrow \lambda_{hhh} = (7/3)\lambda_{hhh}^{SM}$$

$$\to \frac{1}{2}\lambda(h^\dagger h)^2 \log\left[\frac{(h^\dagger h)}{m^2}\right] \rightarrow \lambda_{hhh} = (5/3)\lambda_{hhh}^{SM}$$

EW phase transition strong 1st order!

→ O(1) deviation on λ_{hhh}

X.M.Zhang (1993); C. Grojean et al. (2005)

Question 2: The "Naturalness"

"… scalar particles are the only kind of free particles whose mass term does not break either an internal or a gauge symmetry."

Ken Wilson, 1970

Question 2: The "Naturalness"

"… scalar particles are the only kind of free particles whose mass term does not break either an internal or a gauge symmetry."

Ken Wilson, 1970

"unnatural" in the 't Hooft sense:



$$m_H^2 = m_{H0}^2 \quad - \frac{3}{8\pi^2} y_t^2 \Lambda^2 \qquad + \frac{1}{16\pi^2} g^2 \Lambda^2 \qquad + \frac{1}{16\pi^2} \lambda^2 \Lambda^2$$

If $\Lambda^2 \gg m_H^2$, then unnaturally large cancellations must occur.

Question 2: The "Naturalness"

"… scalar particles are the only kind of free particles whose mass term does not break either an internal or a gauge symmetry."

Ken Wilson, 1970

"unnatural" in the 't Hooft sense:



$$m_H^2 = m_{H0}^2 \quad - \frac{3}{8\pi^2} y_t^2 \Lambda^2 \qquad + \frac{1}{16\pi^2} g^2 \Lambda^2 \qquad + \frac{1}{16\pi^2} \lambda^2 \Lambda^2$$

If $\Lambda^2 \gg m_H^2$, then unnaturally large cancellations must occur.

- Natural: O(1 TeV) new physics, associated with ttH.
- Unknown: Deep UV-IR correlations: gravity at UV?
- Agnostic: Multiverse/anthropic?

A "NATURAL" EW THEORY?

- "Natural SUSY":

Relevant to the Higgs and the "Most Wanted":

$\tilde{H}^{0,\pm},\ \tilde{t},\ \tilde{b},\ (\tilde{g});\ S,\ \tilde{S}...$

LHC Run 2 bounds:

$$m_{\tilde{t}} > 800 - 1100 \text{ GeV}$$

$$m_{\tilde{\chi}^\pm} > 600 - 1100 \text{ GeV}$$



A "NATURAL" EW THEORY?

- "Natural SUSY":

Relevant to the Higgs and the "Most Wanted":

$\tilde{H}^{0,\pm},\ \tilde{t},\ \tilde{b},\ (\tilde{g});\ S,\ \tilde{S}...$

LHC Run 2 bounds:

$m_{\tilde{t}} > 800 - 1100$ GeV

$m_{\tilde{\chi}^\pm} > 600 - 1100$ GeV



- New strong dynamics, "Compositeness":

The top-quark partner T',

Current ATLAS/CMS limit:

$M_T > 1400$ GeV, for $M_A < 100$ GeV.

Question 3: The Dark Sector

The un-protected operator may reveal secret

Higgs portal: $k_s H^\dagger H \; S^* S, \quad \frac{k_\chi}{\Lambda} H^\dagger H \; \bar{\chi}\chi.$





Question 4: The "Flavor Puzzle"

- Particle mass hierarchy
- Patterns of quark, neutrino mixings
- New CP-violation sources?

Higgs Yukawa couplings as the pivot!



Nature News, July '14

COLLISION COURSE

Particle physicists around the world are designing colliders that are much larger in size than the Large Hadron Collider at CERN, Europe's particle-physics laboratory.

LHC Leads the Way (2015-2030)



Nature News, July '14

COLLISION COURSE

Particle physicists around the world are designing colliders that are much larger in size than the Large Hadron Collider at CERN, Europe's particle-physics laboratory.



Nature News, July '14

COLLISION COURSE

Particle physicists around the world are designing colliders that are much larger in size than the Large Hadron Collider at CERN, Europe's particle-physics laboratory.

LHC Leads the Way (2015-2030)

CERN's Large Hadron Collider
Circumference: 27 km
Energy: 14 TeV

China's electron–positron collider
52 km; 240 GeV
China's super proton collider
52 km; ≤70 TeV

US/European super proton collider
100 km; 100 TeV
e+e-&Z,240-350GeV

CEPC/SppC?

China-hosted international electron–positron collider
80 km; 240 GeV
China-hosted international super proton collider
80 km; ≤100 TeV

FCC?

International Linear Collider
Length: 31 km
≤1 TeV

Existing Proposed

GeV, gigaelectronvolt

ILC as Higgs Factory & beyond



Nature News, July '14

COLLISION COURSE

Particle physicists around the world are designing colliders that are much larger in size than the Large Hadron Collider at CERN, Europe's particle-physics laboratory.

LHC Leads the Way (2015-2030)



CEPC/SppC?

FCC?

ILC as Higgs Factory & beyond

Table 1-1. *Proposed running periods and integrated luminosities at each of the center-of-mass energies for each facility.*

Snowmass 1310.8361

Facility	HL-LHC	ILC	ILC(LumiUp)	CLIC	TLEP (4 IPs)	HE-LHC	VLHC
$\overline{s}$ (GeV)	14,000	250/500/1000	250/500/1000	350/1400/3000	240/350	33,000	100,000
$\mathcal{L}dt$ (fb^{-1})	3000/expt	250+500+1000	1150+1600+2500	500+1500+2000	10,000+2600	3000	3000
dt (10^7s)	6	3+3+3	(ILC 3+3+3) + 3+3+3	3.1+4+3.3	5+5	6	6

International Linear Collider as a Higgs Factory & beyond

Ecm (GeV) = 250, 350, 500, 1000 GeV

Under serious consideration in Japan:



International Linear Collider as a Higgs Factory & beyond

Ecm (GeV) = 250, 350, 500, 1000 GeV

Under serious consideration in Japan:







Linear Collider Case study working group, K. Fujii et al. :
Stage studies: 1710.07261; 1801.02841

Future Circular Collider Study
CDR for European Strategy Update 2019/20

international FCC collaboration (CERN as host lab) to design:

- ***pp*-collider (*FCC-hh*)**
  → main emphasis, defining infrastructure requirements

~16 T ⇒ 100 TeV *pp* in 100 km

- **80-100 km tunnel infrastructure** in Geneva area, site specific
- ***e⁺e⁻* collider (*FCC-ee*),** as a possible first step
- ***p-e* (*FCC-he*) option,** one IP, FCC-hh & ERL
- **HE-LHC** w *FCC-hh* technology



M. Benedikt, F. Zimmermann, 2017

FCC Timeline



M. Benedikt

CERN Yellow Reports on "a 100 TeV pp Collider":
Vol. 1. SM; 2. Higgs; 3. BSM; 4. Accelerator

Fastest Possible Technical Schedules



technical schedule defined by magnets program and by CE
→ earliest possible physics starting dates:

- FCC-hh: 2043
- FCC-ee: 2039
- HE-LHC: 2040 (with HL-LHC stop at LS5 / 2034)

HE-LHC
design & construction

M. Benedikt

CEPC (circular e^-e^+)/SppC: China



CEPC-SPPC Timeline



Y. Wang

CEPC/SppC Preliminary Conceptual Design Reports:
Vol. 1: Physics & Detector; Vol. 2: Accelerator
http://cepc.ihep.ac.cn/preCDR/volume.html

e^+e^- colliders: Energy/Lumi projection



TLEP Report: 1308.6176

E_{cm}	running time	statistics (FCC-ee)
	b,c,τ	10^{11} b,c,τ
90 GeV	1-2 yrs	10^{12} Z (Tera Z)
160 GeV	1-2 yrs	10^8- 10^9 WW(Oku W)
240 GeV	4-5 yrs	$2x10^6$ ZH (Mega H)
350 GeV	4-5 yrs	10^6 tt (Mega top)

Z-Factory: Tera (10^{12}) Z Physics

TLEP Report: 1308.6176

- Clean environment, $\Delta Ecm < 1$ MeV, 10^5 x LEP-I
- possible longitudinal polarization
- Z-ploe: ΔM_Z, $\Delta\Gamma_Z < 0.1$ MeV, $\Delta\sin^2\theta_w < 10^{-6}$;
- Thr. scan: $\Delta M_W \sim O(1\ \text{MeV})$, $\Delta m_t \sim O(10\ \text{MeV})$, $\Delta m_H \sim O(10\ \text{MeV})$.



Higgs-Factory: Mega (10^6) Higgs

TLEP Report: 1308.6176



Higgs-Factory: Mega (10^6) Higgs

TLEP Report: 1308.6176



"Recoil mass"

$$m_h^2 = (p_{e^-} + p_{e^+} - q_{\mu^-} - q_{\mu^+})^2$$



Higgs-Factory:

- ILC: E_{cm} = 250 (500) GeV, 250 (500) fb^{-1} (0.5×10^5 Higgs)

Model-independent measurement:

ILC Report: 1308.6176

$\Gamma_H \sim 6\%$, $\Delta m_H \sim 30$ MeV

(HL-LHC: assume SM, $\Gamma_H \sim 5\text{-}8\%$, $\Delta m_H \sim 50$ MeV)

- FCCee/CEPC 10^6 Higgs: $\Gamma_H \sim 1\%$, $\Delta m_H \sim 5$ MeV.

Higgs-Factory:

- ILC: E_{cm} = 250 (500) GeV, 250 (500) fb^{-1} (0.5×10^5 Higgs)

Model-independent measurement:

ILC Report: 1308.6176

$\Gamma_H \sim 6\%$, $\Delta m_H \sim 30$ MeV

(HL-LHC: assume SM, $\Gamma_H \sim 5\text{-}8\%$, $\Delta m_H \sim 50$ MeV)

- FCC/CEPC 10^6 Higgs: $\Gamma_H \sim 1\%$, $\Delta m_H \sim 5$ MeV.

Couplings to sub-percent:



The Next Energy Frontier: 100 TeV Hadron Collider







Higgs Production @ FCC_{hh}/SPPC



Process	σ (100 TeV)/σ (14 TeV)
Total pp	1.25
W	~7
Z	~7
WW	~10
ZZ	~10
tt	~30
H	~15 (ttH ~60)
HH	~40
stop (m=1 TeV)	~10^3

Snowmass QCD Working Group: 1310.5189

Higgs Self-couplings:

$$\mathcal{L} = -\frac{1}{2}m_H^2 H^2 - \frac{g_{HHH}}{3!}H^3 - \frac{g_{HHHH}}{4!}H^4$$

$$g_{HHH} = 6 \quad v = \frac{3m_H^2}{v}, \qquad g_{HHHH} = 6 \quad = \frac{3m_H^2}{v^2}.$$



Triple Higgs boson coupling λ_{hhh} :

Test the shape of the Higgs potential,
and the fate of the EW-phase transition (EWPT):
O(100%) deviation needed for 1st order EWPT;
O(10%) accuracy needed for a conclusive test.

Higgs Self-couplings:

$$\mathcal{L} = -\frac{1}{2}m_H^2 H^2 - \frac{g_{HHH}}{3!}H^3 - \frac{g_{HHHH}}{4!}H^4$$

$$g_{HHH} = 6 \quad v = \frac{3m_H^2}{v}, \qquad g_{HHHH} = 6 \quad = \frac{3m_H^2}{v^2}.$$



Triple Higgs boson coupling λ_{hhh} :

Test the shape of the Higgs potential,
and the fate of the EW-phase transition (EWPT):
O(100%) deviation needed for 1st order EWPT;
O(10%) accuracy needed for a conclusive test.

HL-LHC ~ 50%; ILC(1 TeV), CLIC(3 TeV) ~ 10%;

FCC_{hh} @ 3 ab^{-1}: ~ 8%

SUSY @ FCC_{hh}/SPPC

M.Mangano et al.: 1407.5066







Mass reach at 100 TeV:
~ 7x over LHC

Pushing the "Naturalness" limit



The Higgs mass fine-tune: $\delta m_H/m_H \sim 1\%\ (1\ \text{TeV}/\Lambda)^2$

Thus, $m_{stop} > 8$ TeV → 10^{-4} fine-tune!

DM Searches



GeV low mass:
DD difficult;
Collider complementary

100 GeV or higher mass:
DD + ID + HE Collider

WIMP DM: $M_{\rm DM} < 1.8\ {\rm TeV}\left(\frac{g^2_{\rm eff}}{0.3}\right)$









New Particle Searches

Electroweak Resonances: Z',W'



~ 6x over LHC

Colored Resonances:



Conclusions



- Higgs boson is a new class.

NP BSM → "under the Higgs lamppost"

LHC will lead the way: $g \sim 10\%$; $\lambda_{HHH} \sim 50\%$; $Br_{inv.} \sim 20\%$

but it also calls for new colliders:

Precision: FCC_{ee}/CEPC

Tera Z: ΔM_Z, $\Delta\Gamma_Z < 0.1$ MeV, $\Delta\sin^2\theta_w < 10^{-6}$.

At thresholds: $\Delta M_W \sim 1$ MeV, $\Delta m_t \sim 10$ MeV

Mega Higgs: $\kappa_v \sim 0.2\%$, $\Gamma_H \sim 1\%$, $Br_{inv.} \sim 1\%$, $\Delta m_H \sim 5$ MeV.

Conclusions



- Higgs boson is a new class.

NP BSM → "under the Higgs lamppost"

LHC will lead the way: $g \sim 10\%$; $\lambda_{HHH} \sim 50\%$; $Br_{inv.} \sim 20\%$

but it also calls for new colliders:

Precision: FCC_{ee}/CEPC

Tera Z: ΔM_Z, $\Delta\Gamma_Z < 0.1$ MeV, $\Delta\sin^2\theta_w < 10^{-6}$.

At thresholds: $\Delta M_W \sim 1$ MeV, $\Delta m_t \sim 10$ MeV

Mega Higgs: $\kappa_v \sim 0.2\%$, $\Gamma_H \sim 1\%$, $Br_{inv.} \sim 1\%$, $\Delta m_H \sim 5$ MeV.

Energy frontier: FCC_{hh}/SPPC

$g \sim 10\%$; $\lambda_{hhh} < 10\%$ → Conclusive for EWPT

6x LHC reach: 10 – 30 TeV → fine-tune $< 10^{-4}$

WIPM DM mass ~ 1 – 5 TeV

Conclusions



- Higgs boson is a new class.

NP BSM → "under the Higgs lamppost"

LHC will lead the way: $g \sim 10\%$; $\lambda_{HHH} \sim 50\%$; $Br_{inv.} \sim 20\%$

but it also calls for new colliders:

Precision: FCC_{ee}/CEPC

Tera Z: ΔM_Z, $\Delta\Gamma_Z < 0.1$ MeV, $\Delta\sin^2\theta_w < 10^{-6}$.

At thresholds: $\Delta M_W \sim 1$ MeV, $\Delta m_t \sim 10$ MeV

Mega Higgs: $\kappa_v \sim 0.2\%$, $\Gamma_H \sim 1\%$, $Br_{inv.} \sim 1\%$, $\Delta m_H \sim 5$ MeV.

Energy frontier: FCC_{hh}/SPPC

$g \sim 10\%$; $\lambda_{hhh} < 10\%$ → Conclusive for EWPT

6x LHC reach: 10 – 30 TeV → fine-tune $< 10^{-4}$

WIPM DM mass ~ 1 – 5 TeV

An exciting journey ahead!

Site selections (a few main candidates)

1)



2)

3)

1) Qinhuangdao

2) Shaanxi Province

3) Near Shenzhen and Hongkong

"Canonical" energy / luminosity:

100 TeV, 3 – 30 ab^{-1}

(Perhaps)

- Technology limitation (high field magnets?)
- Budgetary consideration (> 10 B$?)
- Geological / geographic consideration

Atomic physics:

Rydberg const. $E_0 \sim \alpha^2 m_e \rightarrow O(25\ \text{eV})$, very natural!



Nuclear physics?



Solar eclipses:



$\delta\theta/\theta \sim 10^{-2}$

rather unnatural!