

$\mu - \tau$ reflection symmetry with a texture zero

Celso C. Nishi ¹



Universidade Federal do ABC
Santo André, SP, Brazil

7th International Conference on
High Energy Physics in the LHC Era
Valparaiso, 8-12 Jan 2018


¹celso.nishi@ufabc.edu.br

The flavor problem: leptonic mixing

Neutrinos oscillate





'15 Kajita & McDonald



2012 $\theta_{13} \approx 8.5^\circ \rightarrow$ **CP**
 Daya-Bay, RENO


$$|V_{\text{PMNS}}| = \begin{pmatrix} 0.82 & 0.55 & 0.15 \\ 0.35 & 0.71 & 0.61 \\ 0.45 & 0.45 & 0.77 \end{pmatrix}$$




Strumia, Vissani, hep-ph/0606054




The flavor problem: neutrino masses




- If lepton number is broken at a scale $\Lambda \gg v$



- $m_\nu \lesssim 0.1\text{ eV}$ for $\Lambda \gtrsim 10^{12}\text{ GeV}$
- The seesaw mechanism
- Type I: heavy singlets N_1, N_2, N_3



Maximal atmospheric angle and Dirac phase?



Capozzi, et al., NPB'16.

$CP^{\mu\tau}$ symmetry or $\mu\tau$ reflection

$$\nu_e \rightarrow \nu_e^{\text{CP}}, \quad \nu_\mu \rightarrow \nu_\tau^{\text{CP}}, \quad \nu_\tau \rightarrow \nu_\mu^{\text{CP}}$$

$\theta_{23} = 45^\circ$, $\delta_{\text{CP}} = \pm 90^\circ$, **trivial** Majorana phases, θ_{13}, θ_{12} free

• $CP^{\mu\tau} = \text{CP}$ with $\mu - \tau$ interchange

• Also known as $\mu - \tau$ reflection

Harrison, Scott, PLB'02

• **Badly broken** in the charged lepton sector: $y_\mu \ll y_\tau$


• Hint from $|V_{\mu i}| \approx |V_{\tau i}|$

• **Predict all CP phases still allowing ~~CP~~**

• **Accidental maximality**

He, Rodejohann, Xu, PLB'15
Joshi-pura & Patel, PLB'15


CP^{μτ} and neutrinoless double beta decay




generic $m_{ee} = m_1 U_{e1}^2 + m_2 U_{e2}^2 + m_3 U_{e3}^2$

CP^{μτ} $m_{ee} = \pm m_1 |U_{e1}|^2 \pm m_2 |U_{e2}|^2 \pm m_3 |U_{e3}|^2$

four CP parities



PDG




Mohapatra, Nishi, JHEP'15



Baryon asymmetry of the Universe

Baryogenesis mechanism
needed at $T \gtrsim 40 \text{ MeV}$



- $\frac{n_B - n_{\bar{B}}}{n_\gamma} \sim 10^{-9}$
- Sakharov conditions '67
 - violation of CP, C
 - violation of B
 - departure of thermal equilibrium
- SM is not enough


Leptogenesis

Matter-antimatter asymmetry


$$Y_{\Delta B} = \frac{n_B - n_{\bar{B}}}{s} \sim -10^{-3} \times \epsilon \times \eta$$

Fukugita, Yanagida, '86

- $Y_B \Big|_{\text{exp}} = (8.65 \pm 0.09) \times 10^{-11}$
- N_1 decay generates L asymmetry




- Spharelon processes $\Delta L \rightarrow \Delta B$
 $\Delta B = \Delta L = 3$
 Klinkhamer, Manton, '84



Cline, hep-ph/0609145

CP^{μτ} and leptogenesis



Garbrecht, Schwaller, JCAP'14

Washout in different flavors can be different

Abada et al., JCAP'06

Nardi et al. JHEP'06

- **Unflavored** leptogenesis does not work Grimus & Lavoura, PLB'04

- For $T \sim M_1 > 10^{12} \text{ GeV}$: $Y_{\Delta B} \sim -10^{-3} \times \eta \times [\epsilon_\tau + \epsilon_\mu + \epsilon_e]$

- With CP^{μτ}: $\epsilon_e = 0, \epsilon_\mu + \epsilon_\tau = 0 \implies \epsilon_e + \epsilon_\mu + \epsilon_\tau = 0$

Purely flavored scenario

- **Flavored** leptogenesis works for intermediate scales

h_τ interactions fast

Mohapatra, Nishi, JHEP'15

$$10^9 \text{ GeV} \lesssim T \sim M_1 \lesssim 10^{12} \text{ GeV}$$

Increasing predictivity with texture-zeros

- Flavor symmetries increase predictivity
 - Nonabelian: relate entries and/or fix mixing angles
 - Abelian: vanishing entries = texture-zeroes

- Texture zeroes in the lepton sector

Frampton,Glashow,Marfatia,PLB'02

- Can be always enforced by abelian symmetries

Grimus,Joshipura,Lavoura,Tanimoto,EPJC'04

- $CP^{\mu\tau}$ corresponds to the first case
- With only $CP^{\mu\tau}$ there is no sharp prediction besides maximal θ_{23} , δ_{CP} and trivial Majorana phases
- Can we enforce an additional texture-zero on top of $CP^{\mu\tau}$?



Increasing predictivity with texture-zeros

- $CP^{\mu\tau}$ enforces (flavor basis)

$$M_\nu = \begin{pmatrix} a & d & d^* \\ d & c & b \\ d^* & b & c^* \end{pmatrix}, \quad \text{with real } a, b \text{ and } \text{Im}(d^2 c^*) \neq 0.$$

- 5 parameters $a, b, \text{Re } c, \text{Im } c, |d|$
to describe $m_1, m_2, m_3, \theta_{12}, \theta_{13}$ $\theta_{23}, \delta, \alpha, \beta$ fixed by $CP^{\mu\tau}$
- Can we enforce an additional texture-zero on top of this structure?
- Only $a = 0$ or $b = 0$ are phenomenologically viable
- With one less parameters we can predict the lightest neutrino mass!

$CP^{\mu\tau}$ and \mathbb{Z}_8

Nishi, Sánchez-Vega, JHEP 1701 (2017) 068

- Symmetries in (e, μ, τ) and $(\nu_e, \nu_\mu, \nu_\tau)$

$$\mathbb{Z}_8 : T = \begin{pmatrix} -1 & & \\ & \omega_8 & \\ & & \omega_8^3 \end{pmatrix}, \quad \omega_8 = e^{i2\pi/8}$$

$$CP^{\mu\tau} : \nu_{\alpha L} \rightarrow X_{\alpha\beta} \nu_{\beta L}^{cp}, \quad X = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

- They are consistent $T \rightarrow XT^*X^{-1} = T^5$
- Extends the simplest case of $L_\mu - L_\tau$ with $CP^{\mu\tau}$

Mohapatra, Nishi, JHEP15



CP^{μτ} and ℤ₈

- Different entries have different ℤ₈ charges

$$\bar{\nu}_{\alpha L}^c \nu_{\beta L} \sim \begin{pmatrix} 1 & \omega_8^5 & \omega_8^{-1} \\ \star & \omega_8^2 & -1 \\ \star & \star & \omega_8^{-2} \end{pmatrix}, \quad \omega_8 = e^{i2\pi/8}$$

- Couple with scalars of different charges $\eta_k \sim \omega_8^k$
- Additional ℤ₄^{B-L} symmetry: $L_\alpha \sim -i$ and $\eta_k \sim -1$

$$\begin{aligned} & \frac{1}{2} \frac{C_{ee}}{\Lambda^2} \eta_0 L_e H L_e H + \frac{1}{2} \frac{C_{\mu\mu}}{\Lambda^2} \eta_2^* L_\mu H L_\mu H + \frac{1}{2} \frac{C_{\tau\tau}}{\Lambda^2} \eta_2 L_\tau H L_\tau H \\ & + \frac{C_{\mu\tau}}{\Lambda^2} \eta_4 L_\mu H L_\tau H + \frac{C_{e\mu}}{\Lambda^2} \eta_3 L_e H L_\mu H + \frac{C_{e\tau}}{\Lambda^2} \eta_1 L_e H L_\tau H \end{aligned}$$

$$C_{e\tau} = C_{e\mu}^*, C_{\tau\tau} = C_{\mu\mu}^* \text{ and real } C_{ee}, C_{\mu\tau}$$


- Texture-zeros: $a = 0$ if η_0 absent $b = 0$ if η_4 absent



CP ^{$\mu\tau$} and \mathbb{Z}_8

Nishi, Sánchez-Vega, JHEP 1701 (2017) 068


Case	$(M_\nu)_{\alpha\beta}=0$	ordering	CP parities	m_0	$m_{\beta\beta}$	$\sum m_\nu$
I	(ee)	NO	(-++)	4.4 – 9.0	0	63 – 74
II	(ee)	NO	(+ - +)	1.1 – 3.9	0	59 – 65
III	($\mu\tau$)	NO	(++-)	151 – 185	142 – 178	460 – 561
IV	($\mu\tau$)	IO	(+ - +)	15 – 30	14.3 – 29.3	116 – 148



Conclusions

- $\text{CP}^{\mu\tau}$ is a viable symmetry that predicts all CP phases and it is compatible with flavored leptogenesis.
- Maximal θ_{23} and δ_{CP} will be tested in the future
- We have shown by explicit construction a highly predictive scenario where the neutrino mass matrix is symmetric by $\text{CP}^{\mu\tau}$ and *additionally* contains one texture-zero in the (ee) or $(\mu\tau)$ entry.
- The possibility of a neutrino mass matrix with $\text{CP}^{\mu\tau}$ symmetry *simultaneously* with a texture-zero that is enforced by symmetry was first shown here and it is only allowed by combining in a non-usual way a discrete abelian symmetry at least as large as \mathbb{Z}_8 and $\text{CP}^{\mu\tau}$.
- Distinct predictions for the lightest neutrino mass and m_{ee} ; in one case, testable in the near future.

Leptogenesis and $\mu\tau$ reflection



Mohapatra, Nishi, JHEP15

$$M_3 \rightarrow \infty$$

$$M_1 = 10^{12} \text{ GeV}$$

- Leptogenesis is only **possible** for $10^9 \text{ GeV} \lesssim M_1 \lesssim 10^{12} \text{ GeV}$
- For $M_1 \gtrsim 10^{12} \text{ GeV}$, we cannot distinguish any flavor and $Y_B \approx 0$
- For $M_1 \lesssim 10^9 \text{ GeV}$, we can distinguish e, μ, τ and $Y_B \approx 0$