

Contribution ID: 40

Type: Talk

The strong coupling from $e^+e^- \rightarrow$ hadrons

We use a new compilation of the hadronic R-ratio from available data for the process $e^+e^- \rightarrow$ hadrons below the charm mass to determine the strong coupling α_s , using finite-energy sum rules. Quoting our results at the τ mass to facilitate comparison to the results obtained from similar analyses of hadronic τ -decay data, we find $\alpha_s(m_\tau^2) = 0.298 \pm 0.016 \pm 0.006$ in fixed-order perturbation theory, and $\alpha_s(m_\tau^2) = 0.304 \pm 0.018 \pm 0.006$ in contour-improved perturbation theory, where the first error is statistical, and the second error combines various systematic effects. These values are in good agreement with a recent determination from the OPAL and ALEPH data for hadronic τ decays.

Primary author: GOLTERMAN, Maarten (San Francisco State University)

Co-authors: BOITO, Diogo (Universidade de São Paulo); Dr KESHAVARZI, Alexander (University of Liverpool); MALTMAN, Kim (York University); NOMURA, Daisuke; PERIS, Santiago (Univ. Autonoma de Barcelona); TEUB-NER, Thomas (University of Liverpool)

Presenter: GOLTERMAN, Maarten (San Francisco State University)

Track Classification: Tau properties/QCD