Hard Probes 2018: International Conference on Hard & Electromagnetic Probes of High-Energy Nuclear Collisions

Contribution ID: 168

Type: 1c) Initial State (POSTER FROM TALK)

## Hydrodynamic collectivity in Pb + Pb and proton + proton collisions

In this talk, we study the hydrodynamic collectivity in Pb + Pb collisions at 2.76 TeV and 5.02 TeV and proton + proton collisions at 13 TeV using the iEBE-VISHNU hybrid model. For Pb + Pb collision systems, we use TRENTo and AMPT initial conditions. With properly chosen and turned parameter sets, our model calculations can nicely describe various flow obsevables in 2.76 TeV and 5.02 TeV Pb + Pb collisions, including 2- and 4- particle cumulant, differential flow, event-by-event distribution, mode-coupling effects. For proton + proton systems, with properly tuned parameters, iEBE-VISHNU hybrid model with HIJING initial condition can describe the measured 2-particle correlations, including integrated and differential elliptic flow and all charged and identified hadrons ( $K_s^0$ ,  $\Lambda$ ). However, our model calculations shows positive 4-particle cumulant to simultaneously describe the 2- and 4- particle cumulant, it is required to have singnificant improvements on initial condition for p-p collisions.

## **Summary**

Our work focus on the hydrodynamic behavior in proton + proton system, finding the current initial conditions can describe the 4-particle cumulant within the viscous hydro framework, and require to improve the descriptions of the initial state in proton + proton collisions.

**Primary authors:** ZHAO, Wenbin (Peking University); SONG, Huichao; ZHOU, You (University of Arizona (US)); Dr XU, Haojie (Huzhou University)

Presenter: ZHAO, Wenbin (Peking University)

Session Classification: Poster Session