

HARD PROBES 2018

Probing Heavy Ion Collisions Using Quark and Gluon Jet Substructure

Yang-Ting Chien

LHC Theory Initiative Fellow, MIT Center for Theoretical Physics

In collaboration with Raghav Kunnawalkam Elayavalli

arXiv:1803.03589



October 3, 2018, Aix les Bains





Outline



- ▶ Jet substructure: hard and soft probes at all energy scales
- ▶ Quark and gluon jets as two different probes
- ▶ Jet representation and analysis
 - ▶ physics-motivated multivariate analysis: constructive
 - ▶ unbiased machine-learning features: comprehensive
- ▶ Telescoping deconstruction: a complete jet observable basis
- ▶ Conclusion

The era of precision jet substructure studies




SCET Chien-Vitev JHEP05(2016)023



Relate precise jet modifications to medium properties




Quark jets and gluon jets




- ▶ All the jets are a mixture of quark jets and gluon jets
- ▶ Quark and gluon jets have different color charges and substructure
- ▶ Quark/gluon jet fraction affects jet substructure
 - ▶ 40% quark 60% gluon \rightarrow 60% quark 40% gluon: jets become more quark-jet like
- ▶ Substructure of each jet is modified
 - ▶ Quark jets and gluon jets are modified differently

Use quark jets and gluon jets as independent probes



- ▶ Classify quark jets and gluon jets in pp and AA
- ▶ Distinguish pp jets from AA jets
 - ▶ identify **all** jet features which encode **all** jet modifications
- ▶ Closely related to quark/gluon discrimination
 - ▶ highlight quark and gluon jet differences






Jet representations



- ▶ Different multivariate techniques suit different jet representations
 - ▶ list of physics-motivated observables
 - ▶ unbiased and raw input
 - ▶ complete basis and expansion
 - ▶ Modern computation power and deep learning tools help benchmark jet feature identification
-
- ▶ Illustrate using supervised learning in classification task
 - ▶ Quark and gluon enriched jet samples are generated from Monte Carlo simulations
 - ▶ e.g. using JEWEL $q + \gamma$ and $g + \gamma$ channels (Zapp et al)
 - ▶ methods applicable to all simulations and experimentally quark/gluon-enriched data

Physics-motivated multivariate analysis

- ▶ Representative variables capturing quark and gluon jet features
- ▶ Exploiting observable correlations in high-dimensional space
- ▶ jet mass and radial moments $\sum_i p_T^i \Delta\theta_{jet,i}^\kappa / p_T^{\text{jet}}$ with $\kappa = 0.5, 1$
- ▶ $p_T^D = \sqrt{\sum_i p_T^i{}^2} / p_T^{\text{jet}}$ and pixel multiplicity



Jet image








Image recognition using convolutional neural network






- ▶ A convolutional neural network is trained on GPU using quark and gluon jet images
- ▶ New jet features are learned with significantly improved tagging performance
- ▶ In this work we use grey scale jet images encoding jet energy distribution

Schwartz et al, Deep learning in color, JHEP01(2017)110

Telescoping Deconstruction: a complete subjet fragmentation basis





- ▶ A fixed-order N subjet expansion with subjet kinematics
 - ▶ identify dominant energy flow directions using N soft recoil-free axes
 - ▶ reconstruct subjets around the axes with multiple subjet radii R
 - ▶ TD observables represent *subjet topology* and *subjet substructure*
- ▶ Closely related to perturbative expansion and parton shower picture



Chien et al, arXiv:1711.11041

Telescoping subjet topology

- Enhancement of soft, wide angle radiation





 z  θ 

Telescoping subjet substructure

- Reveal subjet flavor dependence in first splitting $q \rightarrow qg$ and $g \rightarrow gg$ using $m^{\text{sub}}/p_T^{\text{sub}}$



Hard Subjet

Soft Subjet





Collinear-drop: probing soft-dropped radiation

- ▶ Variation of m^2 between ungroomed and groomed jets: $\delta m = \sqrt{m_{\text{ungroomed}}^2 - m_{\text{groomed}}^2}$
- ▶ Quark/gluon jet difference disappearing in AA collisions





Quark gluon jet classification



- ▶ ROC curves: the lower the curve, the better the performance
- ▶ Performance drops in heavy ion collisions
- ▶ Information contained in subleading subjets is washed out in JEWEL

pp and AA jet classification



- ▶ Gluon jets are modified more than quark jets
- ▶ Pixel multiplicity is the dominant feature distinguishing pp and AA jets in JEWEL

Conclusion

- ▶ Quark and gluon jet classification provides a new method of studying jet modification
- ▶ Modifications of collective jet substructure observables provide qualitatively new insights
- ▶ Machine-learning techniques are powerful tools in jet modification studies
- ▶ Quark/gluon classification performance drops in JEWEL-simulated AA collisions
- ▶ Telescoping deconstruction provides a complete and systematic jet substructure framework