


Measurements of strange and non-strange charm production in PbPb collisions at 5.02 TeV with the CMS detector

Cheng-Chieh Peng

Purdue University



for the CMS Collaboration

Hard Probes 2018



Motivation


- Heavy quarks produced early , experience the full evolution of the medium
- $D^0 R_{AA}$: nuclear modification factor
 - Flavor dependent energy loss
 - Dead cone effect (Phys. Lett. B **519** (2001) 199)
- $D^0 v_n$ harmonics in PbPb :
 - At low p_T , the degree of medium thermalization
 - At high p_T , the path length dependence of energy loss
- D^0 elliptic flow v_2 in high multiplicity pPb
 - Evidence of QGP in small system?
 - Heavy flavor hydrodynamic flow?



D^0 Meson Reconstruction & Selection

- $D^0 \rightarrow K^- \pi^+$, BR = 3.89%, $c\tau \simeq 120 \mu\text{m}$ Prompt:

- D^0 candidates :
 - pairing two charged tracks
 - kinematic fitter





- D^0 candidates selection (TMVA Rectangular Cuts)
 - Pointing angle $\alpha < 0.12$
 - 3D decay length significance
 - D^0 candidate vertex probability
 - Distance of Closet Approach (DCA) $< 0.008 \text{ cm}$

D⁰ Signal Extraction by Invariant Mass Fit

D⁰ invariant mass distributions are fitted by

- Double Gaussian (Signal)
- 3rd order polynomial (Combinatorial)
- Single Gaussian (K- π swapped. No PID. Candidates with wrong mass assignment on tracks)




PLB 782,474(2018)




Extract Prompt Fraction from Data

- D^0 in data is a mixture of prompt and non-prompt D^0
- Fit DCA of data with prompt and non-prompt D^0 DCA MC templates

PLB 782,474(2018)




$D^0 R_{AA}$ and Comparison with Model Calculations



- Charm quarks lose a significant fraction of energy in the QGP medium
- R_{AA} minimal near $p_T \sim 10$ GeV/c and then increases
- At high p_T , both pQCD and AdS/CFT predictions reasonably agree with R_{AA} results
- At low p_T , PHSD with shadowing describes data better

PLB **782**,474(2018)

R_{AA} Zoo Plot and Comparison



- At low p_T , a hint of smaller suppression of D^0 and non-prompt J/ψ than charged particles
- At high p_T , the D^0 R_{AA} is similar to charged particles R_{AA}
- The non-prompt J/ψ appear to be less suppressed than the D^0 for p_T smaller than ~ 15 GeV

JHEP **04**(2017)39

PRL **119**,152301(2017)

PLB **782**,474(2018)


EPJC **78** (2018) 509

$D^0 \nu_n$ in PbPb collisions at 5.02 TeV

- ν_n obtained by scalar product method
(Luzum, Ollitrault PRC **87** (2013), 044907)
- Simultaneous fit on mass distribution and ν_n vs. mass


$$\nu_n^{S+B}(m) = \alpha(m) \nu_n^S(m) + [1 - \alpha(m)] \nu_n^B(m)$$

$$\alpha(m) = \frac{\text{Sig}(m) + \text{Swap}(m)}{\text{Sig}(m) + \text{Swap}(m) + \text{Bkgd}(m)}$$



PRL **120**,202301(2018)


Prompt D^0 v_2 Result



- Positive prompt D^0 v_2 is observed:
 - Low p_T : charm quark collective motion
 - High p_T : path length dependence of energy loss
- Similar p_T dependence to charged particle
- At centrality 10-30% and 30-50% , the $v_2(D^0) < v_2(\text{charged particle})$
 - Mass ordering or other effect?

PRL **120**,202301(2018)


Prompt D^0 v_3 Result



- Low p_T : $v_3(\text{prompt } D^0) > 0$; Hight p_T : $v_3(\text{prompt } D^0) \approx 0$
- Similar p_T dependence to charged particle

PRL **120**,202301(2018)

Prompt D^0 v_3 Result



- Low p_T : $v_3(\text{prompt } D^0) > 0$; Hight p_T : $v_3(\text{prompt } D^0) \approx 0$
- Similar p_T dependence to charged particle
- Little centrality dependence
 - Indicate a constant initial geometry
- v_2 and v_3 results provide constrain on models

PRL **120**,202301(2018)

$D^0 \nu_2$ in pPb Collisions at 8.16 TeV


- Two-particle correlation method

to extract ν_2

- Correlate D^0 and charged hadrons ($|\Delta\eta|_{\text{gap}} = 1$)
- Perform Fourier fits the two particle correlation


$$- \nu_2^{D^0}(p_T) = \frac{V_{2\Delta}(p_T^{D^0}, p_T^{\text{assoc}})}{\sqrt{V_{2\Delta}(p_T^{\text{assoc}}, p_T^{\text{assoc}})}}$$

- $D^0 \nu_2^{\text{sub}}$, to reduce the non-flow contributions
 - subtracting the $V_{2\Delta}$ in low multiplicity ($N_{\text{trk}} < 35$)
- Simultaneous fit on mass distribution and ν_2 vs. mass




PRL **121**,082301(2018)

D^0 Meson and Light Hadrons v_2 vs p_T




PRL **121**,082301(2018)




- Significant D^0 v_2 have been observed in high multiplicity pPb

- $v_2^{D^0} < v_2^{\text{light hadrons}}$

D^0 Meson v_2 vs p_T and PbPb Collisions




PRL **121**,082301(2018)




- $D^0 v_2^{pPb} < v_2^{PbPb}$ for a given p_T
- Similar mass ordering for pPb and PbPb

D^0 NCQ Scaling v_2 in pPb and PbPb



PRL 121,082301(2018)




- Number of constituent quarks (NCQ) scaling is motivated by quark coalescence model
- In pPb, D^0 v_2/n_q is smaller than strange hadrons for $KE_T/n_q < 2$
- In PbPb, D^0 v_2/n_q follow the same trend as other particle species

Summary


- $D^0 R_{AA}$ at 5.02 TeV PbPb**

- Strong suppression of $D^0 R_{AA}$
- $R_{AA}(D^0) \sim R_{AA}(h^\pm)$ at high p_T
- $R_{AA}(D^0) > R_{AA}(h^\pm)$ at low p_T



- $D^0 v_2$ at 8.16 TeV pPb**

- Significant v_2 in high multiplicity events
- $v_2(D^0) < v_2$ (strange hadrons)



Back Up




The CMS Trigger and Data Sets

Data sets

- LHC Run II 2015 pp and PbPb at $\sqrt{s_{NN}} = 5.02$ TeV and 2016 pPb data at $\sqrt{s_{NN}} = 8.16$ TeV
- Minimum bias sample for $p_T < 20$ GeV/c and triggered samples for $p_T > 20$ GeV/c
- Dedicated HLT D meson filters to enhance the statistics of very high p_T D mesons
- High multiplicity trigger to select high multiplicity pPb events comparable to peripheral PbPb


Triggering system

Hardware Level 1 Jet Trigger Selections




Level 1 (L1) jet algorithm
with online background
subtraction

Track Selections in Software Triggers




Track seed p_T cut applied:
 $p_T > 2$ GeV/c for pp/pPb
 $p_T > 8$ GeV/c for PbPb

D^0 Selections



D^0 online reconstruction
Loose selections based
on D^0 vertex displacement

Scalar Product Method



$$Q_n = \sum_j w_j e^{i n \phi_j}$$

Sum over tracks (tracker),
or towers (HF)
 w_j : tower E_T for HF, track
 p_T for tracker

$$v_n \{ \text{SP} \} = \frac{\langle Q_n \cdot Q_{nA}^* \rangle}{\sqrt{\frac{\langle Q_{nA} \cdot Q_{nB}^* \rangle \langle Q_{nA} \cdot Q_{nC}^* \rangle}{\langle Q_{nB} \cdot Q_{nC}^* \rangle}}}$$

Scaling factor from 3 sub events

- Large η gap applied ($|\Delta\eta| > 3.0$)
- $v_n \{ \text{SP} \}$, non-ambiguous measure of $\sqrt{\langle v_n^2 \rangle}$

Luzum, Ollitrault PRC **87** (2013), 044907

D^0 v_2 in pPb Collisions at 8.16 TeV

- Fourier series describing the azimuthal anisotropy of particle spectrum


$$\frac{dN}{d\phi} \propto 1 + \sum 2v_n(p_T, \eta) \cos[n(\phi - \psi_n)]$$

- Two-particle correlation method to extract v_2**
 - Correlate D^0 and charged hadrons ($\Delta\eta$ gap = 1)
 - Perform Fourier fits the two particle correlation

distribution for D^0 to extract $V_{2\Delta}(p_T^{D^0}, p_T^{assoc})$

- D^0 $v_2(p_T)$ can be obtain by :

$$v_2^{D^0}(p_T) = \frac{V_{2\Delta}(p_T^{D^0}, p_T^{assoc})}{\sqrt{V_{2\Delta}(p_T^{assoc}, p_T^{assoc})}}$$



CMS HIN-17-003

Phys. Rev. Lett **121**,082301(2018)