

# SUMMARY OF CMS RESULTS IN HEAVY ION COLLISIONS

Javier Martin Blanco

Laboratoire Leprince-Ringuet, École Polytechnique, Palaiseau



RENCONTRES QGP FRANCE 2017

October 9<sup>th</sup>



# OUTLINE

## 3 collision systems at LHC: pp, pPb and PbPb



### Initial stage

- ✦ Constrain quark and gluon (nuclear) PDFs  
Scanning  $(x, Q^2)$  phase space
  - 📍 W and Z bosons
  - 📍 Dijets, b and c -jets
  - 📍 Top quarks
  - 📍 Hidden and open heavy flavor
- ✦ Initial state geometry and fluctuations
  - 📍 Correlation of Fourier harmonics ( $v_2$  and  $v_3$ )

### Final stage

- ✦ Study QGP properties  
Debye screening, energy loss...
  - 📍 Jet quenching
  - 📍 Nuclear modification factors
- Transport coefficients
  - 📍 Fourier harmonics and their correlation ( $v_2$  and  $v_4$ )


# W AND Z BOSONS IN pPb

Alice Florent thesis

W and Z sensitive to quark PDF


$W \rightarrow \ell \nu$

$Z \rightarrow \ell \ell$



PLB 750 (2015) 565-586

- ◆ Different modification of u and d PDFs ?



PLB 759 (2016) 36-57

- ◆ Nuclear effects modify asymmetrically the rapidity distribution


**New measurement of W production in pPb at 8 TeV ongoing  
Stay tuned!**

# DIJET PSEUDORAPIDITY IN pp and pPb

Provide **important constraints to nPDFs** in a wide range of  $x$  and  $Q^2$  ( $\eta_{\text{dijet}}$  and  $p_{\text{T}}^{\text{avg}}$ )

$$\eta_{\text{dijet}} = (\eta_1 + \eta_2) / 2 \sim \log(x_1/x_2)$$

$$p_{\text{T}}^{\text{avg}} = (p_{\text{T},1} + p_{\text{T},2}) / 2$$



**Significant modification of pPb pseudorapidity distribution wrt pp**



# c & b -JETS IN pPb

Important role of  
Matt Nguyen

HF jets **sensitive to gluon PDF**

**c-jets**

**b-jets**



- ♦ First measurement of c-jets in pp and pPb

**Compatible spectra in pp and pPb**




# TOP QUARK CROSS SECTION IN pPb


Gluon PDF poorly constrained in the high  $x$  and  $Q^2$  region

Measurement of **top cross section in pPb** allows to scan this region

## Decay topology




- ♦ 4 jets + 1 isolated lepton
- ♦ 2 b-jet requirement reduce background



**First observation of top quarks in nuclear collisions**

# TOP QUARK CROSS SECTION IN pPb




**Measured cross section  $\sigma_{\text{tt}} = 45 \pm 8$  nb**

**Consistent with theoretical calculations and pp scaled reference**


# PROMPT J/ψ IN pPb and PbPb

Andre's talk: Tuesday 10am



**pPb**: small modification

**Constraint to nPDFs**



**PbPb**: consistent with previous measurement



**Low p<sub>T</sub>: regeneration?**

**High p<sub>T</sub>: energy loss?**



# PROMPT $\psi(2S)$ vs $J/\psi$ IN pPb and PbPb

Andre's talk: Tuesday 10am



Contact: Javier

$\psi(2S)$  **more suppressed** than  $J/\psi$  both in pPb and PbPb



**pPb**: importance of final state effects for excited states

**PbPb**: medium effects stronger for excited states

**Theoretical challenge, especially in pPb**

# $\Upsilon(1S, 2S, 3S)$ IN PbPb

Measurements of  $\Upsilon$  family provide insight to **thermal properties of the medium**




- ✦ Excited states more suppressed than ground state
- ✦ Increasing suppression with centrality
- ✦ Hydrodynamical model consistent with measurements: Initial medium temperature raises from  $T \sim 550 \text{ MeV}$  @ 2.76 TeV to  $T \sim 630 \text{ MeV}$  @ 5.02 TeV


**Nicolas Filipovic**  
thesis @2.76TeV

# D AND B MESONS IN PbPb

Measurements of open heavy flavor allow to study **properties of in-medium energy loss**



arXiv: 1708.04962




arXiv: 1705.04727

Similar suppression of D and B mesons

Theoretical models differ on modelling of the medium, energy loss sources and shadowing

- ✦ D meson suppression at high  $p_T$  is qualitatively reproduced but not at low  $p_T$
- ✦ More precise measurements needed for B mesons


# D MESON AND NONPROMPT J/ψ IN PbPb



- ✦ Light quarks and charm flow with the medium
- ✦ If  $v_2(B) > 0$  b quarks also flow
- ✦ Large uncertainties do not allow to draw conclusions on flavor dependence

# FLAVOR DEPENDENCE OF $E_{\text{loss}}$

PbPb 368  $\mu\text{b}^{-1}$ , pp 28.0  $\text{pb}^{-1}$  (5.02 TeV) **Andre's talk: Tuesday 10am**




- ✦ High  $p_T$ : similar suppression of  $D^0$ , light hadrons and nonprompt  $J/\psi$ : **universal behaviour of  $E_{\text{loss}}$  at high  $p_T$ ?**
- ✦ Low  $p_T$ : hints of  $R_{AA}(B \rightarrow J/\psi) > R_{AA}(D^0) \sim R_{AA}(\text{light hadrons})$ : **smaller  $E_{\text{loss}}$  of b quarks at low  $p_T$ ?**

**Constrains on flavor dependence of  $E_{\text{loss}}$**





# HF DIJETS: MOMENTUM IMBALANCE

Momentum imbalance is ascribed to jet quenching



$$\langle x_J \rangle = \frac{p_{T,2}}{p_{T,1}}$$




CMS-PAS-HIN-16-005

Stas Lisniak thesis

- ♦ b dijet removes ambiguity regarding production mechanism
- ♦ Consistent modification in inclusive and b dijets


**Constrains on mass and flavor dependence of energy loss**

# BOSON-JET MOMENTUM RATIO




$\gamma$  and  $Z$  are not affected by the medium

$\gamma + \text{jet}$




$Z + \text{jet}$



- ✦ Measures energy loss of jets wrt EW probe
- ✦ Average value of transverse momentum ratio smaller in PbPb than pp

**Constrains to quenching with well-defined parton flavor and kinematics**

# COLLECTIVITY IN SMALL SYSTEMS



**Similar ridge effect observed in all hadronic systems at LHC**

Sensitive to:

- ✦ Initial state geometry and fluctuations
- ✦ Transport coefficients of the medium


**What is the origin of the ridge? Common paradigm in all systems?**



# COLLECTIVITY IN SMALL SYSTEMS

$v_2$ ,  $v_3$  and  $v_4$  harmonics studied with multi-particle correlations

CMS-PAS-HIN-16-022





- ♦ Similar pattern of  $v_n$  observed for all systems
- ♦ Very small energy dependence in pPb collisions

**Evidence of collectivity in all hadronic systems at the LHC**

# COLLECTIVITY IN SMALL SYSTEMS

Correlation between harmonics studied with normalised symmetric cumulants (SC)



- ♦ Similar in pPb and PbPb at high multiplicity

→ **Points to the same nature of initial state fluctuations**

- ♦ The same in high multiplicity pp ?

- ♦ Ordering observed:  $pp > pPb > PbPb$

→ **May point to different medium transport coefficients**



# SUMMARY

- Many **new CMS measurements** to:
  - ✦ Constrain PDFs and understand shadowing effects
  - ✦ Understand initial state geometry/fluctuations in all hadronic systems
  - ✦ Better understand QGP properties and interaction of partons with the medium

We still have to make full use of all the data collected so stay tuned !

**THANK YOU FOR YOUR ATTENTION**

**BACKUP**

