



Measurements of charmonia production in b-hadron inclusive decays in LHCb

A. Usachov

Mini-workshop on charmonium production at LHCb
June 16, 2017

Introduction

- Test of **NRQCD factorization** framework
 - Universality assumption: long-distance hadronization of $c\bar{c}$ pair to charmonium is independent from the way $c\bar{c}$ pair was produced
 - => **long-distance matrix elements (LDME) are the same for prompt and b-decays production**
 - Spin symmetry:
 - => linked LDMEs parameters for J/ψ and $\eta_c(1S)$
 - => linked LDMEs parameters for $\chi_{c0,1,2}$ and h_c
- Powerful tests of FONLL using $b \rightarrow J/\psi X$
- Nice to measure production of other states
- Mass and natural width of $\eta_c(1S)$

The most precise measurements of mass and natural width of $\eta_c(1S)$ were performed using η_c produced in radiative decays (PRL 108, 222002)

 - > *significant asymmetry in the lineshapes was observed [PRD 73, 054005]*
 - > *the most precise BES results shifted world average of $\Gamma(\eta_c(1S))$ by $> 2\sigma$*
- Properties of $\eta_c(2S)$ are not well studied

Decays of charmonia

	$\mu\mu$	$J/\psi\gamma$ $J/\psi\pi^+\pi^-$	$p\bar{p}$	$\phi\phi$	$p\bar{p}\pi^+\pi^-$	$\phi f_0(980)$	$\phi f_2(1545)$	baryons
$\eta_c(1S)$	forb.	-	0.15%	0.18%	0.5%			~0.1%
J/ψ	6%	-	0.2%	forb.	0.6%	0.03%	~0.1%	~0.1%
χ_{c0}	forb.	1.3%	0.02%	0.08%				~0.04%
χ_{c1}	forb.	34%	0.01%	0.04%	0.05%			~0.01%
h_c	forb.		<0.015%	forb.	?			?
χ_{c2}	forb.	19%	0.1%	0.01%	0.1%			~0.01%
$\eta_c(2S)$	forb.		<0.1%	?	?			?
$\psi(2S)$	0.8%		0.03%	forb.	0.06%			~0.02%


- The most precise studies were performed for J/ψ (and $\psi(2S)$) using clean $\mu\mu$ channel
- χ_{c1} and χ_{c2} are accessed using $J/\psi\gamma$ channel
- Use charmonia decays to hadrons to measure production of other charmonium states ($\eta_c(1S)$, $\eta_c(2S)$, $h_c?$, χ_{c0})
 - *LHCb is well suited to measure charged hadron final states*

$\Lambda\Lambda$
 $\Xi\Xi$
 $\Sigma\Sigma$

J/ψ production in inclusive b-decays



- Simultaneous **invariant mass** and **pseudo-proper decay time t_z** fit to separate prompt J/ψ and J/ψ from b-decays

$$t_z = \frac{(z_{SV} - z_{PV}) \times M_{J/\psi}}{p_z}$$



Results:

- Differential cross-section measurement described by FONLL prediction




$\psi(2S)$ production in inclusive b-decays

- Simultaneous **invariant mass** and **pseudo-proper decay time** fit to separate prompt $\psi(2S)$ and $\psi(2S)$ from b-decays


$$\frac{\mathcal{B}(b \rightarrow \psi(2S)X)}{\mathcal{B}(b \rightarrow J/\psi X)} = 0.235 \pm 0.005 \text{ (stat)} \pm 0.015 \text{ (syst)},$$

$$\mathcal{B}(b \rightarrow \psi(2S)X) = (2.73 \pm 0.06 \text{ (stat)} \pm 0.16 \text{ (syst)} \pm 0.24 \text{ (BF)}) \times 10^{-3}$$




J/ψ production in inclusive b-decays

Measurements were performed for $\sqrt{s} = 2.76, 7, 8, 13 \text{ TeV}$



Ratio of 13 TeV/ 8 TeV production:

- Powerful test of FONLL: experimental and theoretical uncertainties partially cancel



$\eta_c(1S)$ production in inclusive b-decays using $p\bar{p}$ at $\sqrt{s} = 7,8 TeV$

- Hadronic charmonia decays give access to non 1^{--} charmonia

Analysis:

- Normalization channel: $J/\psi \rightarrow p\bar{p}$
- Prompt and b-decays charmonia separated by t_z cut




Results:

- First measurement of η_c production in inclusive b-decays:



$$\mathcal{B}(b \rightarrow \eta_c(1S)X) / \mathcal{B}(b \rightarrow J/\psi X) = 0.421 \pm 0.055 \pm 0.025 \pm 0.045_{\mathcal{B}}$$

$$\mathcal{B}(b \rightarrow \eta_c(1S)X) = (4.88 \pm 0.64 \pm 0.29 \pm 0.67_{\mathcal{B}}) \times 10^{-3}$$

- Differential production cross-section:




$\eta_c(1S)$ production in inclusive b-decays using $p\bar{p}$ at $\sqrt{s} = 13 \text{ TeV}$ (internal results)



Strategy:

1. Use "b-decays" sample to determine masses and resolution parameter
2. Perform invariant mass fit in bins of t_z
3. t_z fit to separate prompt from b-decays
4. Measure differential production in PT bins


Simultaneous (J/ψ and η_c) χ^2 integral t_z -fit:




Agreement between Run I and Run II measurements

χ_c and $\eta_c(2S)$ production in inclusive b-decays using $\phi\phi$ at $\sqrt{s} = 7,8 \text{ TeV}$

- Normalization channel: $\eta_c(1S) \rightarrow \phi\phi$
- Prompt charmonia removed by flight distance cut
- 2D fit of $M(K^+K^-_1) \times M(K^+K^-_2)$ in bins of $M(KKKK)$ to remove ϕKK and $KKKK$ backgrounds




- Invariant mass spectrum of true $\phi\phi$ combinations:



χ_c and $\eta_c(2S)$ production in inclusive b-decays using $\phi\phi$ at $\sqrt{s} = 7,8 \text{ TeV}$


- First measurement of χ_{c0} production in inclusive b-decays
- The most precise measurements of $BR(b \rightarrow \chi_{c1} X)$ and $BR(b \rightarrow \chi_{c2} X)$
- $BR(b \rightarrow \chi_{c1} X)$ and $BR(b \rightarrow \chi_{c2} X)$ are in agreement with measurements at B-factories



- First measurement of $\eta_c(2S)$ production in inclusive b-decays; first evidence of $\eta_c(2S) \rightarrow \phi\phi$

$$\frac{BR(b \rightarrow \eta_c(2S)X) BR(\eta_c(2S) \rightarrow \phi\phi)}{BR(b \rightarrow \eta_c(1S)X) BR(\eta_c(1S) \rightarrow \phi\phi)} = 0.040 \pm 0.011 \pm 0.004 \quad (3.7\sigma \text{ significance})$$

$\eta_c(2S)$ production as a function of assumed $\Gamma[\eta_c(2S)]$



Search for $X(3872), X(3915)$ and $\chi_{c2}(2P)$ in inclusive b-decays using $\phi\phi$ at $\sqrt{s} = 7,8 \text{ TeV}$

- Bayesian upper limits on the $X(3872), X(3915)$ and $\chi_{c2}(2P)$ production rates:

$$\frac{\mathcal{B}(b \rightarrow X(3872)X) \times \mathcal{B}(X(3872) \rightarrow \phi\phi)}{\mathcal{B}(b \rightarrow \chi_{c1}X) \times \mathcal{B}(\chi_{c1} \rightarrow \phi\phi)} < 0.39(0.34), \quad @ 90(95)\% \text{ CL}$$

$$\frac{\mathcal{B}(b \rightarrow \chi_{c0}(2P)X) \times \mathcal{B}(\chi_{c0}(2P) \rightarrow \phi\phi)}{\mathcal{B}(b \rightarrow \chi_{c0}X) \times \mathcal{B}(\chi_{c0} \rightarrow \phi\phi)} < 0.14(0.12),$$


$$\frac{\mathcal{B}(b \rightarrow \chi_{c2}(2P)X) \times \mathcal{B}(\chi_{c2}(2P) \rightarrow \phi\phi)}{\mathcal{B}(b \rightarrow \chi_{c2}X) \times \mathcal{B}(\chi_{c2} \rightarrow \phi\phi)} < 0.20(0.16)$$

$$\mathcal{B}(b \rightarrow X(3872)X) \times \mathcal{B}(X(3872) \rightarrow \phi\phi) < 4.5(3.9) \times 10^{-7}, \quad @ 90(95)\% \text{ CL}$$

$$\mathcal{B}(b \rightarrow \chi_{c0}(2P)X) \times \mathcal{B}(\chi_{c0}(2P) \rightarrow \phi\phi) < 3.1(2.7) \times 10^{-7},$$

$$\mathcal{B}(b \rightarrow \chi_{c2}(2P)X) \times \mathcal{B}(\chi_{c2}(2P) \rightarrow \phi\phi) < 2.8(2.3) \times 10^{-7}.$$

χ_c and $\eta_c(2S)$ production in inclusive b-decays using $\phi\phi$ at $\sqrt{s} = 7,8 \text{ TeV}$.
 Normalized differential production





Exponential slopes of normalized differential production cross-section are extracted:

	$\eta_c(1S)$	χ_{c0}	χ_{c1}	χ_{c2}
$\sqrt{s} = 7 \text{ TeV}$	0.41 ± 0.02	0.32 ± 0.04	0.31 ± 0.06	0.30 ± 0.05
$\sqrt{s} = 8 \text{ TeV}$	0.39 ± 0.02	0.37 ± 0.04	0.41 ± 0.06	0.33 ± 0.04

Mass and natural width of $\eta_c(1S)$

LHCb measurements:

- $b \rightarrow (\eta_c(1S) \rightarrow p\bar{p})X$
- $b \rightarrow (\eta_c(1S) \rightarrow \phi\phi)X$
- **Another determination from exclusive $B^+ \rightarrow (\eta_c(1S) \rightarrow p\bar{p})K^+ \rightarrow$**



Precision of η_c mass is competitive to PDG

Status of charmonia production measurements

	Prompt hadroproduction	$BR(B^0 B^\pm b - \text{baryons} \rightarrow (c\bar{c})X)$	$BR(B^0 B^\pm \rightarrow (c\bar{c})X)$ (B-factories)
$\eta_c(1S)$	LHCb - $p\bar{p}$	$(4.88 \pm 0.96) \times 10^{-3}$ LHCb - $p\bar{p}$	-
J/ψ	LHCb, ATLAS, CMS - $\mu\mu$	$(1.16 \pm 0.10) \times 10^{-3}$ LEP - ll	$(1.094 \pm 0.032) \times 10^{-2}$ direct: $(7.8 \pm 0.4) \times 10^{-3}$ BABAR, CLEO - ll
χ_{c0}	-	$(3.02 \pm 0.47 \pm 0.23 \pm 0.94B) \times 10^{-3}$ LHCb - $\phi\phi$	-
χ_{c1}	ATLAS, LHCb, CMS - $J/\psi \gamma$	$(1.4 \pm 0.4) \times 10^{-2}$ LEP - $J/\psi \gamma$	$(3.86 \pm 0.27) \times 10^{-3}$ direct: $(3.24 \pm 0.25) \times 10^{-3}$ BABAR, Belle, CLEO - $J/\psi \gamma$
		$(2.76 \pm 0.59 \pm 0.23 \pm 0.89B) \times 10^{-3}$ LHCb - $\phi\phi$	
h_c	-	-	-
χ_{c2}	ATLAS, LHCb, CMS - $J/\psi \gamma$	$(1.15 \pm 0.20 \pm 0.07 \pm 0.36B) \times 10^{-3}$ LHCb - $\phi\phi$	$(1.4 \pm 0.4) \times 10^{-3}$ direct: $(1.65 \pm 0.31) \times 10^{-3}$ BABAR, Belle - $J/\psi \gamma$
$\eta_c(2S)$	-	LHCb - $\phi\phi$ BR($\eta_c(2S) \rightarrow \phi\phi$) was not measured	-
$\psi(2S)$	LHCb, ATLAS, CMS - $\mu\mu$	$(2.83 \pm 0.29) \times 10^{-3}$ LHCb, CMS - $\mu\mu$	$(3.07 \pm 0.21) \times 10^{-3}$ BABAR, CLEO - ll

Summary

- LHCb measured J/ψ production in inclusive b-decays in complementary PT and rapidity range to ATLAS and CMS
 - Differential cross-section is in agreement with FONLL prediction
 - The ratio $R_{13/8}$ is a powerful test of FONLL
- LHCb is well suited to measure hadronic final states to access non 1^{--} charmonia
 - First measurement of $\text{BR}(b \rightarrow \eta_c X)$ using $\eta_c \rightarrow p\bar{p}$
 - First measurement of $\text{BR}(b \rightarrow \chi_{c0} X)$, $\text{BR}(b \rightarrow \eta_c(2S)X)$ and the most precise $\text{BR}(b \rightarrow \chi_{c1,2} X)$ using decays to $\phi\phi$
 - Mass measurements for $\eta_c(1S)$
- Still no measurements of $\text{BR}(b \rightarrow h_c X)$
- Determination of mass width and resolution for further prompt measurements
- Tempting to simultaneous constraint LDMEs by both prompt and b-decays measurements