

HARD SCATTERING AND ELECTROWEAK CORRECTIONS AT THE LHC

J.H. Kühn

- I. Introduction
- II. Form Factors and Four-Fermion Scattering Jantzen, J.H.K., Penin, Smirnov
- III. Z, W and Photon Production J.H.K., Kulesza, Pozzorini, Schulze
- IV. Top Production J.H.K., Scharf, Uwer
- V. W-Pair Production J.H.K., Metzler, Penin, Uccirati
- IV. Real Radiation Bell, J.H.K., Rittinger
- V. Conclusions




I. Introduction

"Typical" size of electroweak corrections: $\frac{\alpha_{\text{weak}}}{\pi} \approx 10^{-2}$

new aspects at LHC: $\sqrt{s} \approx 1\text{-}2\text{TeV} \gg M_{W,Z}$

strong enhancement of negative corrections



one-loop example: massive U(1)

$$\Rightarrow \text{Born} * \left[1 + \frac{\alpha}{4\pi} \left(-\ln^2 \frac{s}{M^2} + 3 \ln \frac{s}{M^2} - \frac{7}{2} + \frac{\pi^2}{3} \right) \right]$$

$\frac{s}{M^2}$	$-\ln^2 \frac{s}{M^2}$	$+3 \ln \frac{s}{M^2}$	$-\frac{7}{2} + \frac{\pi^2}{3}$	Σ	$* 4 \frac{\alpha_{\text{weak}}}{4\pi}$
$\left(\frac{1000}{80}\right)^2$	-25.52	+15.15	-0.21	-10.6	-13%
$\left(\frac{2000}{80}\right)^2$	-41.44	+19.31	-0.21	-22.3	-27%

(four-fermion cross section \Rightarrow factor 4)

- leading \log^2 multiplied by $(\text{charge})^2 = I(I + 1) = \begin{cases} 3/4 & I = 1/2 \\ 2 & I = 1 \end{cases}$
- important subleading logarithms (NLL+...)
- two-loop terms may be relevant
- interplay between electroweak and QCD corrections
- important differences between fermions and electroweak gauge bosons

II. Form Factors & Four-Fermion Scattering at Two Loop

LL: Fadin et al. (2000)

NLL: J.H.K., Penin, Smirnov (2000)
Large (!) subleading corrections
important angular dependent terms


NNLL: J.H.K., Moch, Penin, Smirnov (2001)
Large (!) NNLL terms,
oscillating signs of LL, NLL, NNLL
⇒ compensations

$N^3LL + N^4LL$ Jantzen, J.H.K., Penin, Smirnov (2003-2005)

Additional complication in SM: massless photon

$$|Q^2| \gg M_{W,Z}^2 \gg m_\gamma^2$$

A) Form Factor and Evolution Equations



Born:

$$\mathcal{F}_{\text{Born}} = \bar{\psi}(p_2) \gamma_\mu \psi(p_1)$$

$$\frac{\partial}{\partial \ln Q^2} \mathcal{F} = \left[\int_{M^2}^{Q^2} \frac{dx}{x} \gamma(\alpha(x)) + \zeta(\alpha(Q^2)) + \xi(\alpha(M^2)) \right] \mathcal{F}$$

Collins, Sen

$$\Rightarrow \mathcal{F} = \mathcal{F}_{\text{Born}} F_0(\alpha(M^2)) \exp \left\{ \int_{M^2}^{Q^2} \frac{dx}{x} \left[\int_{M^2}^x \frac{dx'}{x'} \gamma(\alpha(x')) + \zeta(\alpha(x)) + \xi(\alpha(M^2)) \right] \right\}$$

aim: N^4LL \Rightarrow corresponds to all terms of the form:

$$\alpha^n \left[\begin{array}{ccccc} \mathcal{L}^{2n} & + & \mathcal{L}^{2n-1} & + & \mathcal{L}^{2n-2} & + & \mathcal{L}^{2n-3} & + & \mathcal{L}^{2n-4} \\ \text{LL} & & \text{NLL} & & \text{NNLL} & & \text{N}^3\text{LL} & & \text{N}^4\text{LL} \end{array} \right]$$
$$\mathcal{L} \equiv \ln(Q^2/M^2)$$

NNLL requires running of α (i.e. β_0 and β_1) and:


$$\begin{array}{ll} \zeta(\alpha), \xi(\alpha), F_0(\alpha) & \text{up to } \mathcal{O}(\alpha) \quad (\text{one-loop}) \\ \gamma(\alpha) & \text{up to } \mathcal{O}(\alpha^2) \quad (\text{massless two loop}) \end{array}$$

N^3LL requires two-loop calculation in high-energy limit including linear logarithms (available for non-abelian theory)

N^4LL requires complete two-loop calculation in high-energy limit (available for abelian theory)

B) Two-Loop Results: Massive U(1) Model

$$\mathcal{F}_\alpha(M, Q) = \mathcal{F}_{\text{Born}} \left[1 + \frac{\alpha}{4\pi} f^{(1)} + \left(\frac{\alpha}{4\pi} \right)^2 f^{(2)} + \dots \right]$$




$$f^{(2)} = \frac{1}{2} \mathcal{L}^4 - 3 \mathcal{L}^3 + \left(8 + \frac{2}{3}\pi^2 \right) \mathcal{L}^2 - (9 + 4\pi^2 - 24\zeta_3) \mathcal{L}$$

$$+ \frac{25}{2} + \frac{52}{3}\pi^2 + 80\zeta_3 - \frac{52}{15}\pi^4 - \frac{32}{3}\pi^2 \ln^2 2 + \frac{32}{3} \ln^4 2 + 256 \text{Li}_4 \left(\frac{1}{2} \right)$$

$$\mathcal{L} \equiv \ln(Q^2/M^2)$$

C) Massive SU(2) form factor in 2-loop approximation


2-loop vertex diagrams (massless fermions, massive bosons):



$$f_2 = +\frac{9}{32} \mathcal{L}^4 - \frac{19}{48} \mathcal{L}^3 - \left(-\frac{7}{8} \pi^2 + \frac{463}{48} \right) \mathcal{L}^2 \\ + \left(\frac{39}{2} \frac{\text{Cl}_2\left(\frac{\pi}{3}\right)}{\sqrt{3}} + \frac{45}{4} \frac{\pi}{\sqrt{3}} - \frac{61}{2} \zeta_3 - \frac{11}{24} \pi^2 + 29 \right) \mathcal{L}$$

individual contributions

(N^3LL approximation, $M_{\text{Higgs}} = M$, $n_f = 3$, Feynman-'t Hooft gauge)



4-fermion cross section \Rightarrow factor 4!

D) Four fermion scattering

Evaluation in the high energy limit

define

$$\begin{aligned}\mathcal{A}^\lambda &= \bar{\psi}_2 t^a \gamma_\mu \psi_1 \bar{\psi}_4 t^a \gamma_\mu \psi_3 \\ \mathcal{A}_{LL}^\lambda &= \bar{\psi}_{2L} t^a \gamma_\mu \psi_{1L} \bar{\psi}_{4L} t^a \gamma_\mu \psi_{3L} \\ \mathcal{A}_{LR}^d &= \bar{\psi}_{2L} \gamma_\mu \psi_{1L} \bar{\psi}_{4R} \gamma_\mu \psi_{3R}\end{aligned}$$

define “reduced” amplitude $\tilde{\mathcal{A}}$

$$\mathcal{A} = \frac{ig^2}{s} \mathcal{F}^2 \tilde{\mathcal{A}}$$

evolution equation

$$\frac{\partial}{\partial \ln Q^2} \tilde{\mathcal{A}} = \chi(\alpha(Q^2)) \tilde{\mathcal{A}}$$

$\tilde{\mathcal{A}}$: vector in isospin/chiral basis

χ : matrix

N^3LL requires:

- form factor up to N^3LL
- χ up to two loop, as obtained from hard contribution to single pole part of 4-fermion scattering amplitude

e.g. pure massive $SU(2)$ theory with SSB:

$$\begin{aligned}\sigma^{(2)} = & \left[\frac{9}{2} \mathcal{L}^4 - \frac{449}{6} \mathcal{L}^3 + \left(\frac{4855}{18} + \frac{37}{3} \pi^2 \right) \mathcal{L}^2 \right. \\ & \left. + \left(\frac{34441}{216} - \frac{1247}{18} \pi^2 - 122\zeta(3) + 15\sqrt{3}\pi + 26\sqrt{3}\text{Cl}_2\left(\frac{\pi}{3}\right) \right) \mathcal{L} \right] \sigma_B\end{aligned}$$

for identical isospin in initial and final state

Electroweak theory

- infrared logs must be separated
- NNLL
 - result insensitive to form of gauge-boson mass generation
 - term of order $1 - M_W^2/M_Z^2 = \sin^2 \theta$ included
- N³LL
 - sensitive to details of mass generation, gauge boson mixing
 - Approximation: terms of $\mathcal{O}(\sin^2 \theta)$ neglected

Result for the correction factor

$$R(e^+e^- \rightarrow Q\bar{Q}) = 1 - 1.66 L(s) + 5.60 l(s) - 8.39 a + 1.93 L^2(s) \\ - 11.28 L(s)l(s) + 33.79 l^2(s) - 150.95 l(s)a$$

$$R(e^+e^- \rightarrow q\bar{q}) = 1 - 2.18 L(s) + 20.94 l(s) - 35.07 a + 2.79 L^2(s) \\ - 51.98 L(s)l(s) + 321.34 l^2(s) - 603.43 l(s)a$$

$$R(e^+e^- \rightarrow \mu^+\mu^-) = 1 - 1.39 L(s) + 10.12 l(s) - 21.26 a + 1.42 L^2(s) \\ - 20.33 L(s)l(s) + 112.57 l^2(s) - 260.15 l(s)a$$

with


$$L(s) = \frac{g^2}{16\pi^2} \ln^2 \left(\frac{s}{M^2} \right) = 0.07 \quad (0.11)$$

$$l(s) = \frac{g^2}{16\pi^2} \ln \left(\frac{s}{M^2} \right) = 0.014 \quad (0.017)$$


$$a = \frac{g^2}{16\pi^2} = 0.003$$

for $\sqrt{s} = 1 \text{ TeV}$ (2 TeV)

Separate logarithmic contributions to $R(e^+e^- \rightarrow q\bar{q})$ in % to the Born approximation



one-loop LL ($\ln^2(s/M^2)$), NLL ($\ln^1(s/M^2)$)
and $N^2\text{LL}$ ($\ln^0(s/M^2)$)



two-loop LL ($\ln^4(s/M^2)$), NLL ($\ln^3(s/M^2)$),
NNLL ($\ln^2(s/M^2)$) and $N^3\text{LL}$ ($\ln^1(s/M^2)$)

Large cancellations!

III. Z, W and Photon Production at LHC

J.H.K., Kulesza, Pozzorini, Schulze

Phys. Lett. B609(2005) 277, Nucl. Phys. B727(2005) 368,


JHEP 0603:059,2006,

Phys.Lett. B651(2007),

NPB797 (2008) 27

sizable rate at large p_T (1-2 TeV)

Large electroweak corrections ($\hat{s} \gg M_{W,Z}^2$)



High energy limit

consider $q\bar{q} \rightarrow Zg$

NLL $\hat{=}$ double + single logarithmic terms

$$\begin{aligned} H_1^A &\stackrel{\text{NLL}}{=} - \left[\log^2 \left(\frac{|\hat{s}|}{M_W^2} \right) - 3 \log \left(\frac{|\hat{s}|}{M_W^2} \right) \right] H_0, \\ H_1^N &\stackrel{\text{NLL}}{=} - \left[\log^2 \left(\frac{|\hat{t}|}{M_W^2} \right) + \log^2 \left(\frac{|\hat{u}|}{M_W^2} \right) - \log^2 \left(\frac{|\hat{s}|}{M_W^2} \right) \right] H_0 \end{aligned}$$

- remaining subleading terms $\leq 2.5\%$
- NNLL includes non-enhanced terms (angular dependent)
- compact formulae

size of the correction:

$$\sqrt{\hat{s}} = 200 \text{ GeV} : \quad \frac{\delta\sigma}{\sigma} \leq 0.3\%$$

$$\sqrt{\hat{s}} = 4000 \text{ GeV} : \quad \frac{\delta\sigma}{\sigma} \approx 20 - 30\%$$


Result consistent with general considerations

two-loop (NLL):

$$A^{(2)} = \sum_{\lambda=L,R} \left\{ \frac{1}{2} \left(I_{q_\lambda}^Z C_{q_\lambda}^{\text{ew}} + \frac{c_W}{s_W^3} T_{q_\lambda}^3 \right) \left[I_{q_\lambda}^Z C_{q_\lambda}^{\text{ew}} \left(\textcolor{red}{L}_{\hat{s}}^4 - 6 \textcolor{red}{L}_{\hat{s}}^3 \right) \right. \right. \\ \left. \left. + \frac{c_W}{s_W^3} T_{q_\lambda}^3 \left(\textcolor{red}{L}_{\hat{t}}^4 + \textcolor{black}{L}_{\hat{u}}^4 - \textcolor{red}{L}_{\hat{s}}^4 \right) \right] - \frac{T_{q_\lambda}^3 Y_{q_\lambda}}{8 s_W^4} \left(\textcolor{red}{L}_{\hat{t}}^4 + \textcolor{black}{L}_{\hat{u}}^4 - \textcolor{red}{L}_{\hat{s}}^4 \right) \right. \\ \left. + \frac{1}{6} I_{q_\lambda}^Z \left[I_{q_\lambda}^Z \left(\frac{b_1}{c_W^2} \left(\frac{Y_{q_\lambda}}{2} \right)^2 + \frac{b_2}{s_W^2} C_{q_\lambda} \right) + \frac{c_W}{s_W^3} T_{q_\lambda}^3 b_2 \right] \textcolor{red}{L}_{\hat{s}}^3 \right\}$$

with $\textcolor{red}{L}_{\hat{r}}^n = \log^n \left(\frac{|\hat{r}|}{M_W^2} \right)$, $b_1 = -41/(6 c_W^2)$ and $b_2 = 19/(6 s_W^2)$

Complete one loop calculation NLL approximation at two loops



- one-loop $\sim 30\%$ at $p_T \sim 1 \text{ TeV}$
- two-loop relevant above 1 TeV
- important angular-dependent logarithmic terms
- experiment: p_T up to 2 TeV


Relative **NLO** and **NNLO** corrections w.r.t. the **LO**
and **statistical error** for the unpolarized integrated
cross section for $pp \rightarrow Zj$ at $\sqrt{s} = 14 \text{ TeV}$.

(Similarly, but smaller by a factor 2 for jet+ γ)



W production

additional complications:

- photon radiation as necessary part of virtual corrections (gauge invariance)
- IR singularities must be compensated by real radiation
- $p_T(W) = p_T(\text{jet}) + p_T(\gamma)$



(related results: Dittmaier, Kasprzik, ...)



ratios are less sensitive to QCD corrections

IV. Top and Bottom Production:


J.H.K., Scharf, Uwer

Eur. Phys. J. C45(2006) 139


Eur. Phys. J. C51(2007) 37

arXiv: 0909.0059



$q \bar{q} \rightarrow t \bar{t}$:



no
interference
with




$gg \rightarrow t\bar{t}$:





$\sim \mathcal{O}(\alpha_s)$


$\mathcal{O}(\alpha_s^2 \alpha_{\text{weak}})$ weak corrections ($q \bar{q} \rightarrow t \bar{t}$)



+








+





cuts of second group individually IR-divergent

$\mathcal{O}(\alpha_s^2 \alpha_{\text{weak}})$ weak corrections ($g g \rightarrow t \bar{t}$)





- analytical & numerical results available
 (earlier partial results by Beenakker *et al.*, some disagreements)
 independent evaluation by Bernreuther & Fücker
- $(\text{box contribution})_{\text{up-quark}} = -(\text{box contribution})_{\text{down-quark}}$
 \Rightarrow suppression
- box contribution moderately \hat{s} -dependent
- strong increase with \hat{s}
- sizable M_h -dependence, large effect close to threshold

large corrections for large \sqrt{s}
sizable M_h -dependence



(relative weak corrections [%])

Transverse momentum dependence




V. W-Pair Production ($e^+e^- \rightarrow W^+W^-$; $q\bar{q} \rightarrow W^+W^-$)

J.H.K., Metzler, Penin, Uccirati


NPB 795,277 (2008); + in preparation

similar techniques: evolution equation & separation of QED
new aspects:

- longitudinal vs transverse Ws ($I=1$ vs $I=1/2$; equivalence theorem)




- Yukawa coupling in NLL:




- NNLL: \log^2 , \log , const. required in 1 loop

Logarithmic correction to the differential cross section (relative to the Born) at $\sqrt{s} = 1 \text{ TeV}$

1-loop:





transverse







longitudinal

2-loop:



Invariant Mass Distribution ($\theta_{cut} = 30^\circ$)




(larger effects for W_T ! I=1)

VI. Impact of real radiation


Bell, J.H.K., Ritter

arXiv:1004.4117

soft and/or collinear radiation might (partially) compensate the reduction



only collinear radiation




collinear and soft radiation




Summation over isospin in the final state (Z and W radiation allowed)

$$e^+ e^- \rightarrow q\bar{q} + (V):$$

$Q > 0.71 \sqrt{s}$



$Q > 0.84 \sqrt{s}$



$(\sigma_R + \sigma_V)/\sigma_0$ with $1 \text{ TeV} < \sqrt{s} < 5 \text{ TeV}$. Virtual only (lower solid line), only initial state radiation (black dotted line), initial and final state radiation (green, red and blue line) and real radiation with full angular phase space (upper solid line)

IV. Conclusions

- LHC will explore the TeV-region: $\hat{s}/M_W^2 \gg 1$
- electroweak corrections amount to $\mathcal{O}(10\% - 20\%)$ in the interesting kinematic region
- form factors and four-fermion scattering in two loop
- p_T -distributions of Z, W, γ and their ratios will be strongly affected
- two-loop terms might become relevant
- top-quark distributions at large \hat{s} are strongly modified
- sizable M_h -dependence
- large effects for W pair production
- how well can we separate real radiation?