

Multi-Parton Interaction in jet-gap-jet processes

Izabela Babiarz

in collaboration with R. Staszewski and A. Szczurek

Rzeszów University

September 5, 2017

Contents

Introduction

Particle production in jet events


Multi-parton interaction in jets events

Process dynamics

Conclusions

Introduction



The jet-gap-jet process is an example of the diffractive jet production, in which the pomeron is exchanged between the produced jets.



Jet-gap-jet event can be understood as an process where particle production in the rapidity region between the jets is suppressed.

Measurements of the jet-gap-jet process were done by collaborations: DØ, CDF (discussed in *O. Kepka, C. Marquet, and C. Royon, Phys. Rev. D 83, 034036*) and CMS (*Dijet production with a large rapidity between jets, CMS-PAS-FSQ-12-001*)


MPI- suppressing mechanism



Mechanism of destroying rapidity gap, here this will be Multi-Parton Interactions (MPI)

Particle production in jet events

We selected the following example:




with fixed position of jets and fixed $p_T = 50 \text{ GeV}$

The difference between nondiffractive and jet-gap-jet:

- nondiffractive ($gg \rightarrow gg$), normal flow of colour charges
- jet-gap-jet, no colour flow - color singlet


Hadronisation process was obtained with Pythia 8.



The density of produced particles is reduced by two orders of magnitude, when no colour is transferred between the jets.

Particle production in jet events

We define gap between the jets as follows:




Rapidity gap distributions for the selected kinematical configuration.

$$\bar{\eta} = 0$$

$$\eta_{jet1} = -3$$


$$\eta_{jet2} = +3$$




Multi-parton interaction

The MPIs are modeled in Pythia with the help of minijets calculated in collinear factorization approach with a special treatment at low transverse momenta by multiplying standard cross section by a suppression factor $F_{sup}(p_t)$.


$$F_{sup}(p_t) = \frac{p_t^4}{(p_{t0}^2 + p_t^2)^2} \theta(p_t - p_{t,cut})$$



Multi parton interactions in jet-gap-jet events






- additional particles are produced in the region where gap was expected



- the blue curve contains MPI effects but only for events without additional interactions
- gap greater than 5 - no additional interactions occurred

Rapidity gap survival probability

Kinematics dependence of gap survival probability, defined as a fraction of events in which any additional parton-parton interactions occurred.



$$S_G = \frac{\text{result with MPI, } n_{MPI}=0}{\text{result without MPI's}}$$

Process dynamics and MPI modeling

More realistic situations - integration over phase space (Monte Carlo simulations).

We impose only cuts on transverse momentum $200\text{GeV} > p_T > 40\text{GeV}$.

For illustration process dynamics we take LL BFKL amplitude as discussed in *O. Kepka, C. Marquet, and C. Royon, Phys. Rev. D 83, 034036*

$$A(\Delta\eta, p_T^2) = \frac{16N_C\pi\alpha_s^2}{C_F p_T^2}$$

$$\sum_{p=-\infty}^{\infty} \int \frac{d\gamma}{2i\pi} \frac{[p^2 - (\gamma - 1/2)^2] \exp(\alpha\chi_{\text{eff}}[2p, \gamma, \alpha]\Delta\eta)}{[(\gamma - 1/2)^2 - (p - 1/2)^2][(\gamma - 1/2)^2 - (p + 1/2)^2]}$$

BFKL amplitude for $gg \rightarrow gg$ with color singlet exchange

$$A(\Delta\eta, p_T^2) = \frac{16N_C\pi\alpha_s^2}{C_F p_T^2}$$

$$\sum_{p=-\infty}^{\infty} \int \frac{d\gamma}{2i\pi} \frac{[p^2 - (\gamma - 1/2)^2] \exp(\alpha\chi_{eff}[2p, \gamma, \alpha]\Delta\eta)}{[(\gamma - 1/2)^2 - (p - 1/2)^2][(\gamma - 1/2)^2 - (p + 1/2)^2]}$$

with the normalization: $\frac{d\sigma}{dt} = \frac{|A_{gg \rightarrow gg}|^2}{16\pi}$


$$\chi_{eff} = 2\psi(1) - \psi\left(1 - \gamma + \frac{|p|}{2}\right) - \psi\left(\gamma + \frac{|p|}{2}\right)$$

$\psi(\gamma) = d \log \Gamma(\gamma) / d\gamma$ is derivative of

Gamma function


α_s in leading-logarithmic calculations is taken as a constant value

p is called conformal spin




The gap between jets

We find gaps between jets as follows: $\bar{\eta} = \frac{\eta_{jet1} + \eta_{jet2}}{2}$





$$\Delta\eta_{gap} < |\eta_{jet1} - \eta_{jet2}|$$



η_1, η_2 for particles at edges of rapidity gap





Results



Results



Jet-gap-jet vs two-gluon color singlet exchange



Conclusions

- The role of multi parton interactions in jet-gap-jet processes was discussed.
- Fixed kinematical configurations were done.
- LL BFKL framework was used to describe dynamics of jet-gap-jet process.
- The two-gluon simple approximation was performed for comparison.
- The subprocess amplitudes for the color singlet exchange was implemented in PYTHIA 8.

The MPI effects lead to dependence on kinematical variables of the gap survival factor, in contrast what is usually assumed in the literature.

Thank you for your attention.