

PREDICTING THE BARYON ASYMMETRY

P. HERNÁNDEZ
(IFIC, U. VALENCIA & CSIC)

SM+3 massive neutrinos

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U_{PMNS}(\theta_{12}, \theta_{23}, \theta_{13}, \delta, \dots) \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$


$$\theta_{12} \sim 34^\circ$$

$$\theta_{23} \sim 42^\circ \text{ o } 48^\circ$$


$$\theta_{13} \sim 8.5^\circ$$

$$\delta \sim ?$$

normal hierarchy



inverted hierarchy




$$\updownarrow 7.5 \cdot 10^{-5} \text{eV}^2$$

$$2.5 \cdot 10^{-3} \text{eV}^2$$

Why are neutrinos so much lighter ?

Neutral vs charged hierarchy ?



Why so different mixing ?

CKM

$$V_{CKM} \simeq \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

PMNS

$$|V_{PMNS}| \simeq \begin{pmatrix} \sqrt{\frac{2}{3}} & \sqrt{\frac{1}{3}} & 0 \\ \sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \\ \sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \end{pmatrix}$$

Harrison, Perkins, Scott

A new physics scale ?


Neutrinos are different...they can have majorana masses:

$$-\mathcal{L}_{\text{Majorana}} = \bar{\nu}_L m_\nu \nu_L^c + h.c. \leftrightarrow \bar{L} \tilde{\Phi} \alpha \tilde{\Phi} L^c + h.c.$$

Weinberg

$$[\alpha] = -1$$

$$\alpha = \frac{\lambda}{\Lambda}$$




$$m_\nu = \lambda \frac{v^2}{\Lambda}$$

Scale at which new particles will show up
(mass of the neutrino mass mediators)

Seesaw mechanism:

Minkowski
Gell-Mann, Ramond Slansky
Yanagida, Glashow
Mohapatra, Senjanovic



What originates the neutrino mass ?

Could be $\Lambda \gg v \dots$ the standard lore (theoretical prejudice ?)

$$\left. \begin{array}{l} \Lambda = M_{\text{GUT}} \\ \lambda \sim \mathcal{O}(1) \end{array} \right\} m_\nu \checkmark$$

What originates the neutrino mass ?

Could be $\Lambda \gg v$... the standard lore (theoretical prejudice ?)


$$\left. \begin{array}{l} \Lambda = M_{\text{GUT}} \\ \lambda \sim \mathcal{O}(1) \end{array} \right\} m_\nu \checkmark$$

Hierarchy problem

$$m_H^2 \propto \Lambda^2$$

not natural in the absence of SUSY/other solution to the hierarchy problem

The Standard Model is healthy as far as we can see...




Could be naturally $\Lambda \sim v$?

Yes ! λ in front of neutrino mass operator must be small...

Resolving the neutrino mass operator at tree level

E. Ma

Type I see-saw:
a heavy singlet scalar




$$m_\nu = \frac{\alpha v^2}{\Lambda} \equiv Y_N^T \frac{v^2}{M_N} Y_N$$

Minkowski;
Yanagida; Glashow;
Gell-Mann, Ramond Slansky;
Mohapatra, Senjanovic...

$$\lambda \sim O(Y^2)$$

Type II see-saw:
a heavy triplet scalar




$$m_\nu = \frac{\alpha v^2}{\Lambda} \equiv Y_\Delta \frac{\mu_\Delta}{M_\Delta^2} v^2$$

Konetschny, Kummer;
Cheng, Li;
Lazarides, Shafi, Wetterich ...

$$\lambda \sim O(Y \mu/M_\Delta)$$

Type III see-saw:
a heavy triplet fermion




$$m_\nu = \frac{\alpha v^2}{\Lambda} \equiv Y_\Sigma^T \frac{v^2}{M_\Sigma} Y_\Sigma$$


Foot et al; Ma;
Bajc, Senjanovic...

$$\lambda \sim O(Y^2)$$


$$M_N \sim \text{GUT}$$



$$M_N \sim \nu$$




Where is the new scale ?



“Once you eliminate the impossible, whatever remains, no matter how improbable/unnatural, must be the truth.”

Where is the new scale ?



Generic predictions

- there is **neutrinoless double beta** decay at some level ($\Lambda > 100\text{MeV}$)

model independent contribution from the neutrino mass

Where is the new scale ?




Generic predictions:

- a **matter-antimatter asymmetry** if there is **CP violation** in the lepton sector via **leptogenesis**

model dependent...

Where is the new scale ?




Generic predictions:

- there are other states out there at scale Λ : **new physics beyond neutrino masses**

potential impact in cosmology, EW precision tests, LHC, rare searches, $\beta\beta 0\nu$, ...

model dependent...

Where is the new scale ?




The EW scale is an interesting region: **new physics underlying the matter-antimatter asymmetry could be tested !**


Minimal model of neutrino masses:

Type I seesaw: SM+right-handed neutrinos

$$\mathcal{L}_\nu = -\bar{l}Y\tilde{\Phi}N_R - \frac{1}{2}\bar{N}_R M N_R + h.c.$$



$$n_R \geq 2$$




$$m_\nu = \lambda \frac{v^2}{\Lambda} \equiv Y^T \frac{v^2}{M} Y$$

Minkowski; Yanagida; Glashow; Gell-Mann, Ramond Slansky; Mohapatra, Senjanovic...

Type I seesaw models



$n_R = 3$: 18 free parameters (6 masses+6 angles+6 phases)
out of which we have measured 2 masses and 3 angles...



Type I seesaw models

Phenomenology (beyond neutrino masses) of these models depends on the heavy spectrum and the size of active-heavy mixing:

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U_{ll} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix} + U_{lh} \begin{pmatrix} N_1 \\ N_2 \\ N_3 \end{pmatrix}$$



Strong correlation between active-heavy mixing and neutrino masses

Baryon asymmetry

Sakharov's necessary conditions for baryogenesis

- ✓ Baryon number violation
- ✓ C and CP violation
- ✓ Deviation from thermal equilibrium

It does not seem to work in the SM with massless neutrinos ...


CP violation too small ✗

EW phase transition too weak ✗

Massive neutrinos provide new sources of CP violation and non-equilibrium conditions

Leptogenesis

Models with massive neutrinos generically lead to generation of lepton and therefore baryon asymmetries



Standard leptogenesis in out-of-equilibrium
decay $M_N > 10^7 \text{ GeV}$

Fukuyita, Yanagida....; Davidson, Nardi, Nir rev


Can be extended to lower scales at the expense of a extreme degeneracy of the heavy states: **resonant leptogenesis**

Pilaftsis,...

High-scale leptogenesis


New sources of CP violation and L violation in the neutrino sector can induce CP asymmetries in decays of heavy Majorana ν

Fukuyita, Yanagida



$$\epsilon_1 = \frac{\Gamma(N \rightarrow \Phi l) - \Gamma(N \rightarrow \Phi \bar{l})}{\Gamma(N \rightarrow \Phi l) + \Gamma(N \rightarrow \Phi \bar{l})}$$

High-scale leptogenesis



$$\Gamma_N \leq H(M_N)$$

(decay rate < hubble expansion)

Leptogenesis




Leptogenesis from neutrino oscillations
 $0.1\text{GeV} < M < 100\text{GeV}$

Akhmedov, Rubakov, Smirnov;
Asaka, Shaposhnikov,...

Low-scale Leptogenesis

CP asymmetries arise in production of sterile states oscillations




$$L_\alpha \rightarrow L_\beta \neq \bar{L}_\alpha \rightarrow \bar{L}_\beta$$

$$\sum_{\alpha} \Delta_{CP}^{\alpha} = 0$$

$$Y_B \propto \sum_{\alpha} \Delta_{CP}^{\alpha} \eta_{\alpha}$$

Different flavours different efficiency in transferring it to the baryons

Low-scale leptogenesis



$$\Gamma_s(T_{EW}) \leq H(T_{EW})$$

(scattering rate < hubble expansion)

Testability/predictivity ?

- Y_B cannot be determined from neutrino masses and mixings only
- More information from the heavy sector is needed:

High-scale scenarios: very difficult for $M_N > 10^7$ GeV

Low-scale scenarios: N 's can be produced in the lab
and could be in principle detectable !

Full exploration of the minimal model N=2

Bayesian posterior probabilities (using nested sampling Montecarlo Multinest)

$$\mathcal{L} = - \left(\frac{Y_B(\text{param}) - Y_B^{\text{obs}}}{\sigma_{Y_B}} \right)^2$$

Use **Casas-Ibarra** parametrization: fix light neutrino masses and mixings to the best fit oscillation points (IH/NH) and vary

$$R(\theta + i\gamma); U_{PMNS}(\delta, \phi_1); M_1, M_2$$


Flat priors in:

$$\theta = [0, \pi]; \delta = [0, 2\pi]; \phi_1 = [0, 2\pi]; \gamma = [-9, 9];$$
$$\log_{10} M_1 \text{ and } \log_{10} M_2 / \log_{10}(M_2 - M_1)$$


Full exploration of the minimal model N=2

Inverted neutrino ordering (IH)

PH, Kekic, Lopez-Pavon, Racker, Salvado
arxiv:1606.06719



In the minimal model with just $n_R=2$ neutrinos (IH)



PH, Kekic, Lopez-Pavon, Racker, Salvado
arxiv:1606.06719

Colored regions: posterior probabilities of successful Y_B

Predicting Y_B in the minimal model $n_R=2$?


Assume a point within SHIP reach that gives the right baryon asymmetry

- SHIP measurement could provide (if states not too degenerate)

$$M_1, M_2, |U_{e1}|^2, |U_{\mu1}|^2, |U_{e2}|^2, |U_{\mu2}|^2$$

- Future neutrino oscillations: δ phase in the U_{PMNS}

Predicting Y_B in the minimal model $n_R=2$ (IH)




PH, Kekic, Lopez-Pavon, Racker, Salvado
arxiv:1606.06719

- Grey band: standard light neutrino contribution to $m_{\beta\beta}$ for IH
- Significant interference between light/heavy neutrino contributions to $m_{\beta\beta}$

Predicting Y_B in the minimal model $n_R=2$

Heavy states also contribute to the $\beta\beta\nu$ amplitude...



$$m_{\beta\beta} = \underbrace{\sum_{i=1}^3 [(U_{PMNS})_{ei}]^2 m_i}_{\text{Light states}} + \underbrace{\sum_{i=j}^3 U_{ej}^2 M_j \frac{\mathcal{M}^{0\nu\beta\beta}(M_j)}{\mathcal{M}^{0\nu\beta\beta}(0)}}_{\text{Heavy states}}$$


$$M_j \rightarrow \infty \quad \frac{\mathcal{M}^{0\nu\beta\beta}(M_j)}{\mathcal{M}^{0\nu\beta\beta}(0)} \propto \left(\frac{100 \text{ MeV}}{M_j} \right)^2$$

the heavy contribution is sizeable for M_i of O(GeV)

Blennow, Fernandez-Martinez, Lopez-Pavon, Menendez;
Lopez-Pavon, Pascoli, Wong; Lopez-Pavon, Molinaro, Petcov

The non standard contributions bring essential information of some CP phases and other unknown parameters


Predicting Y_B in the minimal seesaw model $M \sim \text{GeV}$



PH, Kekic, Lopez-Pavon, Racker, Salvado
arxiv:1606.06719

The GeV-miracle: the measurement of the mixing to e/μ of the sterile states, neutrinoless double-beta decay and δ in neutrino oscillations have a chance to give a prediction for Y_B

What is this Sample point ?



$$\delta = 234^\circ \quad \alpha = 254^\circ$$

$$M_1 \sim M_2 \sim 0.77 \text{ GeV}, \Delta M/M \sim 10^{-2}$$

How fine-tuned is the range of parameters for successful leptogenesis ?

The very degenerate regions could be understood in terms of an approximate global symmetry $U(1)_L$

Wyler, Wolfenstein; Mohapatra, Valle; Branco, Grimus, Lavoura, Malinsky, Romao; Kersten, Smirnov; Abada et al; Gavela et al....many others

$$L(N_1) = +1, \quad L(N_2) = -1 \quad -\mathcal{L}_\nu \supset \bar{N}_1 M N_2^c + Y \bar{L} \tilde{\Phi} N_1 + h.c.$$

$$\begin{pmatrix} 0 & Y v & 0 \\ Y v & 0 & M_N \\ 0 & M_N & 0 \end{pmatrix}$$

Degenerate heavy neutrinos and massless light neutrinos...

How natural is the range of parameters for successful Leptogenesis ?


The very degenerate regions could be understood in terms of an approximate global symmetry $U(1)_L$

Wyler, Wolfenstein; Mohapatra, Valle; Branco, Grimus, Lavoura, Malinsky, Romao; Kersten, Smirnov; Abada et al; Gavela et al....many others

$$\begin{pmatrix} 0 & Y_1 v & \epsilon Y_2 v \\ Y_1 v & \mu' & M_N \\ \epsilon Y_2 v & M_N & \mu \end{pmatrix}$$


➤ How small must the small entries be ?

How natural is the range of parameters for successful ARS leptogenesis ?




- Blue region prefers a mild hierarchy in all $U(1)_L$ breaking terms
- Red region points requires μ, μ' entries significantly smaller than ϵ

How large can the mixing be (even if less probable) and still have successful baryogenesis ?



Drewes, Garbrecht, Gueter, Klaric
arxiv: 1606.06690




Abada, Arcadi, Domcke, Lucente
arxiv: 1709.00415

The seesaw path to leptonic CP violation:

flavour ratios of heavy lepton mixings strongly correlated with ordering, U_{PMNS} matrix: δ, ϕ_1

In minimal model (N=2):




Caputo, PH, Lopez-Pavon, Salvado arxiv:1611.05000

Superb sensitivity to Majorana CP phases: complementary to oscillations

If SHIP/FCC-ee measures the heavy neutrinos and their mixings to e/μ :

Can we exclude a real U_{PMNS} matrix ie.
discover leptonic CP violation in mixing ?



$$(\delta, \phi_1) \neq (0/\pi, 0/\pi)$$



Leptonic CP violation 5σ CL discovery regions (SHIP)

Normal Hierarchy

Inverted Hierarchy




(no systematic error included)

Caputo, PH, Lopez-Pavon, Salvado arxiv:1611.05000


$R_{\text{CP}=5\sigma}$ CP-fraction =
fraction of the area of the CP rectangle which is colored

Leptonic CP violation 5σ CL discovery regions (SHIP)

Normal Hierarchy



Inverted Hierarchy



(no systematic error included)

Caputo, PH, Lopez-Pavon, Salvado [arxiv:1611.05000](https://arxiv.org/abs/1611.05000)

$R_{CP}=5\sigma$ CP-fraction =
fraction of the area of the CP rectangle which is colored

Conclusions

- Exploring the EW- \rightarrow TeV region for NP related to neutrino masses is very well motivated
- A minimal model of neutrino masses with a new scale near GeV can explain the baryon asymmetry and might do so in a testable way (IH more promising)
- Testability in simplest model will require the contribution of very different type of experiments:

SHIP/FCCee: masses and mixings of the heavy neutrino states

DUNE/HyperK: CP violation ν neutrino oscillations

$\beta\beta 0\nu$: non-standard contributions from heavy sector

- Flavour mixings of the heavy states high sensitivity to CP phases in U_{PMNS} (in particular Majorana phases!)
- Mediators of neutrino mass at the EW provide a new portal to BSM physics

Beyond the minimal model


Many possibilities:

Examples: type I + W' , Z' ,
left-right symmetric models
GUTs, etc

Keung, Senjanovic; Pati, Salam, Mohapatra, Pati; Mohapatra, Senjanovic;
Ferrari et al + many recent refs...

- Generically gauge interactions can enhanced the production in colliders
- But they make leptogenesis more challenging (out-of-equilibrium condition harder to meet)

Interesting possibility:



Model independent approach: EFT

$$\mathcal{L}_{BSM} = \mathcal{L}_{\text{mSS}} + \sum_{d,i} \frac{1}{\Lambda^{d-4}} \mathcal{O}_i^{(d)}$$

The **seesaw portal** to BSM:



d=5

$$\mathcal{O}_W = \sum_{\alpha,\beta} \frac{(\alpha_W)_{\alpha\beta}}{\Lambda} \bar{L}_\alpha \tilde{\Phi} \Phi^\dagger L_\beta^c + h.c.,$$

$$\mathcal{O}_{N\Phi} = \sum_{i,j} \frac{(\alpha_{N\Phi})_{ij}}{\Lambda} \bar{N}_i N_j^c \Phi^\dagger \Phi + h.c.,$$

$$\mathcal{O}_{NB} = \sum_{i \neq j} \frac{(\alpha_{NB})_{ij}}{\Lambda} \bar{N}_i \sigma_{\mu\nu} N_j^c B_{\mu\nu} + h.c.$$



Corrections to $|U_{ei}|^2/|U_{\mu i}|^2$ the minimal model correlated to the lightest neutrino mass



$$m_{1(3)} \sim 0.1 \sqrt{|\Delta m_{\text{sol}}^2|}$$

$\mathcal{O}_{N\Phi}$

could lead to spectacular signals at LHC/colliders of two displaced vertices from higgs decays (production independent of U)





Caputo, PH, LopezPavon, Salvado, arxiv:1704.08721

LHC: 300 fb^{-1} , 13TeV

$$\left| \frac{\alpha_{N\Phi} v}{\sqrt{2}\Lambda} \right| \leq 10^{-3} - 10^{-2} \rightarrow \frac{\alpha_{N\Phi}}{\Lambda} \leq 6 \times (10^{-3} - 10^{-2}) \text{TeV}^{-1}.$$

$\mathcal{O}_{N\Phi}$

could lead to spectacular signals at LHC/colliders of two displaced vertices from higgs decays (production independent of U)



Caputo, PH, LopezPavon, Salvado, arxiv:1704.08721

A large hierarchy between the coefficients would be needed since the Weinberg operator is much more strongly suppressed: technically natural with $U(1)_L$