


Wavelet analysis of the Inner Galaxy








Sam McDermott, Paddy Fox, I.C, Samuel Lee JCAP 1607 (2016), (arXiv:1512.00012)
Bhaskar Balaji, I.C., Sam McDermott, Paddy Fox ,arXiv: 171x.xxxxx



Three elephants in the gamma-ray sky
Ilias Cholis 10/23/2017

Why are we all here?



The galactic center and inner galaxy is a very interesting region. Interesting CR activity. Burst? Point Sources. A possible signal of Dark Matter Annihilation?




Also works from Hooper & Goodenough 2009-10, Hooper Linden, Abazajian & Kaplinghat, Gordon & Macias

What are the explanations?

I will focus on the GCE but similar # of suggestions for the FBs.




DM
YES?



YES?

NO?



Alternative work related to the Galactic Center the GeV excess and its interpretations

Millisecond Pulsars:

Hooper, IC, Linden, Siegal-Gaskins & Slatyer
PRD 2013 (1305.0830), (<10% of total)

Calore, Di Mauro, Donato ApJ 2014
(1406.2706) (<10%)

IC, Hooper, Linden JCAP 2015 (1407.5625)
NOT REALLY ABOVE 5deg

Calore, Di Mauro, Donato, Hessels, Weniger
(1512.06825) MAYBE YES

Brandt, Cocsis ApJ 2015 YES BUT SPECIAL
MSPs


O'Leary, Kistler, Kerr, Dexter 2016
PROBABLY

Sensitivity analyses on point-sources and astrophysics modeling:


Bartels, Krishnamurthi, Weniger PRL 2016

Lee, Lisanti, Safdi, Slatyer, Xue PRL 2016

Huang, Ensslin, Selig JPCS 2016.



As reference we need $1-3 \times 10^3$ MSPs in the inner 2 kpc below threshold



Bursts of Cosmic Rays:

Carlson and Profumo PRD 2014
(PROTONS MAYBE?)

Petrovic, Serpico, Zaharijias JCAP 2014
(ELECTRONS ?)

IC, Evoli, Calore, Linden, Weniger, Hooper
JCAP 2015 (ELECTRONS CAN + FB
CONNECTION?)

Radio Limits:


Bringmann, Vollmann, Weniger PRD 2014
(RELEVANT FOR DM)

IC, Hooper, Linden PRD 2015 (NOT
RELEVANT FOR DM)






Wavelet techniques (TIE BRAKER?):

McDermott, Fox, IC, Lee JCAP 2016

Possible Connection to the Fermi Bubbles



DM35: $2.2 < E_\gamma < 4.9$ GeV, significant wavelet levels




What are wavelets?

Wavelets have been used in image compression (JPEG), de-noising, fast signal identification, even in HEP data

Allow analysis of data in both time/space and frequency space



Different type of structures will have a different power at different levels of the decomposition (e.g. edges and other small scale structures vs larger scale variations).



Wavelets can find these different structures.

What are wavelets?

wavelet coefficients original signal

$$W(a, b) = \frac{1}{\sqrt{a}} \int f(x) \psi^* \left(\frac{x - b}{a} \right) dx$$

scale position mother wavelet
(different choices)

$$\int \psi(x) dx = 0$$
$$\int |\psi(x)|^2 dx = 1$$

$\psi(x) \in \mathbb{L}^2(\mathbb{R})$ and

$$\frac{1}{\sqrt{a}} \psi \left(\frac{x - b}{a} \right) \in \mathbb{L}^2(\mathbb{R})$$

for $a, b \in \mathbb{Z}$


Is it better than the Templates?

Technique was developed in McDermott, Fox, Cholis, Lee JCAP 2016 using simulated data. Requires statistics but has less dependence on fore-ground/background assumptions (in the end it does also have some systematics).


Decompose the sky into 8 scales (smallest 6):




j	w_1	w_2	w_3	w_4	w_5	w_6
θ	$[0.7^\circ, 1.4^\circ]$	$[1.4^\circ, 2.8^\circ]$	$[2.8^\circ, 5.6^\circ]$	$[5.6^\circ, 11.3^\circ]$	$[11.3^\circ, 22.5^\circ]$	$[22.5^\circ, 45^\circ]$

Bubbles, templates




Bubbles, wavelets









We mirror the wavelet coeff. for scales w_4 and above (angles larger than 5 degrees) within a chosen window.




Zoom in to the Inner $60^\circ \times 60^\circ$




GCE defined for $j \geq 3$




wavelet higher scales GCE?




wavelet higher scales GCE?



After iterating in the higher scales





Adding everything back together:





Fermi Bubbles:

we do not directly disentangle FB from GCE.




PRELIMINARY



jets?
cocoon?



Profiles



Same as in Calore et al. & the inner 2 degrees




Isotropic Emission is subtracted in our analysis (flux outside of bubbles <0)

- Flux > 0
- Abs(Flux < 0)

Additional Energies/Profiles


GCE:





Statistical smaller than systematics.

*Systematics come from the **location/size of the mirroring area**, the **wavelet scales we choose to mirror and iterate over**. The **collection ISM** that we average over in the first step when we subtract the galactic diffuse emission.*

Spectra



Fermi Bubbles:




In agreement with Su et al. & Fermi. Coll.


The Wavelet Promise:

Is the power more in small scales (e.g. point sources/filaments) or in large scales (diffuse emission as is ICS)?

Fermi Bubbles:



The Fermi Bubbles have *very little power in small scales* in agreement with the leptonic association (WMAP/Planck Haze)




The GCE seems have also *little power in small scales* apart from region 0. Regions 1 and 2 also PS contaminated. For regions 3 and above (>5 degrees), there is agreement with the diffuse association (CE electron bursts or even DM?)

Conclusions, future directions


- Using the wavelet technique we developed in McDermott et. al JCAP 1607 (2016), (arXiv:1512.00012) **we are analyzing data now** and writing the paper Balaji et al. **171x.xxxxx**.
- We also find the Fermi Bubbles and the Galactic Center Excess and are in agreement with most template results.
- Extract spectra both at different regions and also **at different scales!**
- We can ask questions on the underlying properties of these emissions.
- The GCE **only in the inner 5 degrees** has power in low scales (e.g from point sources and miss-modeling of gas distribution).
- Regarding the interpretations we still have to make a connection with simulations we have from 2016 and run some more...
- The Fermi Bubbles are diffuse above 20 degrees. We do find substructure that may be associated with a cocoon/jet.
- Use this technique to study other regions of the sky.
- A GREAT SET OF TOOLS TO STUDY THE GAMMA-RAY DATA

Additional Slides



Bubble excess vs. latitude, all scales





Statistical Errors ONLY




Diffuse Part of the GCE emission




Bubble morphology without mirroring




North bubble spectrum




GCE regions' flux, 0.68 GeV



GCE regions' flux, 35. GeV



Bubble excess vs. latitude, all scales, 3.3 GeV



Iteration procedure:


$$GCE_0 = \sum_{j=3}^9 w_j(R'_0)$$

$$blob_0 = \sum_{j=3}^9 w_j(R_0 - GCE_0)$$


$$GCE_i = \sum_{j=3}^9 w_j(R_0 - blob_{i-1})$$

$$blob_i = \sum_{j=3}^9 w_j(R_0 - GCE_{i-1})$$


sine wave




Fourier




wavelet
Mex. hat




two sine waves




Fourier




wavelet
Mex. hat



sine waves with transition



Fourier



wavelet
Mex. hat

