



# Searching for Beauty-fully Bound Tetraquarks Using Lattice NRQCD


Ciaran Hughes, Estia Eichten, Christine Davies




*Mesons*



*Baryons*



*pentaquarks - LHCb (2015)*




*tetraquarks*

# Searching for Beauty-fully Bound Tetraquarks Using Lattice NRQCD

Ciaran Hughes, Estia Eichten, Christine Davies


---



$2b2\bar{b}$   
*tetraquarks*

# Searching for Beauty-fully Bound Tetraquarks Using Lattice NRQCD

Ciaran Hughes, Estia Eichten, Christine Davies



*tetraquarks*



**THE FORCE  
IS STRONG  
WITH THIS ONE**


**But is it strong enough  
to confine  $2b2\bar{b}$  into a  
stable state below the  
 $2\eta_b$  threshold.**

# What Does Experiments Have To Say

---



# What Does Experiments Have To Say

---



# What Does Experiments Have To Say





---



# What Does Experiments Have To Say: LHCb

---

# What Does Experiments Have To Say: LHCb




arxiv: 1806.09707



# What Does Experiments Have To Say: LHCb


$$\sigma(pp \rightarrow X) \times \mathcal{B}(X \rightarrow \Upsilon(1S)\mu^+\mu^-) \times \mathcal{B}(\Upsilon(1S) \rightarrow \mu^+\mu^-)$$



☛ Total number of events (signal+bkgrnd) low: Don't see  $2\Upsilon$  signal.

# What Does Experiments Have To Say: LHCb


$$\sigma(pp \rightarrow X) \times \mathcal{B}(X \rightarrow \Upsilon(1S)\mu^+\mu^-) \times \mathcal{B}(\Upsilon(1S) \rightarrow \mu^+\mu^-)$$



- Total number of events (signal+bkgrnd) low: Don't see  $2\Upsilon$  signal.
- CMS have clear signal for  $2\Upsilon$  (arxiv:1610.07095). Why CMS and not LHCb?

# What Does Experiments Have To Say: LHCb


$$\sigma(pp \rightarrow X) \times \mathcal{B}(X \rightarrow \Upsilon(1S)\mu^+\mu^-) \times \mathcal{B}(\Upsilon(1S) \rightarrow \mu^+\mu^-)$$



- ☛ Total number of events (signal+bkgrnd) low: Don't see  $2\Upsilon$  signal.
- ☛ CMS have clear signal for  $2\Upsilon$  (arxiv:1610.07095). Why CMS and not LHCb?
- ☛ CMS / Atlas have much larger luminosity

# What Does Experiments Have To Say: LHCb

$$\sigma(pp \rightarrow X) \times \mathcal{B}(X \rightarrow \Upsilon(1S)\mu^+\mu^-) \times \mathcal{B}(\Upsilon(1S) \rightarrow \mu^+\mu^-)$$



- ☛ Total number of events (signal+bkgrnd) low: Don't see  $2\Upsilon$  signal.
- ☛ CMS have clear signal for  $2\Upsilon$  (arxiv:1610.07095). Why CMS and not LHCb?
- ☛ CMS / Atlas have much larger luminosity
- ☛ CMS / Atlas have larger fiducial volume for lepton pairs

# What Does Experiments Have To Say

*<http://meetings.aps.org/Meeting/APR18/Session/U09.6>*

---

- Note: The results are taken from my thesis work and they are not approved by CMS yet. The analysis is still in review.

# What Does Experiments Have To Say

["http://meetings.aps.org/Meeting/APR18/Session/U09.6"](http://meetings.aps.org/Meeting/APR18/Session/U09.6)



- Note: The results are taken from my thesis work and they are not approved by CMS yet. The analysis is still in review.

4/15/18

Suleyman Durgut

2

## Combined Result



- Do a simultaneous fit to both channels, with fixed signal shapes but floating mass value.
- **Best mass :  $18.4 \pm 0.1$  (stat.)  $\pm 0.2$  (syst.) GeV**
- **Local Significance:  $4.86\sigma$  ( $p\_value = 5.8 \times 10^{-7}$ )**

# What Does Experiments Have To Say

["http://meetings.aps.org/Meeting/APR18/Session/U09.6"](http://meetings.aps.org/Meeting/APR18/Session/U09.6)



- Note: The results are taken from my thesis work and they are not approved by CMS yet. The analysis is still in review.

4/15/18

Suleyman Durgut

2

## Combined Result




- Do a simultaneous fit to both channels, with fixed signal shapes but floating mass value.
- **Best mass :  $18.4 \pm 0.1$  (stat.)  $\pm 0.2$  (syst.) GeV**
- **Local Significance:  $4.86\sigma$  ( $p_{\text{value}} = 5.8 \times 10^{-7}$ )**

- In order to calculate global significance, Look-Elsewhere-Effect must be taken into account. Lots of toy MC generations are required, not an efficient method.
- Global significance is calculated using Gross-Vitells method which is used in Higgs discovery.

[Eur.Phys.J.C70:525-530,2010](#)


- **The returned global significance was  $3.6\sigma$ .**

# Model Predictions for $0^{++} 2b2\bar{b}$ tetraquark





# Model Predictions for $0^{++} 2b2\bar{b}$ tetraquark



- Results Very Model Dependent!!
- Not from first-principles
- Inconclusive whether tetraquark bound or not?

# What Is In The Literature (as of Dec 2017)?

Reference	Model
1110.1867	Diquark
1710.0254	Diquark
1605.01647	Sum-rules
1701.04285	Sum-rules
1710.0254	Schrodinger
1612.00012	Schrodinger
1605.01134	Pheno.
1611.00348	Pheno.
1703.00783	String
1709.0965	Production
1710.02738	Production

# Quantum Chromodynamics

*“The fundamental theory of the strong nuclear force”*

---

- QCD is non-perturbative
  - Solution: ‘solve’ numerically via LQCD

# Quantum Chromodynamics

*“The fundamental theory of the strong nuclear force”*

---


- QCD is non-perturbative
  - Solution: ‘solve’ numerically via LQCD
- The Lattice is **not** a model: the Lattice is a UV regulator of a QFT
- Lattice QCD results are from first-principles

# Quantum Chromodynamics

*“The fundamental theory of the strong nuclear force”*

- QCD is non-perturbative
  - Solution: ‘solve’ numerically via LQCD
- The Lattice is **not** a model: the Lattice is a UV regulator of a QFT
- Lattice QCD results are from first-principles

$$\mathcal{L}_{\text{QCD}} = \bar{\psi}_f (i \not{D} - m_f) \psi_f - \frac{1}{4} \text{tr} (GG)$$





# Quantum Chromodynamics

*“The fundamental theory of the strong nuclear force”*

- QCD is non-perturbative
  - Solution: ‘solve’ numerically via LQCD
- The Lattice is **not** a model: the Lattice is a UV regulator of a QFT
- Lattice QCD results are from first-principles

$$\mathcal{L}_{\text{QCD}} = \bar{\psi}_f (i \not{D} - m_f) \psi_f - \frac{1}{4} \text{tr} (GG)$$




# Quantum Chromodynamics

*“The fundamental theory of the strong nuclear force”*

- QCD is non-perturbative
  - Solution: ‘solve’ numerically via LQCD
- The Lattice is **not** a model: the Lattice is a UV regulator of a QFT
- Lattice QCD results are from first-principles

$$\mathcal{L}_{\text{QCD}} = \bar{\psi}_f (i \not{D} - m_f) \psi_f - \frac{1}{4} \text{tr} (GG)$$



??



??

# Lattice QCD Methodology

---

Down  
the  
Rabbit  
Hole






# Searching for Beauty-fully Bound Tetraquarks Using Lattice NRQCD

Ciaran Hughes, Estia Eichten, Christine Davies

---

- 📌 This talk will be a bigger picture sketch of results from [arxiv:1710.03236](https://arxiv.org/abs/1710.03236)
- 📌 For more details, please contact me ([chughes@fnal.gov](mailto:chughes@fnal.gov))!



# A (Bigger Picture) Lattice Spectrum Calculation: The Two-Point Correlator $C_{ab}^{2\text{pt.}}(t) = \langle 0 | \mathcal{O}_b(y_4) \mathcal{O}_a^\dagger(x_4) | 0 \rangle$

---

$$C_{ab}^{2\text{pt.}}(x_4 - y_4) = \langle \quad \rangle$$
The diagram shows a large, empty, light-blue rounded rectangle at the top of the page. Below it, a large white rounded rectangle with a black border contains the equation  $C_{ab}^{2\text{pt.}}(x_4 - y_4) = \langle \quad \rangle$ . The two large angle brackets are drawn with thick black lines, representing the vacuum expectation value of two operators.

# A (Bigger Picture) Lattice Spectrum Calculation: The Two-Point Correlator $C_{ab}^{2\text{pt.}}(t) = \langle 0 | \mathcal{O}_b(y_4) \mathcal{O}_a^\dagger(x_4) | 0 \rangle$


- 👤 Feynman Path Integral Approach to QFT: (Numerically) Integrate over all field configurations

$$C_{ab}^{2\text{pt.}}(x_4 - y_4) = \langle \text{Diagram} \rangle$$


# A (Bigger Picture) Lattice Spectrum Calculation: The Two-Point Correlator $C_{ab}^{2pt.}(t) = \langle 0 | \mathcal{O}_b(y_4) \mathcal{O}_a^\dagger(x_4) | 0 \rangle$

- Feynman Path Integral Approach to QFT: (Numerically) Integrate over all field configurations
  - Create some (superposition of) state with  $\mathcal{O}_a^\dagger(x_4)$


$$C_{ab}^{2pt.}(x_4 - y_4) = \langle \quad \rangle$$



$$t = x_4$$


# A (Bigger Picture) Lattice Spectrum Calculation: The Two-Point Correlator $C_{ab}^{2pt.}(t) = \langle 0 | \mathcal{O}_b(y_4) \mathcal{O}_a^\dagger(x_4) | 0 \rangle$

- Feynman Path Integral Approach to QFT: (Numerically) Integrate over all field configurations
  - Create some (superposition of) state with  $\mathcal{O}_a^\dagger(x_4)$
  - Propagate elementary fields for time  $t$



# A (Bigger Picture) Lattice Spectrum Calculation: The Two-Point Correlator $C_{ab}^{2pt.}(t) = \langle 0 | \mathcal{O}_b(y_4) \mathcal{O}_a^\dagger(x_4) | 0 \rangle$


- Feynman Path Integral Approach to QFT: (Numerically) Integrate over all field configurations
  - Create some (superposition of) state with  $\mathcal{O}_a^\dagger(x_4)$
  - Propagate elementary fields for time  $t$
  - Destroy some (superposition of) state with  $\mathcal{O}_b(y_4)$



# A (Bigger Picture) Lattice Spectrum Calculation: The Two-Point Correlator $C_{ab}^{2pt.}(t) = \langle 0 | \mathcal{O}_b(y_4) \mathcal{O}_a^\dagger(x_4) | 0 \rangle$

## Hilbert Space Formalism:

- Insert a complete set of QCD eigenstates




# A (Bigger Picture) Lattice Spectrum Calculation: The Two-Point Correlator $C_{ab}^{2pt.}(t) = \langle 0 | \mathcal{O}_b(y_4) \mathcal{O}_a^\dagger(x_4) | 0 \rangle$

## Hilbert Space Formalism:

- Insert a complete set of QCD eigenstates

$$C_{ab}^{2pt.}(t, \mathbf{P}) \equiv \langle 0 | \mathcal{O}_b(t, \mathbf{P}) \mathcal{O}_a^\dagger(0, \mathbf{P}) | 0 \rangle = \sum_n Z_{b,n} Z_{a,n}^\dagger e^{-E_n t}$$






# A (Bigger Picture) Lattice Spectrum Calculation: The Two-Point Correlator $C_{ab}^{2pt.}(t) = \langle 0 | \mathcal{O}_b(y_4) \mathcal{O}_a^\dagger(x_4) | 0 \rangle$

## Hilbert Space Formalism:

- Insert a complete set of QCD eigenstates

$$C_{ab}^{2pt.}(t, \mathbf{P}) \equiv \langle 0 | \mathcal{O}_b(t, \mathbf{P}) \mathcal{O}_a^\dagger(0, \mathbf{P}) | 0 \rangle = \sum_n Z_{b,n} Z_{a,n}^\dagger e^{-E_n t}$$



Extract QCD Energy Eigenstates



# Operators Used for $0^{++} 2b2\bar{b}$ State

$0^{++}$	
source	sink
$\mathcal{O}_{(n_b, \bar{n}_b)}^{A_1}$	


$$C_{ab}^{2\text{pt.}}(x_4 - y_4) =$$



# Operators Used for $0^{++} \ 2b2\bar{b}$ State

$0^{++}$	
source	sink
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	


$$C_{ab}^{2\text{pt.}}(x_4 - y_4) =$$



# Operators Used for $0^{++} 2b2\bar{b}$ State

$0^{++}$	
source	sink
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$



$$C_{ab}^{2\text{pt.}}(x_4 - y_4) =$$



# Operators Used for $0^{++} 2b2\bar{b}$ State

$0^{++}$	
source	sink
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$


$$C_{ab}^{2\text{pt.}}(x_4 - y_4) =$$



# Operators Used for $0^{++} 2b2\bar{b}$ State

$0^{++}$	
source	sink
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$
$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$


$$C_{ab}^{2\text{pt.}}(x_4 - y_4) =$$



# Operators Used for $0^{++} 2b2\bar{b}$ State

$0^{++}$	
source	sink
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$
$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$
$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$	$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$

$$C_{ab}^{2\text{pt.}}(x_4 - y_4) =$$



# Fierz Relations

Table 1: Fierz relations in the  $\bar{b}bbb$  system relating the two-meson and the diquark-antidiquark bilinears.


$J^{PC}$	Diquark-AntiDiquark	Two-Meson
$0^{++}$	$\bar{3}_c \times 3_c$	$-\frac{1}{2} 0; \Upsilon\Upsilon\rangle + \frac{\sqrt{3}}{2} 0; \eta_b\eta_b\rangle$
$0^{++}$	$6_c \times \bar{6}_c$	$\frac{\sqrt{3}}{2} 0; \Upsilon\Upsilon\rangle + \frac{1}{2} 0; \eta_b\eta_b\rangle$
$1^{+-}$	$\bar{3}_c \times 3_c$	$\frac{1}{\sqrt{2}}( 1; \Upsilon\eta_b\rangle +  1; \eta_b\Upsilon\rangle)$
$2^{++}$	$\bar{3}_c \times 3_c$	$ 2; \Upsilon\Upsilon\rangle$



# Operators Used for $0^{++} \mathbf{2b2\bar{b}}$ State

$0^{++}$	
source	sink
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$
$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$
$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$	$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$
$\mathcal{O}_{(D_{3_c}, A_{3_c})}^{A_1}$	$\mathcal{O}_{(D_{3_c}, A_{3_c})}^{A_1}$



$$C_{ab}^{2\text{pt.}}(x_4 - y_4) =$$



# Operators Used for $0^{++} 2b2\bar{b}$ State

$0^{++}$	
source	sink
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$
$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$	$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$
$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$	$\mathcal{O}_{(\eta_b, \eta_b)}^{A_1}$
$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$	$\mathcal{O}_{(\Upsilon, \Upsilon)}^{A_1}$
$\mathcal{O}_{(D_{3c}, A_{3c})}^{A_1}$	$\mathcal{O}_{(D_{3c}, A_{3c})}^{A_1}$
$\mathcal{O}_{(D_{6c}, A_{6c})}^{A_1}$	$\mathcal{O}_{(D_{6c}, A_{6c})}^{A_1}$

$$C_{ab}^{2\text{pt.}}(x_4 - y_4) =$$





show me the data!

- 📍 We perform a Bayesian fit to all the data within a certain channel

show me the data!

- 📍 We perform a Bayesian fit to all the data within a certain channel
- 📍 But you want to see the actual data! What can we easily show?

# The Lattice Effective Mass

---

$$aE^{\text{eff}} = \log \left( \frac{C(t)}{C(t+1)} \right)$$

# The Lattice Effective Mass

---

$$aE^{\text{eff}} = \log \left( \frac{C(t)}{C(t+1)} \right)$$

$$= aE_0 + \frac{Z_1^2}{Z_0^2} e^{-(E_1 - E_0)t} (1 - e^{-(E_1 - E_0)}) + \dots$$

# The Lattice Effective Mass

$$aE^{\text{eff}} = \log \left( \frac{C(t)}{C(t+1)} \right)$$

$$= aE_0 + \frac{Z_1^2}{Z_0^2} e^{-(E_1 - E_0)t} (1 - e^{-(E_1 - E_0)}) + \dots$$

*Excited State Contributions  
Exponentially decay away with  
time*

# The Lattice Effective Mass

$$aE^{\text{eff}} = \log \left( \frac{C(t)}{C(t+1)} \right)$$

$$= aE_0 + \frac{Z_1^2}{Z_0^2} e^{-(E_1 - E_0)t} (1 - e^{-(E_1 - E_0)}) + \dots$$

$$\xrightarrow{t \rightarrow \infty} aE_0$$

*Excited State Contributions  
Exponentially decay away with  
time*




# The Lattice Effective Mass

$$aE^{\text{eff}} = \log \left( \frac{C(t)}{C(t+1)} \right)$$

$$= aE_0 + \frac{Z_1^2}{Z_0^2} e^{-(E_1-E_0)t} (1 - e^{-(E_1-E_0)}) + \dots$$

$$\xrightarrow{t \rightarrow \infty} aE_0$$

*Excited State Contributions  
Exponentially decay away with  
time*




# Fake $0^{++}$ data on the $a \approx 0.06$ fm ensemble

*“What might you expect to see?”*

---

# Fake $0^{++}$ data on the $a \approx 0.06$ fm ensemble


*“What might you expect to see?”*



*The non-interacting  
 $2\eta_b$  and  $2\Upsilon$   
thresholds for reference*

# Fake $0^{++}$ data on the $a \approx 0.06$ fm ensemble


*“What might you expect to see?”*



*If there was no new stable tetraquark*

# Fake $0^{++}$ data on the $a \approx 0.06$ fm ensemble


*“What might you expect to see?”*



*If there was ONLY a new state 100 MeV below threshold*

# Fake $0^{++}$ data on the $a \approx 0.06$ fm ensemble


*“What might you expect to see?”*



*If the new state was 100 MeV below threshold and had  $Z_0 = Z_1$*

# Fake $0^{++}$ data on the $a \approx 0.06$ fm ensemble


*“What might you expect to see?”*



*If the new state was 100 MeV below threshold and had different  $Z_0, Z_1$*

# Fake $0^{++}$ data on the $a \approx 0.06$ fm ensemble

*“What might you expect to see?”*




*If the new state was 100 MeV below threshold and had different  $Z_0, Z_1$*

*If tetraquark exists we should see a fall below threshold and a clean signal as in blue curve!*




# The $0^{++}$ data on the $a \approx 0.06$ fm ensemble






For this talk, focus on red data points

Then compare to  $2\eta_b$  non-interacting threshold to determine binding





# The $0^{++}$ data on the $a \approx 0.06$ fm ensemble





# The $0^{++}$ data on the $a \approx 0.06$ fm ensemble




# The $0^{++}$ data on the $a \approx 0.06$ fm ensemble




# The $0^{++}$ data on the $a \approx 0.06$ fm ensemble



# Summary of Energies from Lattice




# Summary of Energies from Lattice



*No evidence of  $0^{++}$  below  $2\eta_b$  threshold*

# Summary of Energies from Lattice



*No evidence of  $0^{++}$  below  $2\eta_b$  threshold*

*If you don't observe a process, need to determine a bound, e.g, proton decay.*



# Bound on $0^{++} 2b2\bar{b}$ state to be stable

*"How would it have missed?"*

📌 If stable tetraquark exists, at a particular time  $t^*$ ,

$$C(t^*) = |\langle 0 | \mathcal{O} | 4b \rangle|^2 e^{-aE_{4b}t^*} + |\langle 0 | \mathcal{O} | 2\eta_b \rangle|^2 e^{-aE_{2\eta_b}t^*}$$

$$= Z_{4b}^2 e^{-aE_{4b}t^*} + \tilde{Z}_{2\eta_b}^2 e^{-aE_{2\eta_b}t^*}$$

# Bound on $0^{++} 2b2\bar{b}$ state to be stable

*"How would it have missed?"*

📌 If stable tetraquark exists, at a particular time  $t^*$ ,

$$C(t^*) = |\langle 0 | \mathcal{O} | 4b \rangle|^2 e^{-aE_{4b}t^*} + |\langle 0 | \mathcal{O} | 2\eta_b \rangle|^2 e^{-aE_{2\eta_b}t^*}$$

Input Data

$$= Z_{4b}^2 e^{-aE_{4b}t^*} + \tilde{Z}_{2\eta_b}^2 e^{-aE_{2\eta_b}t^*}$$

# Bound on $0^{++} 2b2\bar{b}$ state to be stable

*"How would it have missed?"*

📌 If stable tetraquark exists, at a particular time  $t^*$ ,

$$C(t^*) = |\langle 0 | \mathcal{O} | 4b \rangle|^2 e^{-aE_{4b}t^*} + |\langle 0 | \mathcal{O} | 2\eta_b \rangle|^2 e^{-aE_{2\eta_b}t^*}$$

Input Data

$$= Z_{4b}^2 e^{-aE_{4b}t^*} + \tilde{Z}_{2\eta_b}^2 e^{-aE_{2\eta_b}t^*}$$

Output Constraint

# Bound on $0^{++} 2b2\bar{b}$ state to be stable

*"How would it have missed?"*


If stable tetraquark exists, at a particular time  $t^*$ ,

$$C(t^*) = |\langle 0 | \mathcal{O} | 4b \rangle|^2 e^{-aE_{4b}t^*} + |\langle 0 | \mathcal{O} | 2\eta_b \rangle|^2 e^{-aE_{2\eta_b}t^*}$$

Input Data


$$= Z_{4b}^2 e^{-aE_{4b}t^*} + \tilde{Z}_{2\eta_b}^2 e^{-aE_{2\eta_b}t^*}$$

Output Constraint




# Summary

In Summary, lattice QCD finds no evidence of a stable  $2b2\bar{b}$  tetraquark



# What The Models Need!


	Have quarks as elementary particles?	Full $2 \times 2$ potential matrix (including mixing between different color components)?	Both Short and Long distance effects in Gluon exchange?



# What The Models Need!


*Diquarks*

	Have quarks as elementary particles?	Full $2 \times 2$ potential matrix (including mixing between different color components)?	Both Short and Long distance effects in Gluon exchange?
1110.1867	✗	✓	✗
1710.0254	✗	✓	✓



# What The Models Need!


		Have quarks as elementary particles?	Full $2 \times 2$ potential matrix (including mixing between different color components)?	Both Short and Long distance effects in Gluon exchange?
<i>Diquarks</i>	1110.1867	✗	✓	✗
	1710.0254	✗	✓	✓
<i>Sum-Rules</i>	1605.01647	✓	✓	✗
	1701.04285	✓	✓	✗






# What The Models Need!

		Have quarks as elementary particles?	Full $2 \times 2$ potential matrix (including mixing between different color components)?	Both Short and Long distance effects in Gluon exchange?
<i>Diquarks</i>	1110.1867	✗	✓	✗
	1710.0254	✗	✓	✓
<i>Sum-Rules</i>	1605.01647	✓	✓	✗
	1701.04285	✓	✓	✗
<i>Schrodinger Equation</i>	1710.0254	✓	✗	✗
	1612.00012	✓	✗	✓




# What The Models Need!

		Have quarks as elementary particles?	Full $2 \times 2$ potential matrix (including mixing between different color components)?	Both Short and Long distance effects in Gluon exchange?
<i>Diquarks</i>	1110.1867	✗	✓	✗
	1710.0254	✗	✓	✓
<i>Sum-Rules</i>	1605.01647	✓	✓	✗
	1701.04285	✓	✓	✗
<i>Schrodinger Equation</i>	1710.0254	✓	✗	✗
	1612.00012	✓	✗	✓
<i>Pheno.</i>	1605.01134	✓	✗	✗
	1611.00348	✓	✗	✓




# What The Models Need!

		Have quarks as elementary particles?	Full $2 \times 2$ potential matrix (including mixing between different color components)?	Both Short and Long distance effects in Gluon exchange?
<i>Diquarks</i>	1110.1867	✗	✓	✗
	1710.0254	✗	✓	✓
<i>Sum-Rules</i>	1605.01647	✓	✓	✗
	1701.04285	✓	✓	✗
<i>Schrodinger Equation</i>	1710.0254	✓	✗	✗
	1612.00012	✓	✗	✓
<i>Pheno.</i>	1605.01134	✓	✗	✗
	1611.00348	✓	✗	✓
<i>String</i>	1703.00783	✓	✓	✗



# What The Models Need!


	N.B., The lattice is <b>NOT</b> a model	Have quarks as elementary particles?	Full $2 \times 2$ potential matrix (including mixing between different color components)?	Both Short and Long distance effects in Gluon exchange?
<i>Diquarks</i>	1110.1867	✗	✓	✗
	1710.0254	✗	✓	✓
<i>Sum-Rules</i>	1605.01647	✓	✓	✗
	1701.04285	✓	✓	✗
<i>Schrodinger Equation</i>	1710.0254	✓	✗	✗
	1612.00012	✓	✗	✓
<i>Pheno.</i>	1605.01134	✓	✗	✗
	1611.00348	✓	✗	✓
<i>String</i>	1703.00783	✓	✓	✗



# Possible But Not Probable Future Work

---


*Fictional Heavy tetraquarks*



$2Q_1 2\bar{Q}_2$   
*tetraquarks*


# Possible But Not Probable Future Work

*Fictional Heavy tetraquarks*




$2Q_1 2\bar{Q}_2$   
tetraquarks

*tetraquark resonances*




# Possible But Not Probable Future Work

Fictional Heavy tetraquarks




$2Q_1 2\bar{Q}_2$   
tetraquarks

tetraquark resonances



Stable Real tetraquarks




$bb\bar{u}\bar{d}$   
tetraquarks

arxiv:1707.09575

# Ongoing Work


---






# Ongoing Work




*Nucleon properties*



*BSM*



# Ongoing Work



# Thank you!

---




Thank You to Raul Briceno for slide template  
and pretty graphics!

# Back-Up Slides


# Diquark-Antidiquark Wick Contractions

---







# Two-Meson Wick Contractions





---



# Two-Meson Wick Contractions





# Diquark-Antidiquark Wick Contractions





# Individual Wick Contraction Correlator Data



# Correlator Data With Harmonic Oscillator

- Add to the NRQCD Hamiltonian the harmonic oscillator scalar potential

$$\delta H_{HO} = \frac{m_b \omega^2}{2} |\mathbf{x} - \mathbf{x}_0|^2$$

This would bind a hypothetical compact tetraquark more, relative to the lowest threshold, and hence this hypothetical tetraquark would show up more easily in our calculation

# Correlator Data With Harmonic Oscillator



$$C_{i,j}^{JPC}(t, \omega) = \sum_n \frac{Z_n^i Z_n^{j,*}}{(1 + e^{-2\omega t})^{\frac{3}{2}}} e^{-(M(\omega)_n + \frac{3}{2}\omega)t}$$

$$C_{i,j}^{JPC}(t, \omega) = \sum_{X_2} Z_{X_2}^i Z_{X_2}^{j,*} \left( \frac{2\omega\mu_r\pi^{-1}}{1 - e^{-4\omega t}} \right)^{\frac{3}{2}}$$


$$\times e^{-(M_1^S(\omega) + M_2^S(\omega) + 3\omega)t} + \dots$$

The single and two-particle correlators get modified in the presence of the HO

$\eta_b \rightarrow \eta_b$




# Correlator Data With Harmonic Oscillator




🕒 No indication of a new bound state despite the addition of the scalar potential!!!

# Individual Wick Contraction Correlator Data HO

$D1 : 2\eta_b \rightarrow 2\eta_b$



$X2 : 2\eta_b \rightarrow 2\eta_b$



# Lattice QCD Methodology

---

4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

# Lattice QCD Methodology

---


4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}U \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S} \mathcal{O}[U, \psi, \bar{\psi}]$$

# Lattice QCD Methodology

4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}U \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S} \mathcal{O}[U, \psi, \bar{\psi}]$$

$$\approx \frac{1}{N} \sum_{i=1}^N \mathcal{O}[G^{(i)}]$$




# Lattice QCD Methodology


4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}U \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S} \mathcal{O}[U, \psi, \bar{\psi}]$$

$$\approx \frac{1}{N} \sum_{i=1}^N \mathcal{O}[G^{(i)}]$$

- where the integral is approximated as a sum over configurations  $\{G^{(i)}\}$  distributed according to the probability density:  $\exp(-S_{YM}) \prod \det(D + m_q)$

# Bottomonium Elastic Scattering States in FV



*“only a discrete number of modes  
can exist in a finite volume”*

*Spatially Periodic Box:*

$$p_i \in \frac{2\pi}{L} \times \mathbb{Z}$$

# Bottomonium Elastic Scattering States in FV

$$E(X^2) = \sqrt{M_1^2 + |\mathbf{k}|^2} + \sqrt{M_2^2 + |\mathbf{k}|^2}$$
$$\approx M_1^S + M_2^S + \frac{|\mathbf{k}|^2}{2\mu_r}$$

where we have defined the static, kinetic and reduced masses by  $M^S$ ,  $M^K$  and  $\mu_r = M_1^K M_2^K / (M_1^K + M_2^K)$

back-to-back states on our ensembles. As an example, examining the  $a = 0.09$  fm ensemble, and taking  $M_{\eta_b} = 9.399(2)$  GeV from the PDG [4], the smallest allowed  $|\mathbf{k}|^2/2\mu_r \approx 20$  MeV or 0.0092 in lattice units with all other back-to-back states separated by multiples of

$$C(t) = \sum_{X^2} \int \frac{d^3k}{(2\pi)^3} Z_{X^2}(\mathbf{k})^2 e^{-E(X^2)t}$$

$$C(t) = \sum_{X^2} e^{-(M_1^S + M_2^S)t} \sum_k \left\{ \sum_{i=0}^{\infty} Z_{X^2}^{2i} \frac{|\mathbf{k}|^{2i}}{\mu_r^{2i}} \right\} e^{-\frac{|\mathbf{k}|^2}{2\mu_r} t}$$

(A5)

# Bottomonium Elastic Scattering States in FV



*When does the two-body scattering states look like a continuum within stat. precision?*

$$I^{(l)}(t) = \frac{1}{\mu_r^{2l}} \int_{-\infty}^{\infty} d|\mathbf{k}| |\mathbf{k}|^{2l+2} e^{-\frac{|\mathbf{k}|^2}{2\mu_r} t}$$
$$D^{(l)}(t) = \frac{1}{\mu_r^{2l}} \sum_{|\mathbf{k}|} |\mathbf{k}|^{2l+2} e^{-\frac{|\mathbf{k}|^2}{2\mu_r} t}.$$

$$\left| \frac{\sum_{l=0}^{l_{max}} Z^{2,(l)} I^{(l)}(t) - \sum_{l=0}^{\infty} Z^{2,(l)} D^{(l)}(t)}{\sum_{l=0}^{l_{max}} Z^{2,(l)} I^{(l)}(t)} \right|$$
$$\leq \sum_{l=0}^{l_{max}} \frac{|I^{(l)}(t) - D^{(l)}(t)|}{I^{(0)}} + \sum_{l=l_{max}+1}^{\infty} \frac{D^{(l)}(t)}{I^{(0)}}$$
$$\leq \frac{\delta C(t)}{C(t)}$$

# Bottomonium Elastic Scattering States in FV

*When does the two-body scattering states look like a continuum within stat. precision?*



*After 1 fm!!*

# Lattice for Near Term Quantum Computing Era

📌 How many qubits are needed for SU(3) glue and how can lattice help?

📌 10x10x10 Spatial volume dof

*Lattice methodological techniques can reduce the number of qubits needed: Symanzik Improvement, lattice symmetries, etc*

📌 8 Gluon dof

📌  $x$  Occupation number dof (Not known)

*Lattice can help find  $x$*

📌  $x^{8000}$  Total dof  $\implies 8000 \log_2 x$  qubits

*Currently*

- 📌 *Small number of qubits*
- 📌 *Sparse qubit connectivity*
- 📌 *Noisy quantum gates*
- 📌 *Short coherence times*

*Lattice can be used as a testbed for tech, and determined ideal initial wave function (remove adiabatic starting) to remove gate fidelity.*

*Arxiv:1803.03326v2*

📌 *State of the art: 1+1 Schwinger model: 6 qubits reduced to 2 qubits*

# QCD - The Spectrum Enigma

*"We don't understand XYZ since 2003"*

---


# QCD - The Spectrum Enigma

“We don’t understand XYZ since 2003”

TABLE 9: As in Table 4, but for new *unconventional* states in the  $c\bar{c}$  and  $b\bar{b}$  regions, ordered by mass. For  $X(3872)$ , the values given are based only upon decays to  $\pi^+\pi^-J/\psi$ .  $X(3945)$  and  $Y(3940)$  have been subsumed under  $X(3915)$  due to compatible properties. The state known as  $Z(3930)$  appears as the  $\chi_{c2}(2P)$  in Table 4. See also the reviews in [81–84]

State	$m$ (MeV)	$\Gamma$ (MeV)	$J^{PC}$	Process (mode)	Experiment ( $\#\sigma$ )	Year	Status
$X(3872)$	$3871.52 \pm 0.20$	$1.3 \pm 0.6$ ( $< 2.2$ )	$1^{++}/2^{-+}$	$B \rightarrow K(\pi^+\pi^-J/\psi)$ $p\bar{p} \rightarrow (\pi^+\pi^-J/\psi) + \dots$ $B \rightarrow K(\omega J/\psi)$ $B \rightarrow K(D^{*0}\bar{D}^0)$ $B \rightarrow K(\gamma J/\psi)$ $B \rightarrow K(\gamma\psi(2S))$	Belle [85, 86] (12.8), BABAR [87] (8.6) CDF [88–90] (np), DØ [91] (5.2) Belle [92] (4.3), BABAR [93] (4.0) Belle [94, 95] (6.4), BABAR [96] (4.9) Belle [92] (4.0), BABAR [97, 98] (3.6) BABAR [98] (3.5), Belle [99] (0.4)	2003	OK
$X(3915)$	$3915.6 \pm 3.1$	$28 \pm 10$	$0/2^{2+}$	$B \rightarrow K(\omega J/\psi)$ $e^+e^- \rightarrow e^+e^-(\omega J/\psi)$	Belle [100] (8.1), BABAR [101] (19) Belle [102] (7.7)	2004	OK
$X(3940)$	$3942_{-8}^{+9}$	$37_{-17}^{+27}$	$?^{2+}$	$e^+e^- \rightarrow J/\psi(D\bar{D}^*)$ $e^+e^- \rightarrow J/\psi(\dots)$	Belle [103] (6.0) Belle [54] (5.0)	2007	NC!
$G(3900)$	$3943 \pm 21$	$52 \pm 11$	$1^{--}$	$e^+e^- \rightarrow \gamma(D\bar{D})$	BABAR [27] (np), Belle [21] (np)	2007	OK
$Y(4008)$	$4008_{-49}^{+121}$	$226 \pm 97$	$1^{--}$	$e^+e^- \rightarrow \gamma(\pi^+\pi^-J/\psi)$	Belle [104] (7.4)	2007	NC!
$Z_1(4050)^+$	$4051_{-43}^{+24}$	$82_{-55}^{+51}$	$?$	$B \rightarrow K(\pi^+\chi_{c1}(1P))$	Belle [105] (5.0)	2008	NC!
$Y(4140)$	$4143.4 \pm 3.0$	$15_{-7}^{+11}$	$?^{2+}$	$B \rightarrow K(\phi J/\psi)$	CDF [106, 107] (5.0)	2009	NC!
$X(4160)$	$4156_{-25}^{+29}$	$139_{-65}^{+113}$	$?^{2+}$	$e^+e^- \rightarrow J/\psi(D\bar{D}^*)$	Belle [103] (5.5)	2007	NC!
$Z_2(4250)^+$	$4248_{-45}^{+185}$	$177_{-72}^{+321}$	$?$	$B \rightarrow K(\pi^+\chi_{c1}(1P))$	Belle [105] (5.0)	2008	NC!
$Y(4260)$	$4263 \pm 5$	$108 \pm 14$	$1^{--}$	$e^+e^- \rightarrow \gamma(\pi^+\pi^-J/\psi)$ $e^+e^- \rightarrow (\pi^+\pi^-J/\psi)$ $e^+e^- \rightarrow (\pi^0\pi^0J/\psi)$	BABAR [108, 109] (8.0) CLEO [110] (5.4) Belle [104] (15) CLEO [111] (11) CLEO [111] (5.1)	2005	OK
$Y(4274)$	$4274.4_{-6.7}^{+8.4}$	$32_{-15}^{+22}$	$?^{2+}$	$B \rightarrow K(\phi J/\psi)$	CDF [107] (3.1)	2010	NC!
$X(4350)$	$4350.6_{-5.1}^{+4.6}$	$13.3_{-10.0}^{+18.4}$	$0, 2^{++}$	$e^+e^- \rightarrow e^+e^-(\phi J/\psi)$	Belle [112] (3.2)	2009	NC!
$Y(4360)$	$4353 \pm 11$	$96 \pm 42$	$1^{--}$	$e^+e^- \rightarrow \gamma(\pi^+\pi^-\psi(2S))$	BABAR [113] (np), Belle [114] (8.0)	2007	OK
$Z(4430)^+$	$4443_{-18}^{+24}$	$107_{-71}^{+113}$	$?$	$B \rightarrow K(\pi^+\psi(2S))$	Belle [115, 116] (6.4)	2007	NC!
$X(4630)$	$4634_{-11}^{+9}$	$92_{-32}^{+41}$	$1^{--}$	$e^+e^- \rightarrow \gamma(\Lambda_c^+\Lambda_c^-)$	Belle [25] (8.2)	2007	NC!
$Y(4660)$	$4664 \pm 12$	$48 \pm 15$	$1^{--}$	$e^+e^- \rightarrow \gamma(\pi^+\pi^-\psi(2S))$	Belle [114] (5.8)	2007	NC!
$Y_b(10888)$	$10888.4 \pm 3.0$	$30.7_{-7.7}^{+8.9}$	$1^{--}$	$e^+e^- \rightarrow (\pi^+\pi^-\Upsilon(nS))$	Belle [37, 117] (3.2)	2010	NC!

Quarkonium Working Group - 2010



S. Olsen - 2015




# Quantum Chromodynamics

*"We don't understand XYZ since 2003"*

- No general consensus for XYZ's despite being over a decade!!!
- Compact tetraquark
- Loosely bound molecular state
- Kinematical effect - Cusps
- Hydro-quarkonium
- Diquark-quarkonium
- Hybrids
- ....

Simplest Extended System




$2q2\bar{q}$   
tetraquarks

Take  
 $m_q \gg \Lambda_{\text{QCD}}$

# Lattice QCD Methodology

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$



# Lattice QCD Methodology

---

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$

*Complication:  $b$ -quarks do not fit on current lattices!!*

# Lattice QCD Methodology

---

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$

*Complication:  $b$ -quarks do not fit on current lattices!!*

*Solution: Use a Non-Relativistic Effective Field Theory to simulate the  $b$ -quarks*

# Lattice QCD Methodology

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length **L** and spacing **a**

*Complication:  $b$ -quarks do not fit on current lattices!!*

*Solution: Use a Non-Relativistic Effective Field Theory to simulate the  $b$ -quarks*

- Has Expansion Parameter  $v^2 \sim 0.1$
- N.B.: Matching Coefficients Need to be Calculated in Lattice Perturbation Theory

# Lattice QCD Methodology

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length **L** and spacing **a**

$$\begin{aligned} a\delta H &= aH_0 + a\delta H_{v^4} + a\delta H_{v^6}; \\ aH_0 &= -\frac{\Delta^{(2)}}{2am_b} \\ a\delta H_{v^4} &= -c_1 \frac{(\Delta^{(2)})^2}{8(am_b)^3} + c_2 \frac{i}{8(am_b)^2} \left( \nabla \cdot \tilde{\mathbf{E}} - \tilde{\mathbf{E}} \cdot \nabla \right) \\ &\quad - c_3 \frac{1}{8(am_b)^2} \sigma \cdot \left( \tilde{\nabla} \times \tilde{\mathbf{E}} - \tilde{\mathbf{E}} \times \tilde{\nabla} \right) \\ &\quad - c_4 \frac{1}{2am_b} \sigma \cdot \tilde{\mathbf{B}} + c_5 \frac{\Delta^{(4)}}{24am_b} - c_6 \frac{(\Delta^{(2)})^2}{16n(am_b)^2} \\ a\delta H_{v^6} &= -c_7 \frac{1}{8(am_b)^3} \left\{ \Delta^{(2)}, \sigma \cdot \tilde{\mathbf{B}} \right\} \\ &\quad - c_8 \frac{3}{64(am_b)^4} \left\{ \Delta^{(2)}, \sigma \cdot \left( \tilde{\nabla} \times \tilde{\mathbf{E}} - \tilde{\mathbf{E}} \times \tilde{\nabla} \right) \right\} \\ &\quad - c_9 \frac{i}{8(am_b)^3} \sigma \cdot \tilde{\mathbf{E}} \times \tilde{\mathbf{E}} \end{aligned}$$

# Lattice QCD Methodology

---

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length **L** and spacing **a**

# Lattice QCD Methodology

---

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$
2. Get one of these:





# Lattice QCD Methodology

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$
2. Get one of these:



3. Buy one of these:




# Lattice QCD Methodology

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$

2. Get one of these:



3. Buy one of these:



4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)!!

# Lattice QCD Methodology

---

4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

# Lattice QCD Methodology

---


4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}U \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S} \mathcal{O}[U, \psi, \bar{\psi}]$$

# Lattice QCD Methodology

4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}U \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S} \mathcal{O}[U, \psi, \bar{\psi}]$$

$$\approx \frac{1}{N} \sum_{i=1}^N \mathcal{O}[G^{(i)}]$$


# Lattice QCD Methodology

4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}U \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-S} \mathcal{O}[U, \psi, \bar{\psi}]$$

$$\approx \frac{1}{N} \sum_{i=1}^N \mathcal{O}[G^{(i)}]$$

- where the integral is approximated as a sum over configurations  $\{G^{(i)}\}$  distributed according to the probability density:  $\exp(-S_{YM}) \prod \det(D + m_q)$

# Lattice QCD Methodology

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$

2. Get one of these:



3. Buy one of these:



4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

# Lattice QCD Methodology

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$

2. Get one of these:



3. Buy one of these:



4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

5. Do all the computations/analysis



# Lattice QCD Methodology

1. Take Euclidean QCD and discretise it on a **Finite-Volume Lattice** of length  $L$  and spacing  $a$

2. Get one of these:



3. Buy one of these:



4. Numerically evaluate the Feynman Path Integral (the first-principles approach to QFT)

5. Do all the computations/analysis

6. Pay the Electricity Bill....