



Contribution ID: 252

Type: **Invited talk**

## Cornell Model calibration with NRQCD at N<sup>3</sup>LO

Thursday, 2 August 2018 15:40 (20 minutes)

The typical energy scale of heavy hadron spectroscopy makes the system accessible to perturbative calculations in terms of non-relativistic QCD. Within NRQCD the predictions of heavy quarkonium energy levels rely on the accurate description of the static QCD potential  $V_{\text{QCD}}(r)$ . Most recent calculations computed the energy levels of the lower-lying bottomonium states up to  $\mathcal{O}(\alpha_s^5 m)$  and  $\mathcal{O}(\alpha_s^5 m \log \alpha_s)$  utilizing pNRQCD [1]. A closed expression for arbitrary quantum numbers can be found in Ref [2].

Historically, the heavy quarkonium spectroscopy was studied using phenomenological approaches such as the Cornell model  $V_{\text{Cornell}} = -\kappa/r + \sigma r$ , which assumes a short-distance dominant Coulomb potential plus a linear rising potential that emerges at long distances. Such a model works satisfactorily in describing the charmonium and bottomonium spectroscopy. However, even when there are physically-motivated arguments for the construction of the Cornell model, there is no connection a priori between the model and QCD parameters.

Based on a previous work on heavy meson spectroscopy [3], we calibrate the Cornell model with NRQCD predictions for the lowest lying bottomonium states at N<sup>3</sup>LO, in which the bottom mass is varied within a wide range. We show that the Cornell model mass parameter can be identified with the low-scale short-distance MSR mass at the scale  $R = 1$  GeV. This identification holds for any value of  $\alpha_s$  or the bottom mass. Furthermore we show that a) the “string tension” parameter is completely independent of the bottom mass, and b) the Coulomb strength  $\kappa$  of the Cornell model can be related to the QCD strong coupling constant  $\alpha_s$  at a certain scale. Finally we show that for moderate values of  $r$ , the NRQCD and Cornell static potentials are in head-on agreement when switching the pole mass to the MSR scheme, which allows to simultaneously cancel the renormalon and sum up large logarithms.

[1] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Nucl. Phys. B 566, 275 (2000).

[2] Y. Kiyo and Y. Sumino, Nucl. Phys. B 889, 156 (2014).

[3] V. Mateu and P. G. Ortega, JHEP 1801 (2018) 122.

**Primary authors:** Dr GARCIA ORTEGA, Pablo (University of Salamanca); Dr MATEU BARREDA, Vicent (University of Salamanca); RODRIGUEZ ENTEM, David (University of Salamanca); FERNANDEZ, Francisco (Universidad de Salamanca)

**Presenter:** Dr GARCIA ORTEGA, Pablo (University of Salamanca)

**Session Classification:** Heavy quarks

**Track Classification:** C: Heavy quarks