XIIIth Quark Confinement and the Hadron Spectrum



Contribution ID: 42

Type: Talk

Pseudoscalar pole contribution to the hadronic light-by-light piece of a_{μ}

Thursday 2 August 2018 15:00 (20 minutes)

We studied the transition form factor involved in pseudoscalar meson (π,η,η') decays into two virtual photons by means of a chiral-invariant Lagrangian, considering the lowest-lying multiplet of vector and pseudoscalar resonances. Accounting for U(3) breaking effects, we give the most general corrections of order m_P^2 to the form factor. Most parameters are fixed requiring short-distance constraints. The remaining ones are fitted to experimental measurements of the form factors in the space-like $(q^2 < 0)$ region of photon momenta. We, thus, obtain the P-pole contribution to the hadronic light-by-light scattering of the muon g-2 with an improved certainty: $(8.47 \pm 0.16) \times 10^{-10}$. This is obtained neglecting BaBar data for the π^0 Transition Form Factor which, in our analysis, is in conflict with the remaining experimental inputs.

Primary authors: GUEVARA, Adolfo (Madrid University); SANZ-CILLERO, Juan José (Universidad Complutense de Madrid); ROIG GARCÉS, Pablo

Presenter: GUEVARA, Adolfo (Madrid University)

Session Classification: Light quarks

Track Classification: B: Light quarks