

XIIIth Confinement and the Hadron Spectrum

Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results
- Summary

Radiative corrections in Dalitz decays of π^0 , η and η' mesons

Tomáš Husek

IFIC, Universitat de València-CSIC

In collaboration with

K. Kampf, J. Novotný (Charles University), S. Leupold (Uppsala University)

Maynooth

August 2, 2018



Contents

Introduction

 π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

 π^0 Dalitz decay

Introduction

Radiative corrections

Results

 $\eta^{(\prime)}$ Dalitz decays

Introduction

Radiative corrections


Specific contributions

Results

Summary

Decay modes of neutral pion:

Process	Branching ratio
$\pi^0 \rightarrow \gamma\gamma$	$(98.823 \pm 0.034) \%$
$\pi^0 \rightarrow e^+e^-\gamma$	$(1.174 \pm 0.035) \%$
$\pi^0 \rightarrow e^+e^+e^-e^-$	$(3.34 \pm 0.16) \times 10^{-5}$
$\pi^0 \rightarrow e^+e^-$	$(6.46 \pm 0.33) \times 10^{-8}$

Rare decay $\pi^0 \rightarrow e^+e^-$

- interesting way to study low-energy (long-distance) dynamics in the SM
- systematic theoretical treatment dates back to *Drell, NC* (1959)
- suppressed in comparison to the decay $\pi^0 \rightarrow \gamma\gamma$ by a factor of $2(\alpha m_e/M_\pi)^2$
 - ↪ one-loop structure + helicity suppression
 - ↪ may be sensitive to possible effects of new physics

Contents

Introduction

 π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

 π^0 Dalitz decay

Introduction

Radiative corrections

Results

 $\eta^{(\prime)}$ Dalitz decays

Introduction

Radiative corrections

Specific contributions

Results

Summary

KTeV-E799-II experiment at Fermilab (*Abouzaid et al.*, PRD 75 (2007))
→ precise measurements of branching ratio $\pi^0 \rightarrow e^+e^-$ (794 candidates)

$$\frac{\Gamma(\pi^0 \rightarrow e^+e^-(\gamma), x > 0.95)}{\Gamma(\pi^0 \rightarrow e^+e^-\gamma, x > 0.232)} = (1.685 \pm 0.064 \pm 0.027) \times 10^{-4}$$

Extrapolate the Dalitz decay branching ratio to full range of x

$$B^{\text{KTeV}}(\pi^0 \rightarrow e^+e^-(\gamma), x_D > 0.95) = (6.44 \pm 0.25 \pm 0.22) \times 10^{-8}$$

- PDG average value $(6.46 \pm 0.33) \times 10^{-8}$ mainly based on this result
- extrapolate full radiative tail beyond $x > 0.95$ (*Bergström*, Z.Ph.C 20 (1983))
- scale the result back by the overall radiative corrections

→ final result for lowest order (no final state radiation)

$$B_{\text{KTeV}}^{\text{no-rad}}(\pi^0 \rightarrow e^+e^-) = (7.48 \pm 0.29 \pm 0.25) \times 10^{-8}$$

Comparison with SM prediction (*Dorokhov and Ivanov*, PRD 75 (2007))

$$B_{\text{SM}}^{\text{no-rad}}(\pi^0 \rightarrow e^+e^-) = (6.23 \pm 0.09) \times 10^{-8}$$

→ interpreted as 3.3σ discrepancy between theory and experiment

Contents

Introduction

 π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

 π^0 Dalitz decay

Introduction

Radiative corrections

Results

 $\eta^{(\prime)}$ Dalitz decays


Introduction

Radiative corrections

Specific contributions

Results

Summary



- very fashionable to ascribe eventual discrepancies to effects of new physics

BUT

- first, look for more conventional solution (i.e. within SM)
 - radiative corrections (usually very important)
 - form-factor modeling







Radiative corrections to $\pi^0 \rightarrow e^+e^-$

Two-loop virtual radiative corrections

Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results
- Summary

- calculated by Vaško and Novotný, JHEP 1110 (2011)




Radiative corrections to $\pi^0 \rightarrow e^+e^-$

Bremsstrahlung: photon emission from the outer fermion line


Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung**
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results
- Summary

- compensation of IR divergence in 2-loop contributions
 $\hookrightarrow TH, Kampf and Novotný, EPJC 74 (2014)$



- contain UV subdivergences \rightarrow counter-term tree diagrams with coupling χ



Radiative corrections to $\pi^0 \rightarrow e^+e^-$

Bremsstrahlung: photon emission from the inner fermion line (propagator)


Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results
- Summary

Do not forget the third, **box** diagram, necessary to satisfy the Ward identities

$$\mathcal{M}_{(\lambda)} = \varepsilon_{(\lambda)}^{*\rho}(k) \mathcal{M}_\rho^{\text{BS}} \longrightarrow k^\rho \mathcal{M}_\rho^{\text{BS}} = 0$$

- finite contribution to bremsstrahlung amplitude



Contents

[Introduction](#) [\$\pi^0\$ rare decay](#)[Introduction](#)[Radiative corrections](#)[Virtual corrections](#)[Bremsstrahlung](#)[Results](#) [\$\pi^0\$ Dalitz decay](#)[Introduction](#)[Radiative corrections](#)[Results](#) [\$\eta^{\(\prime\)}\$ Dalitz decays](#)[Introduction](#)[Radiative corrections](#)[Specific contributions](#)[Results](#)[Summary](#)Size of the radiative corrections (**newly calculated**)

$$\delta^{\text{NLO}}(0.95) \equiv \delta^{\text{virt.}} + \delta^{\text{BS}}(0.95) = (-5.5 \pm 0.2)\%$$

- can be thought as model-independent
- differs significantly from previous approximate calculations

Bergström, Z.Ph.C 20 (1983): $\delta(0.95) = -13.8\%$

Dorokhov et al., EPJC 55 (2008): $\delta(0.95) = -13.3\%$

- original KTeV vs. SM discrepancy reduced to the 2σ level or less
- contact interaction coupling finite part set to

$$\chi_{\text{LMD}}^{(r)}(M_\rho) = 2.2 \pm 0.9$$

Contents

[Introduction](#) [\$\pi^0\$ rare decay](#)[Introduction](#)[Radiative corrections](#)[Virtual corrections](#)[Bremsstrahlung](#)[Results](#) [\$\pi^0\$ Dalitz decay](#)[Introduction](#)[Radiative corrections](#)[Results](#) [\$\eta^{\(\prime\)}\$ Dalitz decays](#)[Introduction](#)[Radiative corrections](#)[Specific contributions](#)[Results](#)[Summary](#)

Quantity really measured by KTeV

$$\left. \frac{\Gamma(\pi^0 \rightarrow e^+ e^- (\gamma), x > 0.95)}{\Gamma(\pi^0 \rightarrow e^+ e^- \gamma (\gamma), x > 0.2319)} \right|_{\text{KTeV}} = (1.685 \pm 0.064 \pm 0.027) \times 10^{-4}$$

→ Dalitz decay comes into play

- second most important decay channel of a neutral pion
 - ↪ branching ratio $(1.174 \pm 0.035)\%$
- first studied by *Richard H. Dalitz*, PPSA 64 (1951), whose name it carries
- experimental data of this process provide the information about singly virtual pion transition form factor $\mathcal{F}_{\pi^0 \gamma^* \gamma^*}(0, q^2)$
 - ↪ in particular about its slope parameter a_π

Contents

Introduction

 π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

 π^0 Dalitz decay

Introduction

Radiative corrections

Results

 $\eta^{(\prime)}$ Dalitz decays

Introduction

Radiative corrections

Specific contributions

Results

Summary

- radiative corrections to the total decay rate of the Dalitz decay
 - ↪ first addressed (numerically) by *Joseph*, NC 16 (1960)
- pioneering study of corrections to the differential decay rate
 - ↪ *Lautrup and Smith*, PRD 3 (1971)
 - ↪ soft-photon approximation
- extended by *Mikaelian and Smith*, PRD 5 (1972)
 - ↪ hard-photon corrections
 - ↪ whole range of bremsstrahlung photon energy
 - ↪ table of values

Radiative corrections to $\pi^0 \rightarrow e^+e^-\gamma$
Novel approach


Contents

Introduction
 π^0 rare decay
Introduction
Radiative corrections
Virtual corrections
Bremsstrahlung
Results

π^0 Dalitz decay
Introduction
Radiative corrections
Results

$\eta^{(\prime)}$ Dalitz decays
Introduction
Radiative corrections
Results

$\eta^{(\prime)}$ Dalitz decays
Introduction
Radiative corrections
Specific contributions
Results
Summary




- new calculations motivated by needs of NA48/NA62 experiments at CERN
 - ↪ measure the slope a_π of $\mathcal{F}_{\pi^0\gamma^*\gamma^*}(0, q^2)$: *Lazzeroni et al., PLB 768 (2017)*
- unlike before no approximation was used
 - ↪ can be used also for related decays $\eta \rightarrow \ell^+\ell^-\gamma$ etc.
- C++ code returns the correction for any given x and y
 - ↪ propagated into MC generator of NA62 experiment
- *TH, Kampf and Novotný, PRD 92 (2015)*

Radiative corrections to $\pi^0 \rightarrow e^+e^-\gamma$
Results

Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results

Summary



Radiative corrections to $\pi^0 \rightarrow e^+ e^- \gamma$

Precise calculation of ratio R

Contents

Introduction
π^0 rare decay
Introduction
Radiative corrections
Virtual corrections
Bremsstrahlung
Results
π^0 Dalitz decay
Introduction
Radiative corrections
Results
$\eta^{(\prime)}$ Dalitz decays
Introduction
Radiative corrections
Specific contributions
Results
Summary

Precise knowledge of $R \equiv \frac{\Gamma(\pi^0 \rightarrow e^+ e^- \gamma)}{\Gamma(\pi^0 \rightarrow \gamma\gamma)}$

↪ for small slope and up to NLO radiative corrections

$$R \simeq \frac{\alpha}{\pi} \iint (1 + \textcolor{brown}{a}_\pi x)^2 (1 + \delta(x, y)) \frac{(1-x)^3}{4x} \left[1 + y^2 + \frac{4m_e^2}{M_\pi^2 x} \right] dx dy$$

Conservative estimate for uncertainty (a_π , NNLO)

$$R = 1.1978(6) \%$$

↪ chosen $a_\pi = 3.55(70) \%$, covers

$$a_\pi^{\text{VMD}} = 3.0 \%, \quad a_\pi^{\text{PDG}} = 3.35(31) \%, \quad a_\pi^{\text{NA62}} = 3.68(57) \%$$

Constraint: $1 \simeq \mathcal{B}(\pi^0 \rightarrow \gamma\gamma) + \mathcal{B}(\pi^0 \rightarrow e^+ e^- \gamma(\gamma)) + \mathcal{B}(\pi^0 \rightarrow e^+ e^- e^+ e^-)$

$$\mathcal{B}(\pi^0 \rightarrow \gamma\gamma) = 98.8131(6) \%, \quad \mathcal{B}(\pi^0 \rightarrow e^+ e^- \gamma(\gamma)) = 1.1836(6) \%$$

TH, Goudzovski and Kampf, to appear on arXiv soon

PDG

$$R = 1.188(35) \%, \quad \mathcal{B}(\pi^0 \rightarrow \gamma\gamma) = 98.823(34) \%, \quad \mathcal{B}(\pi^0 \rightarrow e^+ e^- \gamma) = 1.174(35) \%$$

Contents

Introduction

 π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

 π^0 Dalitz decay

Introduction

Radiative corrections

Results

 $\eta^{(\prime)}$ Dalitz decays

Introduction

Radiative corrections

Specific contributions

Results

Summary

 $\eta^{(\prime)}$ Dalitz decays

- small branching ratios
 - ↪ hadronic decay modes are open
- access to electromagnetic transition form factors
 - ↪ $\eta^{(\prime)}$ -meson structure
 - ↪ valuable input for other quantities and e.g. $g - 2$ of a muon
 - ↪ radiative corrections crucial to extract relevant information from data

naive rad. corrections for $\eta \rightarrow e^+e^-\gamma$: *Mikaelian and Smith, PRD 5 2890 (1972)*

- numerical values correspond to simple change $M_{\pi^0} \rightarrow M_\eta$
 - ↪ π^0 case: *Mikaelian and Smith, PRD 5 1763 (1972)*
- fully inclusive radiative corrections
 - ↪ no momentum or angular cuts on the bremsstrahlung photon applied

Radiative corrections to $\eta^{(\prime)} \rightarrow \ell^+ \ell^- \gamma$ decays

What should be taken into account?

Contents

Introduction

π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

π^0 Dalitz decay

Introduction

Radiative corrections

Results

$\eta^{(\prime)}$ Dalitz decays

Introduction

Radiative corrections

Specific contributions
Results

Summary

The $\eta^{(\prime)}$ case compared to π^0

- larger rest mass

↪ M_η above muon-pair threshold: $M_\eta > 2m_\mu$

↪ $M_{\eta'}$ above lowest-lying resonances: $M_{\eta'} > M_\rho, M_\omega$

↪ sensitive to the widths of resonances

↪ ω narrow, ρ broad resonance in $\pi\pi$ scattering

- strange-flavor content

↪ quark-flavor basis

Feldmann et al., PLB 449 (1999), Escribano et al., JHEP 06 (2005)

$$j^\ell \equiv \frac{i}{2} [\bar{u}\gamma_5 u + \bar{d}\gamma_5 d], \quad j^s \equiv \frac{i}{\sqrt{2}} [\bar{s}\gamma_5 s]$$

- η - η' mixing: $\langle 0 | j^A | \eta^B \rangle = B_0 F_\pi f_A \delta^{AB}, \langle \eta^A | \eta^B \rangle = \delta^{AB}, A, B \in \{\ell, s\}$

$$|\eta\rangle = \cos \phi |\eta^\ell\rangle - \sin \phi |\eta^s\rangle$$

$$|\eta'\rangle = \sin \phi |\eta^\ell\rangle + \cos \phi |\eta^s\rangle$$

Contents

Introduction

 π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

 π^0 Dalitz decay

Introduction

Radiative corrections

Results

 $\eta^{(\prime)}$ Dalitz decays

Introduction

Radiative corrections

Specific contributions

Results

Summary

Full set of NLO QED radiative corrections:

TH, Kampf, Leupold and Novotný, PRD 97 (2018)

- compared to previous approach:

- ↪ muon loops + hadronic VP
 - ↪ $1\gamma\text{IR}$ at one-loop level
 - ↪ form-factor effects (also in BS)
 - ↪ higher orders in the final-state lepton mass **not** neglected

- general framework: **three** additional processes

- ↪ also muon decay modes

 η case: **most** of the ingredients in *TH, Kampf and Novotný, PRD 92 (2015)* η' case: real challenge

- ↪ resulting framework also **applicable** to the π^0 case (numerically compatible)

- ↪ overkill (correction to the correction of order 1%)

Radiative corrections to $\eta^{(\prime)} \rightarrow \ell^+ \ell^- \gamma$ decays

Virtual corrections

Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results
- Summary

Photon self-energy in the form $\Pi(s) = \Pi_L(s) + \Pi_H(s)$


- lepton loops (electrons and as well **muons**)

$$\Pi_L(M_P^2 x) = \frac{\alpha}{\pi} \sum_{\ell' = e, \mu} \left\{ \frac{8}{9} - \frac{\beta_{\ell'}^2}{3} + \left(1 - \frac{\beta_{\ell'}^2}{3} \right) \frac{\beta_{\ell'}}{2} \log [-\gamma_{\ell'} + i\epsilon] \right\}$$

- hadronic contribution

↪ **Jegerlehner, Z.Ph.C 32 (1986)**

$$\Pi_H(s) = -\frac{s}{4\pi^2 \alpha} \int_{4m_\pi^2}^\infty \frac{\sigma_H(s') ds'}{s - s' + i\epsilon}$$




$$\delta^{\text{virt}}(x, y) = \frac{1}{|1 + \Pi(M_P^2 x)|^2} - 1 + 2 \operatorname{Re} \left\{ F_1(x) + \frac{2F_2(x)}{1 + y^2 + \frac{\nu^2}{x}} \right\}$$

Radiative corrections to $\eta^{(\prime)} \rightarrow \ell^+ \ell^- \gamma$ decays

1 γ IR contribution at one-loop level

Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results
- Summary



1 γ IR contribution at one-loop level

- beyond effective approach
- we don't expect substantial model dependence \hookrightarrow VMD-inspired model
 \hookrightarrow strange-flavor content and η - η' mixing


$$\begin{aligned} e^2 \mathcal{F}_{\eta\gamma^*\gamma^*}^{\text{VMD}}(p^2, q^2) = & -\frac{N_c}{8\pi^2 F_\pi} \frac{2e^2}{3} \\ & \times \left[\frac{5}{3} \frac{\cos \phi}{f_\ell} \frac{M_{\omega/\rho}^4}{(p^2 - M_{\omega/\rho}^2)(q^2 - M_{\omega/\rho}^2)} - \frac{\sqrt{2}}{3} \frac{\sin \phi}{f_s} \frac{M_\phi^4}{(p^2 - M_\phi^2)(q^2 - M_\phi^2)} \right] \end{aligned}$$

Radiative corrections to $\eta^{(\prime)} \rightarrow \ell^+ \ell^- \gamma$ decays

Bremsstrahlung

Contents

Introduction
 π^0 rare decay
 Introduction
 Radiative corrections
 Virtual corrections
 Bremsstrahlung
 Results
 π^0 Dalitz decay
 Introduction
 Radiative corrections
 Results
 $\eta^{(\prime)}$ Dalitz decays
 Introduction
 Radiative corrections
 Specific contributions
 Results
 Summary



- slope not negligible
- for η : expansion in slope a would be **still** (somewhat) suitable

$$\mathcal{F}((p_\gamma + p_{e^+} + p_{e^-})^2) \simeq \mathcal{F}(M_P^2 x) \left[1 + a \frac{2p_\gamma \cdot (p_{e^+} + p_{e^-})}{M_P^2} \right]$$

- for η' : such an expansion **not applicable** anymore
 \hookrightarrow BS necessarily depends on the form-factor model

sensitivity to width of ρ meson \hookrightarrow recent **dispersive** calculations used
Hanhart et al., EPJC 73 (2013), EPJC 77 (2017)

Källén–Lehmann spectral representation \rightarrow common spectral density function

$$\frac{\mathcal{F}(q^2)}{\mathcal{F}(0)} \simeq 1 + q^2 \int_{4m_\pi^2}^{\Lambda^2} \frac{\mathcal{A}(s) \, ds}{q^2 - s + i\epsilon}$$


$$\mathcal{A}(s) = w_\omega \mathcal{A}_\omega(s) + w_\phi \mathcal{A}_\phi(s) - \frac{\kappa}{96\pi^2 F_\pi^2} \left[1 - \frac{4m_\pi^2}{s} \right]^{3/2} P(s) R(s) |\Omega(s)|^2$$

Radiative corrections to $\eta' \rightarrow e^+ e^- \gamma$ decays

The overall NLO correction $\delta(x, 0)$ in comparison to its constituents

Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results
- Summary




Radiative corrections to $\eta' \rightarrow e^+ e^- \gamma$ decays

The two-fold differential decay width $d\Gamma(x, 0)$ at NLO

Contents

- Introduction
- π^0 rare decay
- Introduction
- Radiative corrections
- Virtual corrections
- Bremsstrahlung
- Results
- π^0 Dalitz decay
- Introduction
- Radiative corrections
- Results
- $\eta^{(\prime)}$ Dalitz decays
- Introduction
- Radiative corrections
- Specific contributions
- Results
- Summary



Summary

Contents

Introduction

 π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

 π^0 Dalitz decay

Introduction

Radiative corrections

Results

 $\eta^{(\prime)}$ Dalitz decays

Introduction

Radiative corrections

Specific contributions

Results

Summary

All NLO QED radiative corrections for discussed processes are now available
→ can be taken into account in future experimental analyses

- $\pi^0 \rightarrow e^+ e^-$

Vaško and Novotný, JHEP 1110 (2011)

TH, Kampf and Novotný, EPJC 74 (2014)

- $\pi^0 \rightarrow e^+ e^- \gamma$

TH, Kampf and Novotný, PRD 92 (2015)

→ precise knowledge of R

- $\eta^{(\prime)} \rightarrow \ell^+ \ell^- \gamma$

TH, Kampf, Leupold and Novotný, PRD 97 (2018)

Ancillary files available together with the papers

Goodbye

Contents

Introduction

 π^0 rare decay

Introduction

Radiative corrections

Virtual corrections

Bremsstrahlung

Results

 π^0 Dalitz decay

Introduction

Radiative corrections

Results

 $\eta^{(\prime)}$ Dalitz decays


Introduction

Radiative corrections

Specific contributions

Results

Summary



Thank you for listening!