



Contribution ID: 163

Type: Poster

How to find the glueball among the f_0 s with the QCD counting rules

Friday, 3 August 2018 18:04 (1 minute)

We propose a model-independent method to ascertain the leading valence composition of a hadron: to measure the energy dependence of its production cross section at a fixed angle interval. This E-dependence, by the QCD Brodsky-Farrar counting rules, falls at high energy with a steepness that depends on the leading quark and gluon composition.

We exemplify with a reaction that could help classify the f_0 mesons, exclusive $e^+e^- \rightarrow \phi + f_0$ with an easily reconstructible final state. Some of the f_0 may have a glueball gg component in their wavefunction decomposition; this will dominate at high energy over higher twist quark-antiquark components (because they necessarily have a p-wave) or hybrid/tetraquark components (because of the higher number of particles in the final state). We discuss the prospects to carry out this or similar analysis in Belle II.

Primary authors: LLANES-ESTRADA, Felipe J. (Univ. Complutense de Madrid); BRODSKY, Stanley J. (Stanford Linear Accelerator Center); PELAEZ, Jose R. (Univ. Complutense de Madrid)

Presenter: LLANES-ESTRADA, Felipe J. (Univ. Complutense de Madrid)

Session Classification: Poster

Track Classification: B: Light quarks