

Precision Measurement of β -energy Spectrum in ${}^6\text{He}$ Decay

Precision Measurement of β -energy Spectrum in ${}^6\text{He}$ Decay

- Constraint of $C_T + C_T'$ from neutron and nuclear β decays*:

$$C_T + C_T' \lesssim 3.9 \times 10^{-3}$$

- Constraint of $C_T + C_T'$ from LHC*:

$$C_T + C_T' \lesssim 2.9 \times 10^{-3}$$

Precision Measurement of β -energy Spectrum in ${}^6\text{He}$ Decay

- Constraint of $C_T + C_T'$ from neutron and nuclear β decays*:

$$C_T + C_T' \lesssim 3.9 \times 10^{-3}$$

- Constraint of $C_T + C_T'$ from LHC*:

$$C_T + C_T' \lesssim 2.9 \times 10^{-3}$$

This experiment focuses on a precision measurement of the β -energy spectrum in ${}^6\text{He}$ decay to search for tensor type contributions to the weak interactions.

Precision Measurement of β -energy Spectrum in ${}^6\text{He}$ Decay

- Constraint of $C_T + C_T'$ from neutron and nuclear β decays*:

$$C_T + C_T' \lesssim 3.9 \times 10^{-3}$$

- Constraint of $C_T + C_T'$ from LHC*:

$$C_T + C_T' \lesssim 2.9 \times 10^{-3}$$

This experiment focuses on a precision measurement of the β -energy spectrum in ${}^6\text{He}$ decay to search for tensor type contributions to the weak interactions.

Typical Fit to the Background Subtracted β -energy Spectrum Measured with the CsI(Na) Detector

