



Contribution ID: 177

Type: Talk

Yevgeny Stadnik (Johannes Gutenberg University of Mainz): New Laboratory and Astrophysical Probes for Low-Mass Dark Matter and Dark Bosons

Thursday, 22 February 2018 17:02 (14 minutes)

Low-mass bosonic dark matter particles produced after the Big Bang may form an oscillating classical field, which can be sought for in a variety of low-energy laboratory experiments based on spectroscopic, interferometric and magnetometric techniques, as well as in various astrophysical phenomena. Dark bosons can also mediate anomalous fifth forces between ordinary-matter particles that can be sought for in laboratory experiments. Recent measurements in atoms and astrophysical phenomena have already allowed us to improve on existing constraints on various non-gravitational interactions between dark bosons and ordinary-matter particles by many orders of magnitude.

References:

[1] Y. V. Stadnik and V. V. Flambaum, Phys. Rev. D 89, 043522 (2014).

[2] B. M. Roberts, Y. V. Stadnik, V. A. Dzuba, V. V. Flambaum, N. Leefer and D. Budker, Phys. Rev. Lett. 113, 081601 (2014); Phys. Rev. D 90, 096005 (2014).

[3] Y. V. Stadnik and V. V. Flambaum, Phys. Rev. Lett. 113, 151301 (2014).

[4] Y. V. Stadnik and V. V. Flambaum, Phys. Rev. Lett. 114, 161301 (2015); Phys. Rev. A 93, 063630 (2016).

[5] Y. V. Stadnik and V. V. Flambaum, Phys. Rev. Lett. 115, 201301 (2015); Phys. Rev. A 94, 022111 (2016).

[6] N. Leefer, A. Gerhardus, D. Budker, V. V. Flambaum and Y. V. Stadnik, Phys. Rev. Lett. 117, 271601 (2016).

[7] Y. V. Stadnik, V. A. Dzuba and V. V. Flambaum, arXiv:1708.00486.

[8] C. Abel et al. (nEDM collaboration), arXiv:1708.06367; Phys. Rev. X (In press).

[9] V. A. Dzuba, V. V. Flambaum, and Y. V. Stadnik, arXiv:1709.10009; Phys. Rev. Lett. (In press).

Presenter: STADNIK, Yevgeny (University of New South Wales)

Session Classification: Session 9