



Contribution ID: 124

Type: not specified

Exclusive Photoproduction of $2\pi^+2\pi^-$ Final State at HERA

Tuesday, April 17, 2018 9:00 AM (24 minutes)

Exclusive production of four charged pions at the ep collider HERA is studied at small photon virtualities $Q^2 < 2 \text{ GeV}^2$. The data were taken with the H1 detector in the years 2006 and 2007 at a centre-of-mass energy of $\sqrt{s} = 319 \text{ GeV}$ and correspond to an integrated luminosity of 7.6 pb^{-1} . The cross section of the reaction $\gamma p \rightarrow 2(\pi^+\pi^-)Y$ is determined in the phase space of $45W_g p 100 \text{ GeV}$, $|t| 1 \text{ GeV}^2$ and $M_Y 1.6 \text{ GeV}$. The 4π mass spectra indicate that the reaction proceeds predominately via production and decay of $\rho'(1450)$ and $\rho''(1700)$ resonances. Parameters of these resonances as well as production cross sections times branching ratio into four charged pions are estimated from the mass fit, which includes contributions from non-resonant 4π channel and interference terms.

Primary authors: SCHMITT, Stefan (Deutsches Elektronen-Synchrotron (DE)); LEVONIAN, Sergey (Deutsches Elektronen-Synchrotron (DE)); COLLABORATION, H1 (DESY)

Presenter: SCHMITT, Stefan (Deutsches Elektronen-Synchrotron (DE))

Session Classification: WG2: Small-x and Diffraction

Track Classification: WG2: Small-x and Diffraction