



Contribution ID: 186

Type: not specified

Recent QCD-related studies with the BaBar detector

Thursday 19 April 2018 09:25 (25 minutes)

We report some of the most recent results in studying different aspects of QCD with about 500 fb^{-1} of data collected by the BaBar experiment at the e^+e^- B -factory PEP-II.

Among these, we present a high precision measurement of the mass difference between the $D^*(2010)^+$ and D^+ mesons using the decay chain $D^*(2010)^+ \rightarrow D^+\pi^0$, with $D^+ \rightarrow K^-\pi^+\pi^+$. This result is then combined with a previous BaBar measurement of $m(D^*(2010)^+) - m(D^0)$ to extract the mass difference between the charged and neutral D mesons. We obtain results that are approximately seven times more precise than the present world averages.

We also report on the first evidence for a B -meson decay to four baryons, $B^0 \rightarrow p\bar{p}p\bar{p}$, which can help to shed light on the experimental discrepancy between the inclusive branching fraction of all B meson decay modes with at least a couple of baryons in the final state, measured by ARGUS to be $(6.8 \pm 0.6)\%$, and the sum of exclusive baryonic channels. Finally, we present a measurement of the spectral function for the $\tau^- \rightarrow K^- K_S \nu_\tau$ decay, which can be used to determine the hadronic contribution to the muon $g - 2$ due to the vacuum polarization,

Authors: ANULLI, Fabio (Sapienza Universita e INFN, Roma I (IT)); ZANI, Laura (INFN - National Institute for Nuclear Physics)

Presenter: ZANI, Laura (INFN - National Institute for Nuclear Physics)

Session Classification: WG4-WG5 Joint Session

Track Classification: WG5: Physics with Heavy Flavours