## XXVI International Workshop on Deep Inelastic Scattering and Related Subjects



Contribution ID: 223

Type: not specified

## Deep inelastic scattering in the dipole picture at next-to-leading order

Tuesday 17 April 2018 09:24 (24 minutes)

We study quantitatively [1] the importance of the recently derived NLO corrections [2,3,4] to the DIS structure functions at small x in the dipole formalism. We show that these corrections can be significant and depend on the factorization scheme used to resum large logarithms of energy into renormalization group evolution with the BK equation. This feature is similar to what has recently been observed for single inclusive forward hadron production [5,6]. Using a factorization scheme consistent with the one recently proposed for the single inclusive cross section, we show that it is possible to obtain meaningful results for the DIS cross sections. We also discuss ongoing work to combine these NLO DIS structure functions in the improved factorization scheme with the resummed/NLO BK evolution equation in order to do consistent NLO accuracy comparisons with HERA data.

- [1] B. Ducloué, H. Hänninen, T. Lappi, Y. Zhu, Phys.Rev. D96 (2017) no.9, 094017
- [2] G. Beuf, Phys.Rev. D94 (2016) no.5, 054016
- [3] G. Beuf, Phys.Rev. D96 (2017) no.7, 074033
- [4] H. Hänninen, T. Lappi, R. Paatelainen, arXiv:1711.08207 [hep-ph]
- [5] E. Iancu, A.H. Mueller, D.N. Triantafyllopoulos, JHEP 1612 (2016) 041
- [6] B. Ducloué, T. Lappi, Y. Zhu, Phys.Rev. D95 (2017) no.11, 114007

Primary author: HÄNNINEN, Henri (University of Jyväskylä) Presenter: HÄNNINEN, Henri (University of Jyväskylä)

Session Classification: WG2: Small-x and Diffraction

Track Classification: WG2: Small-x and Diffraction