



Contribution ID: 553

Type: Poster

On the differences among Initial Conditions and their role in the distribution of particles

Tuesday 15 May 2018 19:10 (30 minutes)

There are several theoretical models for initial conditions, which intend to describe the matter created in ultra-relativistic heavy-ion collisions. In this work, we will perform hydrodynamics computations using different initial conditions, as Wood-Saxon, Glauber, Mckln, Nexus and Gubser [1], and calculate several observables, as for instance, anisotropic flows v_n [2], eccentricities ε_n , symmetric cumulants $SC(n, m)$ [3], event-plane correlations, and others quantities, in order to understand the hydrodynamics response to different initials geometry in heavy-ion collisions.

[1] Different references, but you can see “The Fate of the Initial State Fluctuations in Heavy Ion Collisions. II The Fluctuations and Sounds”. Pilar Staig, Edward Shuryak. Phys.Rev. C84 (2011) 034908

[2] “Anisotropic flow in event-by-event ideal hydrodynamic simulations of 200 GeV Au+Au collisions.” Fernando G. Gardim, Frederique Grassi, Matthew Luzum, Jean-Yves Ollitrault. Phys.Rev.Lett. 109 (2012) 202302

[3] “Hydrodynamic Predictions for Mixed Harmonic Correlations in 200 GeV Au+Au Collisions” Fernando G. Gardim, Frederique Grassi, Matthew Luzum, Jacquelyn Noronha-Hostler. Phys.Rev. C95 (2017) no.3, 034901

“This work is in progress, and will be finished up to march 2018.

Content type

Theory

Collaboration

Centralised submission by Collaboration

Presenter name already specified

Primary authors: Dr GARDIM, Fernando (Federal University of Alfenas); Mr MAGALHÃES, Pablo (UNIFAL)

Presenter: Dr GARDIM, Fernando (Federal University of Alfenas)

Session Classification: Poster Session

Track Classification: Initial state physics and approach to equilibrium