



Contribution ID: 244

Type: **Poster**

Measurement of the underlying event in the presence of high pileup at ATLAS

Tuesday 15 May 2018 19:10 (30 minutes)

In measurements of bulk charged-particle production, the kinematic characteristics of produced hadrons and correlation between them in the presence of the hard scattering signatures is a very interesting topic for understanding the physics processes relevant in small collision systems. Extensive data samples have been accumulated by the LHC experiments to conduct these studies in pp collisions. These measurements, however, are complicated by the presence of many pileup vertices in each bunch crossing which contaminate direct studies of the characteristics of the underlying event. This problem is addressed in the ATLAS detector using an event mixing technique which allows to correct for the presence of the pileup on a statistical basis. The implementation, result and limitations of the techniques are presented in this poster. The results use $\sqrt{s} = 8$ and 13 TeV data samples with a combined integrated luminosity exceeding 50 fb^{-1} .

Content type

Experiment

Collaboration

ATLAS

Centralised submission by Collaboration

Presenter name already specified

Authors: WOSIEK, Barbara Krystyna (Institute of Nuclear Physics Polish Academy of Sciences (PL)); MILOV, Alexander (Weizmann Institute of Science (IL))

Presenter: MILOV, Alexander (Weizmann Institute of Science (IL))

Session Classification: Poster Session

Track Classification: Collectivity in small systems