



Contribution ID: 189

Type: Poster

First results on charged $K^*(892)$ resonance production in pp collisions at $\sqrt{s} = 13$ with ALICE at the LHC

Tuesday, May 15, 2018 7:10 PM (30 minutes)

The study of strange hadronic resonances in pp collisions contributes to the study of strangeness production in small systems. Measurements in pp collisions constitute a reference for the study in larger colliding systems and provide constraints for tuning QCD-inspired event generators. Since the lifetimes of short-lived resonances such as $K^*(892)^\pm$ ($\tau \sim 4$ fm/c) are comparable with the lifetime of the fireball produced in heavy-ion collisions, regeneration and rescattering effects can modify the measured yield, especially at low transverse momentum.

The first results for the $K^*(892)^\pm$ resonance obtained in inelastic pp collisions at $\sqrt{s} = 13$ TeV will be shown. The $K^*(892)^\pm$ has been measured at mid-rapidity via its hadronic decay channel $K^*(892)^\pm \rightarrow K_S^0 + \pi^\pm$, with the ALICE detector. In particular, the transverse momentum (p_T) spectrum, integrated yields, $\langle p_T \rangle$ and ratio to stable hadrons will be presented. The $K^*(892)^\pm$ results are compared with $K^*(892)^0$ measurements and with commonly-used Monte Carlo models. Measurements at 13 TeV are in addition a baseline for comparison with pp measurements at other LHC energies.

Content type

Experiment

Collaboration

ALICE

Centralised submission by Collaboration

Presenter name already specified

Primary author: GARG, Kunal (Universita e INFN, Catania (IT))

Co-author: COLLABORATION, ALICE

Presenter: GARG, Kunal (Universita e INFN, Catania (IT))

Session Classification: Poster Session

Track Classification: Thermodynamics and hadron chemistry