

# Parity Doubling of Baryons in QCD Thermodynamics

Chihiro Sasaki

Institute of Theoretical Physics

University of Wroclaw


in collaboration with

Kenji Morita (RIKEN & Wroclaw),

Pok Man Lo, Krzysztof Redlich (Wroclaw)

# Spectra in a chirally restored world

- Lowest scalar meson  $\rightarrow$  O(4) vector with pion
- Parity partners degenerate  $\rightarrow$  chiral partners
- QCD ground-state particles: pions & nucleons





# Lattice QCD tells us ...

❑ Spatial correlations [DeTar-Kogut, 1987]

❑ Temporal correlations [FASTSUM Coll., 2015-17:

$m_{\pi} \approx 400$  MeV,  $m_k \approx 500$  MeV, Wilson fermions,  $T_{ch} = 185$  MeV]



vs.  $M_n \approx 3 \times M_q$

# Non-SCB mass of nucleons


□ SU(2) chiral transformation of 2 nucleons

→ how to assign 2 indep. rotation to them?

$$\psi_{1L} \rightarrow g_l \psi_{1L}, \quad \psi_{1R} \rightarrow g_r \psi_{1R} \sim \psi_{1L} : (1/2, 0) \quad \psi_{1R} : (0, 1/2)$$

$$\psi_{2L} \rightarrow g_r \psi_{2L}, \quad \psi_{2R} \rightarrow g_l \psi_{2R} \sim \psi_{2L} : (0, 1/2) \quad \psi_{2R} : (1/2, 0)$$

$$\mathcal{L}_m = m_0 (\bar{\psi}_2 \gamma_5 \psi_1 - \bar{\psi}_1 \gamma_5 \psi_2) \Rightarrow m_{N_{\pm}} = \frac{1}{2} \left[ \sqrt{c_1 \sigma^2 + 4m_0^2} \mp c_2 \sigma \right]$$



[DeTar-Kunihiro, 1989]

Red: Standard  
Blue: Mirror

# Origin of the survival mass?

□ Emergence of a scale in QCD → trace anomaly

$$\partial_\mu J^\mu = T_\mu^\mu \propto \langle H | G^2 | H \rangle$$

- in hot matter: reduced by 50% at  $T_c$  [Miller, 2007: lattice EoS]
- in nuclear matter: reduced by 5% at normal  $\rho$  [Cohen et al. 1995: Feynman-Hellmann theorem & low-density approx.]

□ How large is  $m_0$ ? --- not conclusive!

- Models: 300-800 MeV
- Lattice (FASTSUM): 800-900 MeV

✓  $m_0 \approx$  a few  $\Lambda_{\text{qcd}}$ , mass diff.  $\approx$  weaker  $m_0$  dep.

# Parity doubling of baryons

- Baryon octet and decuplet with finite  $m_0$
- Consistent with established phenomenology:
  - ✓ Gell-Mann-Okubo mass formula

$$\frac{3}{4}m_\Lambda + \frac{1}{4}m_\Sigma - \frac{1}{2}(m_N + m_\Xi) = 0$$

- ✓ Gell-Mann's equal spacing rule

$$m_{\Sigma^*} - m_\Delta = m_{\Xi^*} - m_{\Sigma^*} = m_\Omega - m_{\Xi^*}$$

- Mass relations [CS, 2017]

$$M_B(\sigma_q, \sigma_s; a, b, m_0)$$

Light-quark condensate    Strange-quark condensate

# Chiral model vs. LQCD (FASTSUM)


❑ Strong  $m_{\pi}$  dependence; SU(2+1) vs. SU(3)

❑ Fitting the LQCD masses [Aarts et al., 2017]

$$M^-(T) = M^-(T=0)\omega(T, b) + M^-(T_c)(1 - \omega(T, b))$$

$$\omega(T, b) = \tanh[(1 - T/T_c)/b] / \tanh(1/b)$$

$$M_-^{\text{lat}}(T_c) \times \frac{M_+^{\text{PDG}}}{M_+^{\text{lat}}}$$



**Any imprint in EoS?**



# Signal of chiral symmetry restoration

- ❑ Lattice QCD shows clearly  $\langle qq\bar{q} \rangle$  dropping!
- ❑ More deviation from HRG in higher-order fluctuations  $\rightarrow$  Missing states? Interactions? and/or in-medium effects?
- ❑ In-medium HRG [Aarts et al., 2017]
  - T-dep. masses motivated by Lattice findings
  - Constant masses for positive-parity states
  - Its verification?  $\rightarrow$  baryon number fluctuations.

corrected  $M_{-}^{\text{lat}}(T_c)$


uncorrected

# Fluctuations of net-baryon number



$$\chi_{ijk}^{BQS} \equiv$$

$$\frac{\partial^{i+j+k}}{\partial^i(\mu_B/T) \partial^j(\mu_Q/T) \partial^k(\mu_S/T)} p(T, \mu_B, \mu_Q, \mu_S) / T^4$$



Accidental!

[Morita et al., arXiv:1711.10779 [hep-ph]]

# What is missing? --- finite width

□ Thermodynamics of broad resonances

→ S matrix approach [Dashen, Ma and Bernstein, 1969]

- Grand canonical potential

$$\Omega = \Omega_0 + \Omega_{\text{int}}$$

$$\Delta \ln Z = \int dE e^{-\beta E} \frac{1}{4\pi i} \text{tr} \left[ S^{-1} \overleftrightarrow{\frac{\partial}{\partial E}} S \right]_c$$

- Leading contribution: 2-body [Beth-Uhlenbeck, 1937]

$$\Delta \ln Z = \int dE e^{-\beta E} \times \frac{1}{\pi} \frac{\partial}{\partial E} \text{tr} (\delta_E) \text{Phase shift}$$

Dynamical information

# What is missing? --- finite width

□  $K_0^*/\kappa$  (800) meson: chiral partner of kaon

NOTE: omitted from PDG summary table


□ S matrix approach [Friman et al. 2015]

✓ Empirical  $\pi$ -K phase shift from experiment


$$\Omega = \Omega_\pi + \Omega_K + \Omega_{\text{int}}$$

cf. HRG

$$\Omega = \Omega_\pi + \Omega_K + \Omega_{\text{res}}$$



# Pi-Nucleon system [Lo et al., 2018]



# Parity doublers in cold dense QCD

- Quark-nucleon hybrid model [Benic et al., 2015]
  - How to suppress quarks at low density?
    - ✓ IR/UV cutoff “b” in Fermi dist. functions
    - ✓ from const. “b” to a VEV of a scalar field b
  - Chiral & deconfinement p.t. in a single framework
- Symmetric matter [Marczenko, CS, 2017]
  - Net-baryon numb. fluct.:  $\chi$ -dynamics dominated
- Asymmetric matter [Marczenko et al., to appear]
  - Constraints on the mass and compactness of a star → hadronic scenario w/o deconf. quarks

# Summary

□ Emergent parity-doubling structure as a manifestation of restored chiral symmetry

Lessons:

- ◆ Naive “in-medium HRG” does not work.
- ◆ Effect of resonance widths – beyond HRG
- ◆ Survival mass  $\approx$  chromo-magnetic sector
  
- ◆ Higher-lying states near QCD p.t.
- ◆ Interplay between CSB and confinement
- ◆ Toward more realistic description of QCD