



Contribution ID: 170

Type: Poster

## Effects of resonance widths on particle distributions and anisotropies in heavy-ion collisions

*Tuesday, May 15, 2018 7:10 PM (30 minutes)*

We extend the S-matrix framework to the Delta-type resonances (spin 3/2, isospin 3/2) in elastic pion-nucleon scatterings up to 1.8 GeV mass. We evaluate not only Deltas, but also rho, f<sub>0</sub>, K\* and K<sub>0</sub> meson properties using the S-matrix framework, and implement them in the hydrodynamical description of Pb+Pb collisions at LHC.

We show that the proper treatment of resonances modifies the spectrum of daughter particles, and thus the final observable distributions. In particular the yield of pions at low p<sub>T</sub> increases, which reduces the average transverse momentum, and thus improves the description of the pion spectrum measured in the heavy-ion experiments.

### Content type

Theory

### Collaboration

### Centralised submission by Collaboration

Presenter name already specified

**Primary authors:** HUOVINEN, Pasi (University of Wroclaw); Dr LO, Pok Man (University of Wroclaw); NIEMI, Harri (Johann Wolfgang Goethe-Universität); REDLICH, Krzysztof (University of Wroclaw)

**Presenter:** REDLICH, Krzysztof (University of Wroclaw)

**Session Classification:** Poster Session

**Track Classification:** Collective dynamics