

# Tests of Universality of Baryon Form Factors In Holographic QCD



A. Cherman, T.D. Cohen, M. Nielsen, PRL103(09)

Univ. of Maryland and Univ. of Sao Paulo




Low x Meeting,  
September 2009

# QCD: strong-coupled theory



# AdS/QCD models: successful at reproducing low energy hadronic observables

- two classes of AdS/QCD models




## Top-down models:

arise from string theory

D4/D8 system  gauge theory confining as QCD

AdS/CFT requires large  $N_c$  and 't Hooft coupling

## Bottom-up models:

QCD large  $N_c$   dual to a classical 5D theory

field content 5D matched

to low energy chiral symmetry of QCD

# Both cases: large $N_c$ is required

- QCD is a weakly-interacting theory of long-lived mesons in the large  $N_c$  limit.
- In the large  $N_c$  limit, baryons are 'soliton-like' configurations of meson fields.
- Baryon masses scale as  $N_c^1$  (composed of  $N_c$  quarks), interactions with mesons scale as  $N_c^{1/2}$ .
  - Unlike mesons, baryons are not narrow at large  $N_c$ .
  - Nucleons, deltas, become degenerate. Mass splitting scales as  $N_c^{-1}$ .
- In contrast to the pure meson sector, meson loops make leading order contributions to baryon properties at large  $N_c$ .
- Variety of baryon models (Skyrme models) use large  $N_c$  properties of baryons for inspiration.
- Baryons modeled as quantum states of slowly rotating hedgehog Skyrmons of meson fields.

't Hooft, 1973

Witten, 1979

Skyrme, 1961

Adkins, Nappi,  
Witten, 1983

Dashen, Manohar,  
Jenkins, 1993-95

# Model-independent relations for baryons valid at large $N_c$

- Goldberger-Treiman relation:  $m_N g_A = f_\pi g_{\pi NN}$
- Relation is model-independent, and follows from chiral symmetry.
- Nucleon and delta couplings with pions are related by

$$2g_{\pi NN} = 3g_{\pi N\Delta}$$


- This relation follows just from the large  $N_c$  limit
- New model-independent relation:
- Ratio of nucleon form factors in position space, evaluated at large distances:  
$$\lim_{r \rightarrow \infty} \frac{G_E^{I=0}(r) G_E^{I=1}(r)}{G_M^{I=0}(r) G_M^{I=1}(r)} = \frac{18}{r^2}$$
- Relation depends on **both** the large  $N_c$  and chiral limits.
- Can serve as a useful and highly non-trivial probe of large  $N_c$  baryon models.

# Model independence of the ratio

Leading large  $r$  behavior of form factors in large  $N_c$  ChiPT are mediated by pions

$$G_E^{I=0}(r)$$

$$G_M^{I=0}(r)$$




Three pion interaction - comes from anomaly.

$$\sim \frac{1}{f_\pi^3} \left( \frac{g_A}{f_\pi} \right)^3$$

$$G_E^{I=1}(r)$$

$$G_M^{I=1}(r)$$



Two pion interaction.

$$\sim \left( \frac{g_A}{f_\pi} \right)^2$$

- In the large  $N_c$  limit, leading contributions to isovector electric and isoscalar magnetic form factors are sensitive to the nucleon-delta mass splitting  $\sim 1/N_c$  (Cohen PLB 359)

$$G_E^{I=1} \propto \Delta \left( \frac{g_A}{f_\pi} \right)^2 \qquad G_M^{I=0} \propto \frac{\Delta}{f_\pi^3} \left( \frac{g_A}{f_\pi} \right)^3$$

$$G_M^{I=1} \propto \left( \frac{g_A}{f_\pi} \right)^2 \qquad G_E^{I=0} \propto \frac{1}{f_\pi^3} \left( \frac{g_A}{f_\pi} \right)^3$$

$$\lim_{r \rightarrow \infty} \frac{G_E^{I=0}(r) G_E^{I=1}(r)}{G_M^{I=0}(r) G_M^{I=1}(r)}$$

ratio is independent  
of

$$g_A \quad f_\pi \quad \Delta$$

# Skyrme-type models

$$\mathbb{L} = \frac{1}{26} f_\pi^2 \text{Tr} [\partial_\mu U \partial^\mu U^\dagger] + \frac{1}{2^5 e^2} \text{Tr} [(\partial_\mu U) U^\dagger, (\partial_\nu U) U^\dagger]^2$$

$$U = e^{\frac{i \vec{\pi} \cdot \vec{\tau}}{f_\pi}} \quad \rightarrow \quad \text{SU(2) matrix}$$

- Original Skyrme model includes only pion fields.
- Baryons appear as quantum states of slowly rotating hedgehog Skyrmons of pion fields.

hedgehog ansatz:  $\vec{\pi} = f(r) \vec{\tau}$

in the large  $r$  limite  $f(r)$  goes like  $\eta/r^2$



in the Skyrme model:

$$g_A = \frac{8\pi}{3} f_\pi^2 \eta$$

$$\lim_{r \rightarrow \infty} G_{I=0}^E = \frac{3^3}{2^9 \pi^5} \frac{1}{f_\pi^3} \left( \frac{g_A}{f_\pi} \right)^3 \frac{1}{r^9}$$

$$\lim_{r \rightarrow \infty} G_{I=0}^M = \frac{3\Delta}{2^9 \pi^5} \frac{1}{f_\pi^3} \left( \frac{g_A}{f_\pi} \right)^3 \frac{1}{r^7}$$

$$\lim_{r \rightarrow \infty} G_{I=1}^M = \frac{1}{2^5 \pi^2} \left( \frac{g_A}{f_\pi} \right)^2 \frac{1}{r^4}$$

$$\lim_{r \rightarrow \infty} G_{I=1}^E = \frac{\Delta}{2^4 \pi^2} \left( \frac{g_A}{f_\pi} \right)^2 \frac{1}{r^4} .$$

$$\lim_{r \rightarrow \infty} \frac{G_E^{I=0}(r) G_E^{I=1}(r)}{G_M^{I=0}(r) G_M^{I=1}(r)} = \frac{18}{r^2}$$

# 5D Skyrmons

## Pomarol-Wulzer holographic baryon model

Pomarol, Wulzer, NPB809:347-361,2009.

Panico, Wulzer, arXiv:0811.2211

- Bottom-up AdS/QCD model: uses hard-wall AdS background, with two  $U(2)$  5D gauge fields,  $L_M$   $R_M$ , associated with 4D left and right quark currents ( AdS/CFT dictionary).
- Chiral symmetry broken by choice of IR boundary conditions for 5D gauge fields.
- Has a 5D CS term to get anomaly physics right.

$$S = -\frac{M_5}{2} \int d^5x \sqrt{g} \operatorname{Tr} [\mathbf{L}_{MN}^2 + \mathbf{R}_{MN}^2] + \frac{-iN_c}{24\pi^2} \int_{5D} [\omega_5(\mathbf{L}) - \omega_5(\mathbf{R})]$$

$$M_5 \sim \mathcal{O}(N_c^1)$$

- this model looks *quite* different from 4D Skyrme models: there are no explicit pion fields.
- Baryons appear as quantum states of slowly rotating Skyrmion-like hedgehog configurations of the 5D gauge field.
- Skyrmons are stabilized by the CS term.

|                                    | Experiment | AdS <sub>5</sub> | Deviation |
|------------------------------------|------------|------------------|-----------|
| $M_N$                              | 940 MeV    | 1130 MeV         | 20%       |
| $\mu_S$                            | 0.44       | 0.34             | 30%       |
| $\mu_V$                            | 2.35       | 1.79             | 31%       |
| $g_A$                              | 1.25       | 0.70             | 79%       |
| $\sqrt{\langle r_{E,S}^2 \rangle}$ | 0.79 fm    | 0.88 fm          | 11%       |
| $\sqrt{\langle r_{E,V}^2 \rangle}$ | 0.93 fm    | $\infty$         |           |
| $\sqrt{\langle r_{M,S}^2 \rangle}$ | 0.82 fm    | 0.92 fm          | 12%       |
| $\sqrt{\langle r_{M,V}^2 \rangle}$ | 0.87 fm    | $\infty$         |           |
| $\sqrt{\langle r_A^2 \rangle}$     | 0.68 fm    | 0.76 fm          | 12%       |
| $\mu_p / \mu_n$                    | -1.461     | -1.459           | 0.1%      |

$g_A = 0.65$  in the original Skyrme model

in the large  $r$  limit

$$\lim_{r \rightarrow \infty} G_E^{I=0} = -\frac{\beta^3 L^6}{\pi^2} \frac{1}{r^9}$$


$$\lim_{r \rightarrow \infty} G_M^{I=0} = \frac{\beta^3 L^6}{6\pi^2 \mathcal{I}} \frac{1}{r^7}$$

$$\lim_{r \rightarrow \infty} G_E^{I=1} = \frac{8\beta^2}{3\mathcal{I}} M_5 L^3 \frac{1}{r^4}$$

$$\lim_{r \rightarrow \infty} G_M^{I=0} = -\frac{8\beta^2}{9} M_5 L^3 \frac{1}{r^4}$$

$$\lim_{r \rightarrow \infty} \frac{G_E^{I=0}(r) G_E^{I=1}(r)}{G_M^{I=0}(r) G_M^{I=1}(r)} = \frac{18}{r^2}$$

large  $N_c$  and chiral physics handled correctly in  
Pomarol-Wulzer model



# Baryons as Holographic Instantons

## Sakai-Sugimoto model

Hata et al, hep-th/0701280

Hashimoto, Sakai, Sugimoto, PTP120:1093-1137,2008

- Top-down AdS/QCD model: baryons are described as instantons in a 5D Yang-Mills and CS theory, formulated in the D4/D8 model.

$$S = -\kappa \int d^4x dz \left( \text{Tr} \frac{\mathbf{F}_{\mu\nu}^2}{2} (1+z^2)^{-1/3} + (1+z^2) \mathbf{F}_{\mu z}^2 \right) + \frac{N_c}{24\pi^2} \int \omega_5(\mathbf{A})$$

- $A_M$  :5D  $U(N_f)$  field,  $F_{MN}$ : field strength,  $\omega_5(A)$  : CS 5-form
- Model expected to make sense when  $N_c$  and the 't Hooft coupling are large.

- CS term stabilizes the instanton size to be of order  $\lambda^{-1/2}$
- At large  $\lambda$  higher derivative terms are not suppressed, and  $1/\lambda$  expansion is not well justified

|                                      | SS model               | Skyrmion <sup>14)</sup> | experiment             |
|--------------------------------------|------------------------|-------------------------|------------------------|
| $\langle r^2 \rangle_{I=0}^{1/2}$    | 0.742 fm               | 0.59 fm                 | 0.806 fm               |
| $\langle r^2 \rangle_{M, I=0}^{1/2}$ | 0.742 fm               | 0.92 fm                 | 0.814 fm               |
| $\langle r^2 \rangle_{E,p}$          | $(0.742 \text{ fm})^2$ | $\infty$                | $(0.875 \text{ fm})^2$ |
| $\langle r^2 \rangle_{E,n}$          | 0                      | $-\infty$               | $-0.116 \text{ fm}^2$  |
| $\langle r^2 \rangle_{M,p}$          | $(0.742 \text{ fm})^2$ | $\infty$                | $(0.855 \text{ fm})^2$ |
| $\langle r^2 \rangle_{M,n}$          | $(0.742 \text{ fm})^2$ | $\infty$                | $(0.873 \text{ fm})^2$ |
| $\langle r^2 \rangle_A^{1/2}$        | 0.537 fm               | —                       | 0.674 fm               |
| $\mu_p$                              | 2.18                   | 1.87                    | 2.79                   |
| $\mu_n$                              | -1.34                  | -1.31                   | -1.91                  |
| $\left  \frac{\mu_p}{\mu_n} \right $ | 1.63                   | 1.43                    | 1.46                   |
| $g_\Lambda$                          | 0.734                  | 0.61                    | 1.27                   |
| $g_{\pi NN}$                         | 7.46                   | 8.9                     | 13.2                   |
| $g_{\rho NN}$                        | 5.80                   | —                       | 4.2 ~ 6.5              |



in the large  $r$  limit

$$\lim_{r \rightarrow \infty} G_E^{I=0}(r) = \frac{g_{v^1} \psi_1(0)}{4\pi r} e^{-\rho_1 r}$$

$$\lim_{r \rightarrow \infty} G_M^{I=0}(r) = \frac{9\pi r}{16\pi \lambda N_c} g_{v^1} \psi_1(0) \rho_1 e^{-\rho_1 r}$$

$$\lim_{r \rightarrow \infty} G_E^{I=1}(r) = \frac{g_{v^1} \psi_1(0)}{4\pi r} e^{-\rho_1 r}$$

$$\lim_{r \rightarrow \infty} G_M^{I=1}(r) = \frac{N_c}{12\pi} \sqrt{\frac{2}{15}} g_{v^1} \psi_1(0) \rho_1 e^{-\rho_1 r}$$

$$\lim_{r \rightarrow \infty} \frac{\tilde{G}_E^{I=0} \tilde{G}_E^{I=1}}{\tilde{G}_M^{I=0} \tilde{G}_M^{I=1}} = \frac{\lambda \sqrt{40/3}}{\pi \rho_1^2 r^2}$$

$\rho_1$   related with meson rho mass

- Model does not satisfy large  $N_c$  relation.
  - Ratio depends on model parameters.
- Model fails because it does not treat chiral symmetry correctly.
  - Another signal: the isovector charge radius is finite in the model. It is known to be infinite in chiral perturbation theory in the chiral limit.
- These results suggest that the Sakai-Sugimoto instanton model fails to correctly describe the long-range part of large  $N_c$  baryon physics.

# Summary

- We discussed a model-independent large  $N_c$  relation for baryons.

$$\lim_{r \rightarrow \infty} \frac{G_E^{I=0}(r) G_E^{I=1}(r)}{G_M^{I=0}(r) G_M^{I=1}(r)} = \frac{18}{r^2}$$

- Relation probes consistency of implementation of chiral and large  $N_c$  physics in baryon models.
- In this case, the probe reveals that some holographic models get large  $N_c$  chiral physics right, while others do not.
- This situation illustrates the utility of large  $N_c$  analysis as a diagnostic tool for probing models.
- Relation should be checked in other new large  $N_c$  baryon models.

<http://www.sbf1.sbfisica.org.br/eventos/extras/hadrons2010/>



## **XI HADRON PHYSICS**

**March, 22 - 27, 2010, São Sebastião, Brazil**

origem of problem: large expansion.

- SS model: pion nucleon coupling scales as

$$\frac{g_A}{f_\pi} \sim \sqrt{\frac{N_c}{\lambda}}$$

- pion loops contributions to nucleon properties are discarded if the large  $\lambda$  limit is taken prior to the large  $N_c$  limit.
- It is necessary to go beyond large  $\lambda$  limit in the SS model to capture large  $N_c$  chiral physics.

# Definitions

$$G_E^{I=0}(r) = \frac{1}{4\pi} \int d\Omega \langle p \uparrow | J_{I=0}^0 | p \uparrow \rangle,$$

$$G_M^{I=0}(r) = \frac{1}{4\pi} \int d\Omega \frac{1}{2} \varepsilon_{ij3} \langle p \uparrow | x_i J_{I=0}^j | p \uparrow \rangle,$$

$$G_E^{I=1}(r) = \frac{1}{4\pi} \int d\Omega \langle p \uparrow | J_{I=1}^{03} | p \uparrow \rangle,$$

$$G_M^{I=1}(r) = \frac{1}{4\pi} \int d\Omega \frac{1}{2} \varepsilon_{ij3} \langle p \uparrow | x_i J_{I=1}^{j3} | p \uparrow \rangle,$$

$$J_{I=1}^{\mu 3} = \text{Tr} [J_{I=1}^{\mu} \sigma^3]$$

These position-space form factors can be related to the standard (experimentally accessible) momentum-space form factors by the appropriate Fourier transforms.