



# Kinematically Dependent Renormalization



**Dmitry Kazakov**



**Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, Russia**

**Moscow Institute of Physics and Technology, Dolgoprudny, Russia**

Based on [arXiv:1804.08387](https://arxiv.org/abs/1804.08387) [hep-th]

**Lev's Lipatov Memorial Session**

**July, 7 2018 Kolymbari, Crete, Greece**

- **This is an attempt to shed some new light on non-renormalizable interactions with the aim to make sense of them at least in some cases**
- **As an example we consider maximally supersymmetric gauge theory in  $D=8$  dimensions and focus on the on-shell scattering amplitudes**
- **The reason is that this case was studied in detail**

**in collaboration with A. Borlakov, D.Tolkachev and D.Vlasenko**

**JHEP 1612 (2016) 154, arXiv:1610.05549v2 [hep-th]  
Phys.Rev. D95 (2017) no.4, 045006 arXiv:1603.05501 [hep-th]  
Phys.Rev. D97 (2018) no.12, 125008 arXiv:1712.04348 [hep-th],**

**and has important advantages**

- **All analysis is performed within dimensional regularization**

# Motivation

## Maximal SYM

**D=4 N=4**

**D=6 N=2**

**D=8 N=1**

**D=10 N=1**

- **Partial or total cancellation of UV divergences (all bubble and triangle diagrams cancel)**
- **First UV divergent diagrams at  $D=4+6/L$**
- **Conformal or dual conformal symmetry**
- **Common structure of the integrands**

*Bern, Dixon & Co 10*  
*Drummond, Henn, Korchemsky, Sokatchev 10*  
*Arkani-Hamed 12*

# Motivation

## Maximal SYM

**D=4 N=4**

**D=6 N=2**

**D=8 N=1**

**D=10 N=1**

- Partial or total cancellations (all bubble and triangle diagrams cancel)
  - First discovered in  $\mathcal{N}=4$  SYM at  $D=4+6/L$
- All of them can be obtained from 10dim superstring by compactification on a torus**
- on structure of the integrands**

Bern, Dixon & Co 10

Drummond, Henn, Korchemsky, Sokatchev 10

Arkani-Hamed 12

# Motivation

## Maximal SYM

**D=4 N=4**

**D=6 N=2**

**D=8 N=1**

**D=10 N=1**

**D=4 N=8 Supergravity**

- Partial or total cancellations (all bubble and triangle diagrams cancel)
- First UV divergence appears at  $D=4+6/L$
- On-shell finiteness is due to the special conformal symmetry
- On-shell finiteness is due to the special structure of the integrands

**All of them can be obtained from 10dim superstring by compactification on a torus**

- On-shell finite up to 8 loops
- Similar to higher dim SYM

Bern, Dixon & Co 10  
Drummond, Henn, Korchemsky, Sokatchev 10  
Arkani-Hamed 12

# Motivation

## Maximal SYM

D=4 N=4

D=6 N=2

D=8 N=1

D=10 N=1

- Partial or total cancellations (all bubble and triangle diagrams cancel)
- First discovered in 1970s at D=4+6/L
- Maximal conformal symmetry
- Hidden structure of the integrands

All of them can be obtained from 10dim superstring by compactification on a torus

Bern, Dixon & Co 10

Drummond, Henn, Korchemsky, Sokatchev 10

Arkani-Hamed 12

D=4 N=8 Supergravity

- On-shell finite up to 8 loops
- Similar to higher dim SYM

Object: Helicity Amplitudes on mass shell with arbitrary number of legs and loops

The case: Planar limit  $N_c \rightarrow \infty$ ,  $g_{YM}^2 \rightarrow 0$  and  $g_{YM}^2 N_c$  - fixed

The aim: to get all loop (exact) result

# Motivation

## Maximal SYM

D=4 N=4

D=6 N=2

D=8 N=1

D=10 N=1

- Partial or total cancellations (all bubble and triangle diagrams cancel)
- First discovered in 1970s at D=4+6/L
- Maximal conformal symmetry
- Hidden structure of the integrands

All of them can be obtained from 10dim superstring by compactification on a torus

Bern, Dixon & Co 10

Drummond, Henn, Korchemsky, Sokatchev 10

Arkani-Hamed 12

D=4 N=8 Supergravity

- On-shell finite up to 8 loops
- Similar to higher dim SYM

Object: Helicity Amplitudes on mass shell with arbitrary number of legs and loops

The case: Planar limit  $N_c \rightarrow \infty$ ,  $g_{YM}^2 \rightarrow 0$  and  $g_{YM}^2 N_c$  - fixed

The aim: to get all loop (exact) result

Study of higher dim SYM gives insight into quantum gravity

# UV divergences in all Loops

Spinor-helicity formalism: S-matrix elements

**D=4 N=4      No UV div      IR div on shell**

**D=6 N=2      UV div from 3 loops      No IR div**

**D=8 N=1      UV div from 1 loop      No IR div**

**D=10 N=1      UV div from 1 loop      No IR div**

All these theories are non-renormalizable by power counting

The coupling  $g^2$  has dimension  $[g^2] = \frac{1}{M^{D-4}}$

The aim: to get all loop (exact) result for the leading (at least) divs


# Perturbation Expansion for the 4-point Amplitudes for any D

$$A_4/A_4^{tree}$$

No bubbles  
No Triangles

First UV div at  
 $L=[6/(D-4)]$  loops

IR finite



T. Dennen Yu-yin Huang 10 ,  
S.Caron-Huot D.O'Connell 10

# Leading Divergences from Generalized «Renormalization Group»

- In renormalizable theories the leading divergences can be found from the 1-loop term due to the renormalization group, in particular, for a single coupling theory the coefficient of  $1/\epsilon^n$  in n loops is

$$\mathcal{R}'G = \sum_n \frac{a_n^{(n)}}{\epsilon^n} \quad a_n^{(n)} = (a_1^{(1)})^n$$

# Leading Divergences from Generalized «Renormalization Group»

- In renormalizable theories the leading divergences can be found from the 1-loop term due to the renormalization group, in particular, for a single coupling theory the coefficient of  $1/\epsilon^n$  in n loops is

$$\mathcal{R}'G = \sum_n \frac{a_n^{(n)}}{\epsilon^n} \quad a_n^{(n)} = (a_1^{(1)})^n$$

- In non-renormalizable theories the leading divergences can be also found from 1-loop due to locality and R-operation

$$\mathcal{R}'G = 1 - \sum_{\gamma} K\mathcal{R}'_{\gamma} + \sum_{\gamma, \gamma'} K\mathcal{R}'_{\gamma} K\mathcal{R}'_{\gamma'} - \dots,$$

# Leading Divergences from Generalized «Renormalization Group»

- In renormalizable theories the leading divergences can be found from the 1-loop term due to the renormalization group, in particular, for a single coupling theory the coefficient of  $1/\epsilon^n$  in n loops is

$$\mathcal{R}'G = \sum_n \frac{a_n^{(n)}}{\epsilon^n} \quad a_n^{(n)} = (a_1^{(1)})^n$$

- In non-renormalizable theories the leading divergences can be also found from 1-loop due to locality and R-operation

$$\mathcal{R}'G = 1 - \sum_{\gamma} K \mathcal{R}'_{\gamma} + \sum_{\gamma, \gamma'} K \mathcal{R}'_{\gamma} K \mathcal{R}'_{\gamma'} - \dots,$$

$$\mathcal{R}'G_n = \frac{A_n^{(n)} (\mu^2)^{n\epsilon}}{\epsilon^n} + \frac{A_{n-1}^{(n)} (\mu^2)^{(n-1)\epsilon}}{\epsilon^n} + \dots + \frac{A_1^{(n)} (\mu^2)^{\epsilon}}{\epsilon^n}$$

$$+ \frac{B_n^{(n)} (\mu^2)^{n\epsilon}}{\epsilon^{n-1}} + \frac{B_{n-1}^{(n)} (\mu^2)^{(n-1)\epsilon}}{\epsilon^{n-1}} + \dots + \frac{B_1^{(n)} (\mu^2)^{\epsilon}}{\epsilon^{n-1}}$$

Leading pole

SubLeading pole

+lower order terms

$$A_1^{(n)}, B_1^{(n)}$$

1-loop graph

$$B_2^{(n)}$$

2-loop graph

# SubLeading Divergences from Generalized «Renormalization Group»

- In non-renormalizable theories the leading divergences can be also found from 1-loop due to locality and R-operation

All terms like  $(\log \mu^2)^m / \epsilon^k$  should cancel

$$A_n^{(n)} = (-1)^{n+1} \frac{A_1^{(n)}}{n},$$

$$B_n^{(n)} = (-1)^n \left( \frac{2}{n} B_2^{(n)} + \frac{n-2}{n} B_1^{(n)} \right)$$



Leading pole  
from 1 loop  
diagrams



SubLeading pole  
from 2 loop  
diagrams

$$\mathcal{KR}'G_n = \sum_{k=1}^n \left( \frac{A_k^{(n)}}{\epsilon^n} + \frac{B_k^{(n)}}{\epsilon^{n-1}} \right) \equiv \frac{A_n^{(n)'}}{\epsilon^n} + \frac{B_n^{(n)'}}{\epsilon^{n-1}}.$$

$$A_n^{(n)'} = (-1)^{n+1} A_n^{(n)} = \frac{A_1^{(n)}}{n},$$

$$B_n^{(n)'} = \left( \frac{2}{n(n-1)} B_2^{(n)} + \frac{2}{n} B_1^{(n)} \right)$$

Just like in  
renormalizable  
theories one can  
deduce the  
leading,  
subleading, etc  
divergences from  
1, 2, etc diagrams

# Kinematically dependent renormalization

## One-loop box

$$st \square = \frac{st}{3!} \frac{1}{\epsilon}$$

Totally defined by 1 loop

Independent term

## Two-loop box

$$s^2t \left[ \text{two boxes side-by-side} \right] + st^2 \left[ \text{two boxes stacked} \right] = \frac{st}{3!4!} \left( \frac{s^2 + t^2}{\epsilon^2} + \frac{27/4 s^2 + 1/3 st + 27/4 t^2}{\epsilon} \right)$$

$R'$




$$\left( \frac{1}{3!\epsilon} \quad \frac{s}{4!\epsilon} \right)$$

**This is true to all orders of PT like in renormalizable theories via the locality of the counterterms due to the R-operation**

# Ladder diagrams (leading divs)

**D=8 N=1**

**Horizontal boxes**



**Leading poles**

$$A_n^{(n)} = s^{n-1} A_n$$

**1 loop box**

$$nA_n = -\frac{2}{4!}A_{n-1} + \frac{2}{5!} \sum_{k=1}^{n-2} A_k A_{n-1-k}, \quad n \geq 3$$

$$A_1 = \frac{1}{3!}$$

$$A_2 = -\frac{1}{3!4!}$$

$$A_3 = \frac{2}{33!4!4!} + \frac{2}{35!3!3!}$$

# Ladder diagrams (leading divs)

**D=8 N=1**

**Horizontal boxes**

$$A_n^{(n)} = s^{n-1} A_n$$

$$nA_n = -\frac{2}{4!}A_{n-1} + \frac{2}{5!} \sum_{k=1}^{n-2} A_k A_{n-1-k}, \quad n \geq 3$$

$$A_1 = 1/6$$

**1 loop box**

# Ladder diagrams (leading divs)

**D=8 N=1**

**Horizontal boxes**

$$A_n^{(n)} = s^{n-1} A_n$$

$$nA_n = -\frac{2}{4!}A_{n-1} + \frac{2}{5!} \sum_{k=1}^{n-2} A_k A_{n-1-k}, \quad n \geq 3$$

$$A_1 = 1/6$$

**1 loop box**

**Summation**

$$\Sigma_m(z) = \sum_{n=m}^{\infty} A_n (-z)^n$$

# Ladder diagrams (leading divs)

**D=8 N=1**

**Horizontal boxes**

$$A_n^{(n)} = s^{n-1} A_n$$

$$nA_n = -\frac{2}{4!}A_{n-1} + \frac{2}{5!} \sum_{k=1}^{n-2} A_k A_{n-1-k}, \quad n \geq 3$$

$$A_1 = 1/6$$

**1 loop box**

**Summation**

$$\Sigma_m(z) = \sum_{n=m}^{\infty} A_n (-z)^n$$

$$-\frac{d}{dz} \Sigma_3 = -\frac{2}{4!} \Sigma_2 + \frac{2}{5!} \Sigma_1 \Sigma_1. \quad \Sigma_3 = \Sigma_1 + A_1 z - A_2 z^2, \quad \Sigma_2 = \Sigma_1 + A_1 z, \quad A_1 = \frac{1}{3!}, \quad A_2 = -\frac{1}{3!4!}$$

$$\Sigma_A \equiv \Sigma_1$$

**Diff eqn**

$$\frac{d}{dz} \Sigma_A = -\frac{1}{3!} + \frac{2}{4!} \Sigma_A - \frac{2}{5!} \Sigma_A^2$$

$$z = g^2 s^2 / \epsilon$$

# Ladder diagrams (leading divs)

**D=8 N=1**

**Horizontal boxes**

$$A_n^{(n)} = s^{n-1} A_n$$

$$nA_n = -\frac{2}{4!}A_{n-1} + \frac{2}{5!} \sum_{k=1}^{n-2} A_k A_{n-1-k}, \quad n \geq 3$$

$$A_1 = 1/6$$

**1 loop box**

**Summation**

$$\Sigma_m(z) = \sum_{n=m}^{\infty} A_n (-z)^n$$

$$-\frac{d}{dz} \Sigma_3 = -\frac{2}{4!} \Sigma_2 + \frac{2}{5!} \Sigma_1 \Sigma_1. \quad \Sigma_3 = \Sigma_1 + A_1 z - A_2 z^2, \quad \Sigma_2 = \Sigma_1 + A_1 z, \quad A_1 = \frac{1}{3!}, \quad A_2 = -\frac{1}{3!4!}$$

$$\Sigma_A \equiv \Sigma_1$$

**Diff eqn**

$$\frac{d}{dz} \Sigma_A = -\frac{1}{3!} + \frac{2}{4!} \Sigma_A - \frac{2}{5!} \Sigma_A^2$$

$$z = g^2 s^2 / \epsilon$$

$$\Sigma_A(z) = -\sqrt{5/3} \frac{4 \tan(z/(8\sqrt{15}))}{1 - \tan(z/(8\sqrt{15}))\sqrt{5/3}} = \sqrt{10} \frac{\sin(z/(8\sqrt{15}))}{\sin(z/(8\sqrt{15}) - z_0)}$$

$$\Sigma(z) = -(z/6 + z^2/144 + z^3/2880 + 7z^4/414720 + \dots)$$

$$z_0 = \arcsin(\sqrt{3/8})$$

# All loop Exact Recurrence Relation

**D=8 N=1**

**s-channel term**  $S_n(s, t)$  **t-channel term**  $T_n(s, t)$   $T_n(s, t) = S_n(t, s)$

**Exact relation for ALL diagrams**

$$\begin{aligned}
 nS_n(s, t) &= -2s^2 \int_0^1 dx \int_0^x dy y(1-x) (S_{n-1}(s, t') + T_{n-1}(s, t'))|_{t'=tx+yu} \\
 + s^4 \int_0^1 dx x^2(1-x)^2 \sum_{k=1}^{n-2} \sum_{p=0}^{2k-2} \frac{1}{p!(p+2)!} \frac{d^p}{dt'^p} (S_k(s, t') + T_k(s, t')) \times \\
 S_1 &= \frac{1}{12}, T_1 = \frac{1}{12} \times \frac{d^p}{dt'^p} (S_{n-1-k}(s, t') + T_{n-1-k}(s, t'))|_{t'=-sx} (tsx(1-x))^p
 \end{aligned}$$

# All loop Exact Recurrence Relation

**D=8 N=1**

**s-channel term**  $S_n(s, t)$  **t-channel term**  $T_n(s, t)$   $T_n(s, t) = S_n(t, s)$

**Exact relation for ALL diagrams**

$$\begin{aligned}
 nS_n(s, t) &= -2s^2 \int_0^1 dx \int_0^x dy y(1-x) (S_{n-1}(s, t') + T_{n-1}(s, t'))|_{t'=tx+yu} \\
 + s^4 \int_0^1 dx x^2(1-x)^2 &\sum_{k=1}^{n-2} \sum_{p=0}^{2k-2} \frac{1}{p!(p+2)!} \frac{d^p}{dt'^p} (S_k(s, t') + T_k(s, t')) \times \\
 S_1 = \frac{1}{12}, T_1 = \frac{1}{12} &\times \frac{d^p}{dt'^p} (S_{n-1-k}(s, t') + T_{n-1-k}(s, t'))|_{t'=-sx} (tsx(1-x))^p
 \end{aligned}$$

**summation**

# All loop Exact Recurrence Relation

**D=8 N=1**

**s-channel term**  $S_n(s, t)$  **t-channel term**  $T_n(s, t)$   $T_n(s, t) = S_n(t, s)$

**Exact relation for ALL diagrams**

$$\begin{aligned}
 nS_n(s, t) &= -2s^2 \int_0^1 dx \int_0^x dy y(1-x) (S_{n-1}(s, t') + T_{n-1}(s, t'))|_{t'=tx+yu} \\
 + s^4 \int_0^1 dx x^2(1-x)^2 \sum_{k=1}^{n-2} \sum_{p=0}^{2k-2} \frac{1}{p!(p+2)!} \frac{d^p}{dt'^p} (S_k(s, t') + T_k(s, t')) \times \\
 S_1 &= \frac{1}{12}, T_1 = \frac{1}{12} \times \frac{d^p}{dt'^p} (S_{n-1-k}(s, t') + T_{n-1-k}(s, t'))|_{t'=-sx} (tsx(1-x))^p
 \end{aligned}$$

**summation**  $\Sigma_3(s, t, z) = \Sigma_1(s, t, z) - S_2(s, t)z^2 + S_1(s, t)z$ ,  $\Sigma_2(s, t, z) = \Sigma_1(s, t, z) + S_1(s, t)z$

# All loop Exact Recurrence Relation

**D=8 N=1**

**s-channel term**  $S_n(s, t)$  **t-channel term**  $T_n(s, t)$   $T_n(s, t) = S_n(t, s)$

**Exact relation for ALL diagrams**

$$\begin{aligned}
 nS_n(s, t) &= -2s^2 \int_0^1 dx \int_0^x dy y(1-x) (S_{n-1}(s, t') + T_{n-1}(s, t'))|_{t'=tx+yu} \\
 + s^4 \int_0^1 dx x^2(1-x)^2 \sum_{k=1}^{n-2} \sum_{p=0}^{2k-2} \frac{1}{p!(p+2)!} \frac{d^p}{dt'^p} (S_k(s, t') + T_k(s, t')) \times \\
 S_1 = \frac{1}{12}, T_1 = \frac{1}{12} &\times \frac{d^p}{dt'^p} (S_{n-1-k}(s, t') + T_{n-1-k}(s, t'))|_{t'=-sx} (tsx(1-x))^p
 \end{aligned}$$




**summation**  $\Sigma_3(s, t, z) = \Sigma_1(s, t, z) - S_2(s, t)z^2 + S_1(s, t)z$ ,  $\Sigma_2(s, t, z) = \Sigma_1(s, t, z) + S_1(s, t)z$

**Diff eqn**


$$\begin{aligned}
 \frac{d}{dz} \Sigma(s, t, z) &= -\frac{1}{12} + 2s^2 \int_0^1 dx \int_0^x dy y(1-x) (\Sigma(s, t', z) + \Sigma(t', s, z))|_{t'=tx+yu} \\
 -s^4 \int_0^1 dx x^2(1-x)^2 \sum_{p=0}^{\infty} \frac{1}{p!(p+2)!} \left( \frac{d^p}{dt'^p} (\Sigma(s, t', z) + \Sigma(t', s, z))|_{t'=-sx} \right)^2 (tsx(1-x))^p.
 \end{aligned}$$

# All loop Solution (leading divs)


**D=8 N=1**



**PT and Pade versus ladder for t=s**



$$z = \frac{g^2}{\epsilon}$$



$$\Sigma_L(s, z) = -\sqrt{5/3} \frac{4 \tan(zs^2 / (8\sqrt{15}))}{1 - \tan(zs^2 / (8\sqrt{15})) \sqrt{5/3}}$$

# Subleading divergences

$$\Sigma_L(z) + \epsilon \Sigma_{NL}(z) + \epsilon^2 \Sigma_{NNL}(z) + \dots$$

$$\Sigma(z) = \sum_n^{\infty} z^n F_n$$

$$D = 4 \quad N = 4 \quad z = g^2/\epsilon$$

$$D = 6 \quad N = 2 \quad z = g^2 s/\epsilon, z = g^2 t/\epsilon$$

$$D = 8 \quad N = 1 \quad z = g^2 s^2/\epsilon, z = g^2 st/\epsilon, ..$$

$$D = 10 \quad N = 1 \quad z = g^2 s^3/\epsilon, z = g^2 s^2 t/\epsilon, ..$$

**D=8 N=1**

**sLadder case**

$$\Sigma_{NL} = s \Sigma_{sB}(z) + t \Sigma_{tB}(z)$$

$$z = \frac{g^2 s^2}{\epsilon}$$

# Subleading divergences

$$\Sigma_L(z) + \epsilon \Sigma_{NL}(z) + \epsilon^2 \Sigma_{NNL}(z) + \dots \qquad \Sigma(z) = \sum_n z^n F_n$$

$$D = 4 \quad N = 4 \quad z = g^2/\epsilon$$

$$D = 6 \quad N = 2 \quad z = g^2 s/\epsilon, z = g^2 t/\epsilon$$

$$D = 8 \quad N = 1 \quad z = g^2 s^2/\epsilon, z = g^2 st/\epsilon, ..$$

$$D = 10 \quad N = 1 \quad z = g^2 s^3/\epsilon, z = g^2 s^2 t/\epsilon, ..$$

**D=8 N=1**

**sLadder case**  $\Sigma_{NL} = s \Sigma_{sB}(z) + t \Sigma_{tB}(z) \qquad z = \frac{g^2 s^2}{\epsilon}$

$$\Sigma'_{tB}(z) = \frac{5}{6} \left[ e^{z/60} (2 \cos(z/30) - \sin(z/30)) - 2 \right]$$

$$\Sigma_{tB} = -\frac{1}{36} \left[ 60 + z + e^{z/60} (-(60 + z) \cos(z/30) - 2(-15 + z) \sin(z/30)) \right]$$

# Sum of Ladder diagrams (subleading divs)

$$\Sigma'_{sB} = \sum_{n=2}^{\infty} z^n B'_{sn}$$

$$\frac{d^2 \Sigma'_{sB}(z)}{dz^2} + f_1(z) \frac{d \Sigma'_{sB}(z)}{dz} + f_2(z) \Sigma'_{sB}(z) = f_3(z)$$

**Diff eqn**

$$f_1(z) = -\frac{1}{6} + \frac{\Sigma_A}{15},$$

$$f_2(z) = \frac{1}{80} - \frac{\Sigma_A}{360} + \frac{\Sigma_A^2}{600} + \frac{1}{15} \frac{d \Sigma_A}{dz},$$

$$f_3(z) = \frac{2321}{5!5!2} \Sigma_A + \frac{11}{1800} \Sigma'_{tB} - \frac{47}{5!45} \Sigma_A^2 - \frac{1}{5!72} \Sigma_A \Sigma'_{tB} + \frac{23}{6750} \Sigma_A^3 + \frac{1}{1200} \Sigma_A^2 \Sigma'_{tB} \\ - \frac{19}{36} \frac{d \Sigma_A}{dz} - \frac{1}{15} \frac{d \Sigma'_{tB}}{dz} + \frac{23}{225} \frac{d \Sigma_A^2}{dz} + \frac{1}{30} \frac{d(\Sigma_A \Sigma'_{tB})}{dz} - \frac{3}{32}$$

# Sum of Ladder diagrams (subleading divs)

$$\Sigma'_{sB} = \sum_{n=2}^{\infty} z^n B'_{sn}$$

$$\frac{d^2 \Sigma'_{sB}(z)}{dz^2} + f_1(z) \frac{d \Sigma'_{sB}(z)}{dz} + f_2(z) \Sigma'_{sB}(z) = f_3(z)$$

**Diff eqn**

$$f_1(z) = -\frac{1}{6} + \frac{\Sigma_A}{15},$$

$$f_2(z) = \frac{1}{80} - \frac{\Sigma_A}{360} + \frac{\Sigma_A^2}{600} + \frac{1}{15} \frac{d \Sigma_A}{dz},$$

$$f_3(z) = \frac{2321}{5!5!2} \Sigma_A + \frac{11}{1800} \Sigma'_{tB} - \frac{47}{5!45} \Sigma_A^2 - \frac{1}{5!72} \Sigma_A \Sigma'_{tB} + \frac{23}{6750} \Sigma_A^3 + \frac{1}{1200} \Sigma_A^2 \Sigma'_{tB} \\ - \frac{19}{36} \frac{d \Sigma_A}{dz} - \frac{1}{15} \frac{d \Sigma'_{tB}}{dz} + \frac{23}{225} \frac{d \Sigma_A^2}{dz} + \frac{1}{30} \frac{d(\Sigma_A \Sigma'_{tB})}{dz} - \frac{3}{32}$$

**Solution to Diff eqn**

$$\Sigma'_{sB}(z) = \frac{d \Sigma_A}{dz} u(z)$$

$$u(z) = \int_0^z dy \int_0^y dx \frac{f_3(x)}{d \Sigma_A(x)/dx}$$

**smooth monotonic function**



# Scheme dependence and arbitrariness of subtraction

## subleading case

$$A'_1 + B'_{s1} = \frac{1}{6\epsilon}(1 + c_1\epsilon) \quad \Delta\Sigma'_{sB} = c_1 z \frac{d\Sigma'_A}{dz} \quad \longrightarrow \quad z \rightarrow z(1 + c_1\epsilon).$$

## sub-subleading case

$$A'_2 + B'_2 = \frac{s}{3!4!\epsilon^2} \left( 1 - \frac{5}{12}\epsilon + 2c_1\epsilon + c_2\epsilon^2 \right) \quad \Delta\Sigma'_{sC} = c_2 z^2 \frac{d\Sigma'_A}{dz}.$$

$$\longrightarrow \quad z \rightarrow z(1 + c_1\epsilon) + z^2 c_2 \epsilon^2.$$

$$\Delta\Sigma'_{sC} = -c_1^2 \frac{z}{4!} \left( \frac{d\Sigma'_A}{dz} - 12 \frac{d^2\Sigma'_A}{dz^2} \right) \quad \longrightarrow \quad z \rightarrow z(1 + c_1\epsilon) + z^2 (c_2 + c_1^2/4!) \epsilon^2$$

# Scheme dependence and arbitrariness of subtraction

## subleading case

$$A'_1 + B'_{s1} = \frac{1}{6\epsilon}(1 + c_1\epsilon) \quad \Delta\Sigma'_{sB} = c_1 z \frac{d\Sigma'_A}{dz} \quad \longrightarrow \quad z \rightarrow z(1 + c_1\epsilon).$$

## sub-subleading case

$$A'_2 + B'_2 = \frac{s}{3!4!\epsilon^2} \left( 1 - \frac{5}{12}\epsilon + 2c_1\epsilon + c_2\epsilon^2 \right) \quad \Delta\Sigma'_{sC} = c_2 z^2 \frac{d\Sigma'_A}{dz}.$$

$$\longrightarrow \quad z \rightarrow z(1 + c_1\epsilon) + z^2 c_2 \epsilon^2.$$

$$\Delta\Sigma'_{sC} = -\frac{c_1^2 z}{4!} \left( \frac{d\Sigma'_A}{dz} - 12 \frac{d^2 \Sigma'_A}{dz^2} \right) \quad \longrightarrow \quad z \rightarrow z(1 + c_1\epsilon) + z^2 (c_2 + c_1^2/4!) \epsilon^2$$

# Scheme dependence and arbitrariness of subtraction

**sub-subleading case**

linear term

$$A'_2 + B'_2 = \frac{s}{3!4!\epsilon^2} \left( 1 - \frac{5}{12}\epsilon + 2c_1\epsilon + c_2\epsilon^2 \right)$$

new contribution from subleading term

$$\Delta\Sigma'_{sC}(3-loop) = -\frac{719c_1s^2}{1036800\epsilon}$$



$$\Sigma'_{sB}(3-loop) = -\frac{71s^2}{345600\epsilon^2}$$

$$\Sigma'^{trunc}_{sB}(3-loop) = -\frac{719s^2}{3110400\epsilon^2}$$

the source of a problem

$$\Delta\Sigma'_{sC}(3-loop) = c_1 z \frac{d\Sigma'^{trunc}_{sB}}{dz}(3-loop)$$

$$z \rightarrow z(1 + c_1\epsilon) + z^2(c_2 - c_1^2/4!)\epsilon^2 + z^3 c_1^3/6!\epsilon^3 - z^4 c_1^4/4!6!\epsilon^4 + \dots$$

# Scheme dependence and arbitrariness of subtraction

**sub-subleading case**

linear term


$$A'_2 + B'_2 = \frac{s}{3!4!\epsilon^2} \left( 1 - \frac{5}{12}\epsilon + 2c_1\epsilon + c_2\epsilon^2 \right)$$

new contribution from subleading term

$$\Delta\Sigma'_{sC}(3-loop) = -\frac{719c_1s^2}{1036800\epsilon}$$



$$\Sigma'_{sB}(3-loop) = -\frac{71s^2}{345600\epsilon^2}$$



$$\Sigma'^{trunc}_{sB}(3-loop) = -\frac{719s^2}{3110400\epsilon^2}$$

the source of a problem

$$\Delta\Sigma'_{sC}(3-loop) = c_1 z \frac{d\Sigma'^{trunc}_{sB}}{dz}(3-loop)$$

$$z \rightarrow z(1 + c_1\epsilon) + z^2(c_2 - c_1^2/4!)\epsilon^2 + z^3c_1^3/6!\epsilon^3 - z^4c_1^4/4!6!\epsilon^4 + \dots$$

# Kinematically dependent renormalization

- R-operation is equivalent to

**renormalizable theories**

**nonrenormalizable theories**

$$\bar{A}_4 = Z_4(g^2) \bar{A}_4^{bare} \Big|_{g_{bare}^2 \rightarrow g^2} Z_4$$

$$g_{bare}^2 = \mu^\epsilon Z_4(g^2) g^2.$$

$$Z = 1 - \sum_i KR'G_i$$

**simple multiplication**

**operator multiplication**

$$Z = 1 + \frac{g^2}{\epsilon} + g^4 \left( \frac{1}{\epsilon^2} + \frac{1}{\epsilon} \right) + \dots$$

$$Z = 1 + \frac{g^2}{\epsilon} st + g^4 st \left( \frac{s^2 + t^2}{\epsilon^2} + \frac{s^2 + st + t^2}{\epsilon} \right) + \dots$$

$$\frac{g^2}{\epsilon} (D_\rho D_\sigma F_{\mu\nu})^2$$

# Kinematically dependent renormalization

## operator kinematically dependent renormalization

at 2 loops

$$\bar{A}_4 = 1 - \frac{g_B^2 st}{3!\epsilon} - \frac{g_B^4 st}{3!4!} \left( \frac{s^2 + t^2}{\epsilon^2} + \frac{27/4s^2 + 1/3st + 27/4t^2}{\epsilon} \right) + \dots$$

$$\bar{A}_4 = Z_4(g^2) \bar{A}_4^{bare} |_{g_{bare}^2 \rightarrow g^2 Z_4}$$

$$Z_4 = 1 + \frac{g^2 st}{3!\epsilon} + \frac{g^4 st}{3!4!} \left( -\frac{s^2 + t^2}{\epsilon^2} + \frac{5/12s^2 + 1/3st + 5/12t^2}{\epsilon} \right)$$

$$g_B^2 = g^2 \left( 1 + \frac{g^2}{3!\epsilon} \right)$$

$$\frac{g^2 st}{\epsilon} \square \Rightarrow \frac{g^2}{\epsilon} \left( s \triangle + t \nabla \right)$$


**this is operator action!**

$$R' \quad \square \square = \square \square - \textcircled{\square} \triangleleft$$

**compare with R-operation**

# Kinematically dependent renormalization

## Two-loop box operator action



**Z-operator reproduces R-operation like in renormalizable theories**

# Kinematically dependent renormalization

## renormalizable theories

### scheme dependence

$$g^2 = zg'^2, \quad z = 1 + g'^2 c_1 + g'^4 c_2 + \dots$$

**infinite number of free parameters lead to a single multiplication constant -> redefinition of a single coupling**

## nonrenormalizable theories

### scheme dependence

$$g^2 = zg'^2, \quad z = 1 + g'^2 st c_1 + g'^4 st(s^2 + t^2) c_2 + \dots$$

**infinite number of free parameters lead to a single multiplication constant acting as an operator -> redefinition of a series of couplings**

# Conclusions

# Conclusions

- **The structure of UV divergences in non-renormalizable theories essentially copies that of renormalizable ones**

# Conclusions

- **The structure of UV divergences in non-renormalizable theories essentially copies that of renormalizable ones**
- **The main difference is that the renormalization constant  $Z$  depends on kinematics and acts like an operator rather than simple multiplication**

# Conclusions

- **The structure of UV divergences in non-renormalizable theories essentially copies that of renormalizable ones**
- **The main difference is that the renormalization constant  $Z$  depends on kinematics and acts like an operator rather than simple multiplication**
- **As a result, one can construct the higher derivative theory that gives the finite scattering amplitudes with a single arbitrary coupling  $g$  defined in PT within the given renormalization scheme.**

# Conclusions

- **The structure of UV divergences in non-renormalizable theories essentially copies that of renormalizable ones**
- **The main difference is that the renormalization constant  $Z$  depends on kinematics and acts like an operator rather than simple multiplication**
- **As a result, one can construct the higher derivative theory that gives the finite scattering amplitudes with a single arbitrary coupling  $g$  defined in PT within the given renormalization scheme.**
- **Transition to another scheme is performed by the action on the amplitude of a finite renormalization operator  $z$  that depends on kinematics.**

# Conclusions

- **The structure of UV divergences in non-renormalizable theories essentially copies that of renormalizable ones**
- **The main difference is that the renormalization constant  $Z$  depends on kinematics and acts like an operator rather than simple multiplication**
- **As a result, one can construct the higher derivative theory that gives the finite scattering amplitudes with a single arbitrary coupling  $g$  defined in PT within the given renormalization scheme.**
- **Transition to another scheme is performed by the action on the amplitude of a finite renormalization operator  $z$  that depends on kinematics.**
- **Assuming that one accepts these arguments, there is still a problem that at each order of PT the amplitude increases with energy, thus violating unitarity. However, apparently, this problem has to be addressed after summation of the whole PT series.**