Multi-particle Production (& Odderon) in the Multi-Regge Limit

Agustín Sabio Vera

Universidad Autónoma de Madrid, Instituto de Física Teórica UAM/CSIC

Seminar at "New Frontiers in Physics", Crete, July 7th, 2018

1. Multi-Regge limit
2. Monte Carlo event generator BFKLex
3. Collinear double logs
4. Odderon, open spin chains & high energy complexity

1. Multi-Regge limit
2. Monte Carlo event generator BFKLex
3. Collinear double logs
4. Odderon, open spin chains & high energy complexity

Regge theory preludes QCD. Pomeron in terms of quarks & gluons?

Perturbation theory with large scale $Q > \Lambda_{\rm QCD} \to \alpha_s(Q) \ll 1$.

$s \gg t, Q^2 \to \alpha_s(Q) \log\left(\frac{s}{t}\right) \sim \mathcal{O}(1)$. Resummation needed.



$$\sigma_{\rm tot}^{\rm LL} = \sum_{n=0}^{\infty} C_n^{\rm LL} \alpha_s^n \int_{y_B}^{y_A} dy_1 \int_{y_B}^{y_1} dy_2 \ldots \int_{y_B}^{y_{n-1}} dy_n = \sum_{n=0}^{\infty} \frac{C_n^{\rm LL}}{n!} \underbrace{\alpha_s^n (y_A - y_B)^n}_{\rm LL}$$

Multi-Regge linked to elastic amplitudes via optical theorem:





New degree of freedom
= Reggeized gluon

Pomeron = Composite state

2-dim interaction Hamiltonian

DIS data: $F_2(x, Q^2) \simeq x^{-\lambda}$

A NLL Multi-Regge approach fits data well (Hentschinski-Salas-ASV)





Transition from a perturbative to a non-perturbative Pomeron not well understood. Need more exclusive observables.

Observable proposed (ASV)^{2006} (Schwennsen,ASV)^{2007}
as ideal to pin down BFKL

$$\mathcal{R}_{2,1} = \frac{<\cos(2\theta)>}{<\cos(\theta)>}$$





Confirmed in 2013

1. Multi-Regge limit
2. Monte Carlo event generator BFKLex
3. Collinear double logs
4. Odderon, open spin chains & high energy complexity

With Grigorios Chachamis

Effective Feynman rules:

Gluon Regge trajectory: $\omega\left(\vec{q}\right) = -\frac{\alpha_s N_c}{\pi} \log \frac{q^2}{\lambda^2}$

Modified propagators in the t-channel:

$$\left(\frac{s_i}{s_0}\right)^{\omega(t_i)} = e^{\omega(t_i)(y_i - y_{i+1})}$$



$$\left(\frac{\alpha_s N_c}{\pi}\right)^2 \int d^2\vec{k}_1 \frac{\theta\left(k_1^2 - \lambda^2\right)}{\pi k_1^2} \int d^2\vec{k}_2 \frac{\theta\left(k_2^2 - \lambda^2\right)}{\pi k_2^2} \delta^{(2)}\left(\vec{k}_A + \vec{k}_1 + \vec{k}_2 - \vec{k}_B\right)$$

$$\times \int_0^Y dy_1 \int_0^{y_1} dy_2 e^{\omega\left(\vec{k}_A\right)(Y - y_1)} e^{\omega\left(\vec{k}_A + \vec{k}_1\right)(y_1 - y_2)} e^{\omega\left(\vec{k}_A + \vec{k}_1 + \vec{k}_2\right) y_2}$$

$$\sigma(Q_1, Q_2, Y) = \int d^2\vec{k}_A d^2\vec{k}_B \underbrace{\phi_A(Q_1, \vec{k}_A)\,\phi_B(Q_2, \vec{k}_B)}_{\text{PROCESS-DEPENDENT}} \underbrace{f(\vec{k}_A, \vec{k}_B, Y)}_{\text{UNIVERSAL}}$$

$$f\left(\vec{k}_A, \vec{k}_B, Y\right) = \sum_n$$



$$= e^{\omega\left(\vec{k}_A\right)Y}\left\{\delta^{(2)}\left(\vec{k}_A - \vec{k}_B\right) + \sum_{n=1}^{\infty}\prod_{i=1}^{n}\frac{\alpha_s N_c}{\pi}\int d^2\vec{k}_i \frac{\theta\left(k_i^2 - \lambda^2\right)}{\pi k_i^2}\right.$$

$$\left.\times \int_0^{y_{i-1}} dy_i\, e^{\left(\omega\left(\vec{k}_A + \sum_{l=1}^{i}\vec{k}_l\right) - \omega\left(\vec{k}_A + \sum_{l=1}^{i-1}\vec{k}_l\right)\right)y_i}\,\delta^{(2)}\left(\vec{k}_A + \sum_{l=1}^{n}\vec{k}_l - \vec{k}_B\right)\right\}$$

BFKLex: Monte Carlo implementation of full NLO BFKL

NLL is more complicated, sensitive to the running & choice of energy scale:

$$\sigma_{\text{tot}}^{\text{NLL}} = \sum_{n=1}^{\infty} \frac{\mathcal{C}_n^{\text{LL}}(\mathbf{k}_i)}{n!} \left(\alpha_s - \mathcal{A}\alpha_s^2\right)^n (y_A - y_B - \mathcal{B})^n$$

$$= \sigma_{\text{tot}}^{\text{LL}} - \sum_{n=1}^{\infty} \frac{\left(\mathcal{B}\,\mathcal{C}_n^{\text{LL}}(\mathbf{k}_i) + (n-1)\,\mathcal{A}\,\mathcal{C}_{n-1}^{\text{LL}}(\mathbf{k}_i)\right)}{(n-1)!} \underbrace{\alpha_s^n (y_A - y_B)^{n-1}}_{\text{NLL}}$$

besides, quarks enter the game ...
All of this is captured by Quasi-Multi-Regge kinematics.

NLL example: non-forward $N = 4$ SUSY in adjoint rep.

$$\mathcal{F}(\mathbf{q}_1,\mathbf{q}_2;\mathbf{q};\mathrm{Y}) = \left(\frac{\mathbf{q}^2\lambda^2}{\mathbf{q}_1^2\mathbf{q}_1'^2}\right)^{\frac{\bar{\alpha}}{2}\left(1-\frac{\zeta_2}{2}\bar{\alpha}\right)\mathrm{Y}} e^{\frac{3}{4}\zeta_3\bar{\alpha}^2\mathrm{Y}}\left\{\delta^{(2)}(\mathbf{q}_1-\mathbf{q}_2)\right.$$

$$+\sum_{n=1}^{\infty}\prod_{i=1}^{n}\left[\int d^2\mathbf{k}_i\frac{\bar{\alpha}}{4}\left(1-\frac{\zeta_2}{2}\bar{\alpha}\right)\frac{\theta\left(\mathbf{k}_i^2-\lambda^2\right)}{\pi\mathbf{k}_i^2}\left(1+\frac{\left(\mathbf{q}_1'+\sum_{l=1}^{i-1}\mathbf{k}_l\right)^2(\mathbf{q}_1+\sum_{l=1}^{i}\mathbf{k}_l)^2-\mathbf{q}^2\mathbf{k}_i^2}{(\mathbf{q}_1'+\sum_{l=1}^{i}\mathbf{k}_l)^2\left(\mathbf{q}_1+\sum_{l=1}^{i-1}\mathbf{k}_l\right)^2}\right)\right.$$

$$\left.\left.+\Phi\left(\mathbf{q}_1+\sum_{l=1}^{i-1}\mathbf{k}_l,\mathbf{q}_1+\sum_{l=1}^{i}\mathbf{k}_l\right)\right]\delta^{(2)}\left(\mathbf{q}_1+\sum_{l=1}^{n}\mathbf{k}_l-\mathbf{q}_2\right)\right.$$

$$\left.\times\int_0^{y_{i-1}}dy_i\left(\frac{\left(\mathbf{q}_1+\sum_{l=1}^{i-1}\mathbf{k}_l\right)^2}{\left(\mathbf{q}_1+\sum_{l=1}^{i}\mathbf{k}_l\right)^2}\right)^{1+\frac{\bar{\alpha}y_i}{2}\left(1-\frac{\zeta_2}{2}\bar{\alpha}\right)}\left(\frac{\left(\mathbf{q}_1'+\sum_{l=1}^{i-1}\mathbf{k}_l\right)^2}{\left(\mathbf{q}_1'+\sum_{l=1}^{i}\mathbf{k}_l\right)^2}\right)^{\frac{\bar{\alpha}y_i}{2}\left(1-\frac{\zeta_2}{2}\bar{\alpha}\right)}\right\}$$

$$
\begin{aligned}
&\Phi\left(\mathbf{q}_1,\mathbf{q}_1+\mathbf{k}\right) \;=\; \\
&\frac{\bar{\alpha}^2}{32\pi}\frac{1}{\mathbf{q}_1^2(\mathbf{k}+\mathbf{q}_1')^2}\Bigg\{\mathbf{q}^2\Bigg[\ln\left(\frac{\mathbf{q}_1^2}{\mathbf{q}^2}\right)\ln\left(\frac{\mathbf{q}_1'^2}{\mathbf{q}^2}\right)+\ln\left(\frac{\left(\mathbf{q}_1+\mathbf{k}\right)^2}{\mathbf{q}^2}\right)\ln\left(\frac{\left(\mathbf{q}_1'+\mathbf{k}\right)^2}{\mathbf{q}^2}\right) \\
&+\frac{1}{2}\ln^2\left(\frac{\mathbf{q}_1^2}{\left(\mathbf{q}_1+\mathbf{k}\right)^2}\right)+\frac{1}{2}\ln^2\left(\frac{\mathbf{q}_1'^2}{\left(\mathbf{q}_1'+\mathbf{k}\right)^2}\right)\Bigg]+\frac{1}{2}\frac{\left(\mathbf{q}_1^2\left(\mathbf{q}_1'+\mathbf{k}\right)^2-\mathbf{q}_1'^2\left(\mathbf{q}_1+\mathbf{k}\right)^2\right)}{\mathbf{k}^2} \\
&\quad\times\left[\ln\left(\frac{\mathbf{q}_1'^2}{\left(\mathbf{q}_1'+\mathbf{k}\right)^2}\right)\ln\left(\frac{\mathbf{q}_1'^2\left(\mathbf{q}_1'+\mathbf{k}\right)^2}{\mathbf{k}^4}\right)-\ln\left(\frac{\mathbf{q}_1^2}{\left(\mathbf{q}_1+\mathbf{k}\right)^2}\right)\ln\left(\frac{\mathbf{q}_1^2\left(\mathbf{q}_1+\mathbf{k}\right)^2}{\mathbf{k}^4}\right)\right] \\
&-\frac{\left(\mathbf{q}_1^2\left(\mathbf{q}_1'+\mathbf{k}\right)^2+\mathbf{q}_1'^2\left(\mathbf{q}_1+\mathbf{k}\right)^2\right)}{\mathbf{k}^2}\left[\ln^2\left(\frac{\mathbf{q}_1^2}{\left(\mathbf{q}_1+\mathbf{k}\right)^2}\right)+\ln^2\left(\frac{\mathbf{q}_1'^2}{\left(\mathbf{q}_1'+\mathbf{k}\right)^2}\right)\right] \\
&+\Bigg[\mathbf{q}^2\left(\mathbf{k}^2-\mathbf{q}_1^2-\left(\mathbf{q}_1+\mathbf{k}\right)^2\right)+2\mathbf{q}_1^2\left(\mathbf{q}_1+\mathbf{k}\right)^2-\mathbf{q}_1^2\left(\mathbf{q}_1'+\mathbf{k}\right)^2-\mathbf{q}_1'^2\left(\mathbf{q}_1+\mathbf{k}\right)^2 \\
&\quad+\frac{\left(\mathbf{q}_1^2\left(\mathbf{q}_1'+\mathbf{k}\right)^2-\mathbf{q}_1'^2\left(\mathbf{q}_1+\mathbf{k}\right)^2\right)}{\mathbf{k}^2}\left(\mathbf{q}_1^2-\left(\mathbf{q}_1+\mathbf{k}\right)^2\right)\Bigg]\mathcal{I}\left(\mathbf{q}_1^2,\left(\mathbf{q}_1+\mathbf{k}\right)^2,\mathbf{k}^2\right) \\
&+\Bigg[\mathbf{q}^2\left(\mathbf{k}^2-\mathbf{q}_1'^2-\left(\mathbf{q}_1'+\mathbf{k}\right)^2\right)+2\mathbf{q}_1'^2\left(\mathbf{q}_1'+\mathbf{k}\right)^2-\mathbf{q}_1'^2\left(\mathbf{q}_1+\mathbf{k}\right)^2-\mathbf{q}_1^2\left(\mathbf{q}_1'+\mathbf{k}\right)^2 \\
&\quad+\frac{\left(\mathbf{q}_1'^2\left(\mathbf{q}_1+\mathbf{k}\right)^2-\mathbf{q}_1^2\left(\mathbf{q}_1'+\mathbf{k}\right)^2\right)}{\mathbf{k}^2}\left(\mathbf{q}_1'^2-\left(\mathbf{q}_1'+\mathbf{k}\right)^2\right)\Bigg]\mathcal{I}\left(\mathbf{q}_1'^2,\left(\mathbf{q}_1'+\mathbf{k}\right)^2,\mathbf{k}^2\right)\Bigg\}
\end{aligned}
$$

$$
\mathcal{I}\left(\mathbf{p}^2,\mathbf{q}^2,\mathbf{r}^2\right) \;=\; \int_0^1\frac{dx}{\mathbf{p}^2(1-x)+\mathbf{q}^2x-\mathbf{r}^2x(1-x)}\ln\left(\frac{\mathbf{p}^2(1-x)+\mathbf{q}^2x}{\mathbf{r}^2x(1-x)}\right).
$$





Cut Pomeron: Number of emissions?

Pomeron: Number of rungs?





BFKL: $\mathbf{k}_i$ dependence in coefficients. $\sigma_{\rm tot}^{\rm LL} = \sum_{n=0}^{\infty} \frac{C_n^{\rm BFKL}(\mathbf{k}_i)}{n!} \alpha_s^n \left(y_A - y_B\right)^n$

Simple model for BFKL multiplicity distributions:



Virtual contributions tame the growth with Y.

Poisson distributions due to gluon Reggeization

Reggeized (virtual) gluon $|p_T|$ at a given rapidity?





Growth with energy? Depends on the azimuthal angle Fourier component:

$$f_n\left(|\vec{k}_A|, |\vec{k}_B|, Y\right) = \int_0^{2\pi} \frac{d\theta}{2\pi} f\left(\vec{k}_A, \vec{k}_B, Y\right) \cos\left(n\,\theta\right)$$



This is a distinct feature of BFKL

BFKL



CCFM



All CCFM projections grow with energy, not in BFKL - 1102.1890

Observables only sensitive to $n > 0$ single out BFKL

1. Multi-Regge limit
2. Monte Carlo event generator BFKLex
3. Collinear double logs
4. Odderon, open spin chains & high energy complexity

With Grigorios Chachamis

We can extend the formalism to include collinear regions

$$f = e^{\omega\left(\vec{k}_A\right)Y}\left\{\delta^{(2)}\left(\vec{k}_A-\vec{k}_B\right)+\sum_{n=1}^{\infty}\prod_{i=1}^{n}\frac{\alpha_s N_c}{\pi}\int d^2\vec{k}_i\,\frac{\theta\left(k_i^2-\lambda^2\right)}{\pi k_i^2}\right.$$
$$\left.\times\int_0^{y_{i-1}}dy_i\,e^{\left(\omega\left(\vec{k}_A+\sum_{l=1}^{i}\vec{k}_l\right)-\omega\left(\vec{k}_A+\sum_{l=1}^{i-1}\vec{k}_l\right)\right)y_i}\,\delta^{(2)}\left(\vec{k}_A+\sum_{l=1}^{n}\vec{k}_l-\vec{k}_B\right)\right\}$$

Key at NLL: $\theta\left(k_i^2-\lambda^2\right)\to\theta\left(k_i^2-\lambda^2\right)-\underbrace{\frac{\bar{\alpha}_s}{4}\ln^2\left(\frac{\vec{k}_A^2}{\left(\vec{k}_A+\vec{k}_i\right)^2}\right)}_{\text{NLL}}$

Resum it to all orders (SV-0507317):

$$\theta\left(k_i^2-\lambda^2\right)\to\theta\left(k_i^2-\lambda^2\right)+\sum_{n=1}^{\infty}\frac{\left(-\bar{\alpha}_s\right)^n}{2^n n!(n+1)!}\ln^{2n}\left(\frac{\vec{k}_A^2}{\left(\vec{k}_A+\vec{k}_i\right)^2}\right)$$

It corresponds to a Bessel function $J_1\left(\sqrt{2\bar{\alpha}_s\ln^2\left(\frac{\vec{k}_A^2}{\left(\vec{k}_A+\vec{k}_i\right)^2}\right)}\right)$

$$\sigma(Q_1, Q_2, Y) = \int d^2\mathbf{k}_a d^2\mathbf{k}_b \, \phi_A(Q_1, \mathbf{k}_a) \, \phi_B(Q_2, \mathbf{k}_b) \, f(\mathbf{k}_a, \mathbf{k}_b, Y)$$



Important to go beyond the MRK limit (Ciafaloni, Colferai, Salam, Stasto).
For BFKL domain we need "δ-like" impact factors $\phi_{A,B}$ & $Q_1 \simeq Q_2$.



Average transverse momentum of emitted mini-jets?

$$\langle p_t \rangle = \frac{1}{N} \sum_{i=1}^{N} |k_i|$$



Average rapidity separation among emitted mini-jets?

$$\langle \mathcal{R}_y \rangle = \frac{1}{N+1} \sum_{i=1}^{N+1} \frac{y_i}{y_{i-1}} \quad \simeq 1 + \frac{\Delta}{Y} \ln \frac{\Delta}{Y}$$

if $Y \simeq N\Delta$ in MRK and $Y \gg \Delta$





Higher $\langle \mathcal{R}_y \rangle_{\max}$ for higher energies: $\Delta_{\rm LO} \simeq 0.62$, $\Delta_{\rm LO+DLs} \simeq 0.81$

Lower mini-jet multiplicity when including higher order corrections

1. Multi-Regge limit
2. Monte Carlo event generator BFKLex
3. Collinear double logs
4. Odderon, open spin chains & high energy complexity

With Grigorios Chachamis

Amplitudes in the Generalized Leading Logarithmic Approximation (GLLA)



This is an old standing problem in High Energy QCD
Mapped onto a Closed Spin Chain
(Lipatov, Faddeev, Korchemsky,
Janik, Wosiek, Kotanski, Derkachov, Manashov ...)

Monte Carlo integration can be applied in this case (Chachamis-ASV)[2016]
Let us consider singlet exchange in t-channel with 3 Reggeized gluons:

ODDERON

Solution of the IR-finite Bartels-Kwiecinski-Praszalowicz equation:

$$\begin{aligned}
\left(\omega - \omega(\mathbf{p}_1) - \omega(\mathbf{p}_2) - \omega(\mathbf{p}_3)\right) f_\omega\left(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\right) \quad = \\
\delta^{(2)}\left(\mathbf{p}_1 - \mathbf{p}_4\right) \delta^{(2)}\left(\mathbf{p}_2 - \mathbf{p}_5\right) \delta^{(2)}\left(\mathbf{p}_3 - \mathbf{p}_6\right) \\
+ \int d^2\mathbf{k}\, \xi\left(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{k}\right) f_\omega\left(\mathbf{p}_1 + \mathbf{k}, \mathbf{p}_2 - \mathbf{k}, \mathbf{p}_3\right) \\
+ \int d^2\mathbf{k}\, \xi\left(\mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_1, \mathbf{k}\right) f_\omega\left(\mathbf{p}_1, \mathbf{p}_2 + \mathbf{k}, \mathbf{p}_3 - \mathbf{k}\right) \\
+ \int d^2\mathbf{k}\, \xi\left(\mathbf{p}_1, \mathbf{p}_3, \mathbf{p}_2, \mathbf{k}\right) f_\omega\left(\mathbf{p}_1 + \mathbf{k}, \mathbf{p}_2, \mathbf{p}_3 - \mathbf{k}\right)
\end{aligned}$$

Square of Lipatov's emission vertex:

$$\xi\left(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{k}\right) = \frac{\alpha_s N_c}{4} \frac{\theta(\mathbf{k}^2 - \lambda^2)}{\pi^2 \mathbf{k}^2} \left(1 + \frac{(\mathbf{p}_1 + \mathbf{k})^2 \mathbf{p}_2^2 - (\mathbf{p}_1 + \mathbf{p}_2)^2 \mathbf{k}^2}{\mathbf{p}_1^2 (\mathbf{k} - \mathbf{p}_2)^2}\right)$$

Gluon Regge trajectory: $\omega(\mathbf{p}) = -\frac{\bar{\alpha}_s}{2} \ln \frac{\mathbf{p}^2}{\lambda^2}$

Multi-particle Production (& Odderon) in the Multi-Regge Limit



$$f\left(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3,Y\right) = \int_{a-i\infty}^{a+i\infty}\frac{d\omega}{2\pi i}e^{\omega Y}f_\omega\left(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3\right)$$

$$\begin{aligned} f\left(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3,Y\right) &= \\ & e^{(\omega(\mathbf{p}_1)+\omega(\mathbf{p}_2)+\omega(\mathbf{p}_3))Y}\delta^{(2)}\left(\mathbf{p}_1-\mathbf{p}_4\right)\delta^{(2)}\left(\mathbf{p}_2-\mathbf{p}_5\right)\delta^{(2)}\left(\mathbf{p}_3-\mathbf{p}_6\right) \\ & +\sum_{n=1}^{\infty}\left\{\prod_{i=1}^{n}\int_0^{y_{i-1}}dy_i\int d^2\mathbf{k}_i e^{(\omega(\mathbf{p}_1)+\omega(\mathbf{p}_2)+\omega(\mathbf{p}_3))(y_{i-1}-y_i)}\mathcal{O}(\mathbf{k}_i)\otimes\right\} \\ & e^{(\omega(\mathbf{p}_1)+\omega(\mathbf{p}_2)+\omega(\mathbf{p}_3))y_n}\delta^{(2)}\left(\mathbf{p}_1-\mathbf{p}_4\right)\delta^{(2)}\left(\mathbf{p}_2-\mathbf{p}_5\right)\delta^{(2)}\left(\mathbf{p}_3-\mathbf{p}_6\right) \end{aligned}$$

$$\begin{aligned} \mathcal{O}(\mathbf{k})\otimes f\left(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3\right) &\equiv \xi\left(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3,\mathbf{k}\right)f\left(\mathbf{p}_1+\mathbf{k},\mathbf{p}_2-\mathbf{k},\mathbf{p}_3\right) \\ &+ \xi\left(\mathbf{p}_2,\mathbf{p}_3,\mathbf{p}_1,\mathbf{k}\right)f\left(\mathbf{p}_1,\mathbf{p}_2+\mathbf{k},\mathbf{p}_3-\mathbf{k}\right) \\ &+ \xi\left(\mathbf{p}_1,\mathbf{p}_3,\mathbf{p}_2,\mathbf{k}\right)f\left(\mathbf{p}_1+\mathbf{k},\mathbf{p}_2,\mathbf{p}_3-\mathbf{k}\right) \end{aligned}$$



$$= \delta^{(2)}\left(\mathbf{p}_1 - \mathbf{p}_4\right)\delta^{(2)}\left(\mathbf{p}_2 - \mathbf{p}_5\right)\delta^{(2)}\left(\mathbf{p}_3 - \mathbf{p}_6\right)e^{(\omega(\mathbf{p}_1)+\omega(\mathbf{p}_2)+\omega(\mathbf{p}_3))Y}$$



$$= \int d^2\mathbf{k}_1 \int_0^Y dy_1 \delta^{(2)}\left(\mathbf{p}_3 - \mathbf{p}_6\right) \delta^{(2)}\left(\mathbf{k}_1 + \mathbf{p}_1 - \mathbf{p}_4\right) \delta^{(2)}\left(-\mathbf{k}_1 + \mathbf{p}_2 - \mathbf{p}_5\right)$$

$$\times \quad \xi\left(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{k}_1\right) e^{\omega(\mathbf{p}_3)Y} e^{(\omega(\mathbf{k}_1+\mathbf{p}_1)+\omega(\mathbf{p}_2-\mathbf{k}_1))y_1} e^{(\omega(\mathbf{p}_1)+\omega(\mathbf{p}_2))(Y-y_1)}$$



$$
\begin{aligned}
= & \int d^2\mathbf{k}_1 \int d^2\mathbf{k}_2 \int_0^Y dy_1 \int_0^{y_1} dy_2 \delta^{(2)}\left(k_1+p_1-p_4\right)\delta^{(2)}\left(-k_1+k_2+p_2-p_5\right) \\
\times & \ \delta^{(2)}\left(-k_2+p_3-p_6\right)\xi\left(p_1,p_2,p_3,k_1\right)\xi\left(p_2-k_1,p_3,k_1+p_1,k_2\right) \\
\times & \ e^{(\omega(p_1)+\omega(p_2))(Y-y_1)}e^{\omega(p_3)(Y-y_2)}e^{\omega(k_1+p_1)y_1}e^{\omega(p_2-k_1)(y_1-y_2)}e^{(\omega(-k_1+k_2+p_2)+\omega(p_3-k_2))y_2}
\end{aligned}
$$

We can pick up two configurations with different transferred momentum:

$$\begin{aligned}
\mathbf{q} &= (4,0) \\
\mathbf{p}_1 &= (10,0) \\
\mathbf{p}_2 &= (20,\pi) \\
\mathbf{p}_3 &= (\mathbf{q}-\mathbf{p}_1)-\mathbf{p}_2 = (14,0) \\
\mathbf{p}_4 &= (20,0) \\
\mathbf{p}_5 &= (25,\pi) \\
\mathbf{p}_6 &= (\mathbf{q}-\mathbf{p}_4)-\mathbf{p}_5 = (9,0)
\end{aligned}
\qquad
\begin{aligned}
\mathbf{q} &= (31,0) \\
\mathbf{p}_1 &= (10,0) \\
\mathbf{p}_2 &= (20,\pi) \\
\mathbf{p}_3 &= (\mathbf{q}-\mathbf{p}_1)-\mathbf{p}_2 = (41,0) \\
\mathbf{p}_4 &= (20,0) \\
\mathbf{p}_5 &= (25,\pi) \\
\mathbf{p}_6 &= (\mathbf{q}-\mathbf{p}_4)-\mathbf{p}_5 = (36,0).
\end{aligned}$$



and study its growth with energy ...

Our solution must contain previous solutions in the literature.
They are singled out by particular impact factors.



On-going work on phenomenological applications ...

Very similar Reggeon diagrams appear in $N = 4$ SUSY amplitudes (Bartels, Lipatov, SV)[2009,2010]

Most complicated contributions at higher order & arbitrary number of legs:



Mapped onto an Open Spin Chain (Lipatov)
MC integration can be applied also here (Chachamis-ASV)[2018]
Let us consider octet exchange in t-channel with 3 Reggeized gluons:

EIGHT-POINT MHV AMPLITUDE

Solution of the IR-divergent BKP-like equation:

$$\begin{aligned}(\omega-\omega(\mathbf{p}_1)-\omega(\mathbf{p}_2)-\omega(\mathbf{p}_3))\,f_\omega(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3) \;=\;& \\ \delta^{(2)}(\mathbf{p}_1-\mathbf{p}_4)\,\delta^{(2)}(\mathbf{p}_2-\mathbf{p}_5)\,\delta^{(2)}(\mathbf{p}_3-\mathbf{p}_6)& \\ +\int d^2\mathbf{k}\,\xi(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3,\mathbf{k})\,f_\omega(\mathbf{p}_1+\mathbf{k},\mathbf{p}_2-\mathbf{k},\mathbf{p}_3)& \\ +\int d^2\mathbf{k}\,\xi(\mathbf{p}_2,\mathbf{p}_3,\mathbf{p}_1,\mathbf{k})\,f_\omega(\mathbf{p}_1,\mathbf{p}_2+\mathbf{k},\mathbf{p}_3-\mathbf{k})&\end{aligned}$$

is

$$\begin{aligned}f^{\text{adjoint}}_{\text{BKP}}(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3,Y) \;=\;& \\ e^{(\omega(\mathbf{p}_1)+\omega(\mathbf{p}_2)+\omega(\mathbf{p}_3))Y}\,\delta^{(2)}(\mathbf{p}_1-\mathbf{p}_4)\,\delta^{(2)}(\mathbf{p}_2-\mathbf{p}_5)\,\delta^{(2)}(\mathbf{p}_3-\mathbf{p}_6)& \\ +\sum_{n=1}^{\infty}\left\{\prod_{i=1}^{n}\int_0^{y_{i-1}}dy_i\int d^2\mathbf{k}_i\,e^{(\omega(\mathbf{p}_1)+\omega(\mathbf{p}_2)+\omega(\mathbf{p}_3))(y_{i-1}-y_i)}\mathcal{O}(\mathbf{k}_i)\otimes\right\}& \\ e^{(\omega(\mathbf{p}_1)+\omega(\mathbf{p}_2)+\omega(\mathbf{p}_3))y_n}\,\delta^{(2)}(\mathbf{p}_1-\mathbf{p}_4)\,\delta^{(2)}(\mathbf{p}_2-\mathbf{p}_5)\,\delta^{(2)}(\mathbf{p}_3-\mathbf{p}_6)&\end{aligned}$$

with

$$\begin{aligned}\mathcal{O}(\mathbf{k})\otimes f(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3) \;\equiv\;& \xi(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3,\mathbf{k})\,f(\mathbf{p}_1+\mathbf{k},\mathbf{p}_2-\mathbf{k},\mathbf{p}_3)\\ +\;& \xi(\mathbf{p}_2,\mathbf{p}_3,\mathbf{p}_1,\mathbf{k})\,f(\mathbf{p}_1,\mathbf{p}_2+\mathbf{k},\mathbf{p}_3-\mathbf{k})\end{aligned}$$

The IR divergencies factor out in a simple form:

$$f_{\mathrm{BKP}}^{\mathrm{adjoint}}(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3,\mathrm{Y}) = \left(\frac{\lambda^2}{\sqrt{\mathbf{p}_1^2\mathbf{p}_3^2}}\right)^{\frac{\bar{\alpha}_s \mathrm{Y}}{2}} \underbrace{\widehat{f}_{\mathrm{BKP}}^{\mathrm{adjoint}}(\mathbf{p}_1,\mathbf{p}_2,\mathbf{p}_3,\mathrm{Y})}_{\mathrm{IR-FINITE}}$$

where $\widehat{f}_{\mathrm{BKP}}^{\mathrm{adjoint}}$ is finite when $\lambda \to 0$.

Associated Laplacian matrix: Closed Chain with 3 Reggeons (Odderon)







$$\boxed{L} = \begin{pmatrix}
2 & -1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
-1 & 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & -1 & 3 & -1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\
0 & 0 & -1 & 2 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\
-1 & 0 & 0 & 0 & 3 & -1 & -1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & -1 & -1 & 3 & 0 & 0 & 0 & -1 & 0 & 0 \\
0 & 0 & 0 & 0 & -1 & 0 & 3 & -1 & 0 & 0 & -1 & 0 \\
0 & 0 & -1 & 0 & 0 & 0 & -1 & 3 & -1 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & -1 & 3 & 0 & -1 \\
0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 2 & -1 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & 2
\end{pmatrix}$$

Associated Laplacian matrix: Open Chain with 3 Reggeons







$$\boxed{L} = \begin{pmatrix} 2 & -1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 3 & -1 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 3 & -1 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 & -1 & 3 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 3 & -1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & -1 & 3 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 3 & -1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 2 \end{pmatrix}$$

Graph Complexity: Number of possible spanning trees
(Connected diagrams crossing all nodes with no loops)

Matrix Tree theorem by Kirchoff: determinant of any principal minor of L



[56]



[15]



[19]



[16]

Multi-particle Production (& Odderon) in the Multi-Regge Limit



[0]



[2.7 x 10^(-8)]



[3.2 x 10^(-8)]



[2.4 x 10^(-8)]



[3.7 x 10^(-8)]



[2.8 x 10^(-8)]



[3.2 x 10^(-8)]



[1.2 x 10^(-8)]



[4.3 x 10^(-8)]



[3.2 x 10^(-8)]



[3.7 x 10^(-8)]



[1.5 x 10^(-8)]



[4.2 x 10^(-8)]



[1.9 x 10^(-8)]



[2.3 x 10^(-8)]



[0]

Number of Rungs = 4

Complexity	# diagrams	Average weight in GGF
15	4	2.6×10^{-8}
16	2	3.3×10^{-8}
19	4	2.6×10^{-8}
23	2	3.2×10^{-8}
24	2	3.3×10^{-8}
56	2	0

A "Complexity Democracy" emerges ...

Average weight per complexity class for Reggeon webs.
Emerging scaling.



Due to integrability?



This is the solution to the "adjoint Odderon"
(open spin chain) relevant in $N = 4$ SUSY amplitudes.

1. Multi-Regge limit
2. Monte Carlo event generator BFKLex
3. Collinear double logs
4. Odderon, open spin chains & high energy complexity