

Total and differential cross sections for Higgs and top-quark production

Nikolaos Kidonakis

- Higher-order soft-gluon corrections
- Charged-Higgs production
- Single-top production and $t\bar{t}$ production



College of Science and Mathematics
Department of Physics

ICNFP 2018



Higher-order corrections

QCD corrections are very significant for Higgs and top production

Soft-gluon corrections are important
and they approximate exact results very well

We calculate/resum these soft corrections at NNLL accuracy
for the double-differential cross section

Finite-order expansions

Approximate NNLO (aNNLO) and $N^3\text{LO}$ ($aN^3\text{LO}$) predictions
for cross sections and differential distributions

Soft-gluon corrections

partonic processes

$$f_1(p_1) + f_2(p_2) \rightarrow t(p_t) + X$$

define

$$s = (p_1 + p_2)^2, \quad t = (p_1 - p_t)^2, \quad u = (p_2 - p_t)^2$$

and $s_4 = s + t + u - \sum m^2$

At partonic threshold $s_4 \rightarrow 0$

$$\text{Soft corrections} \left[\frac{\ln^k(s_4/m_t^2)}{s_4} \right]_+$$

For the order α_s^n corrections $k \leq 2n - 1$

Resum these soft corrections for the double-differential cross section

At NNLL accuracy we need two-loop soft anomalous dimensions

Soft-gluon Resummation

moments of the partonic cross section with moment variable N :

$$\hat{\sigma}(N) = \int (ds_4/s) e^{-Ns_4/s} \hat{\sigma}(s_4)$$

factorized expression for the cross section in $4 - \epsilon$ dimensions

$$\begin{aligned} \sigma^{f_1 f_2 \rightarrow tX}(N, \epsilon) &= H_{IL}^{f_1 f_2 \rightarrow tX}(\alpha_s(\mu_R)) S_{LI}^{f_1 f_2 \rightarrow tX}\left(\frac{m_t}{N\mu_F}, \alpha_s(\mu_R)\right) \\ &\quad \times \prod J_{\text{in}}(N, \mu_F, \epsilon) \prod J_{\text{out}}(N, \mu_F, \epsilon) \end{aligned}$$


$H_{IL}^{f_1 f_2 \rightarrow tX}$ is hard function and $S_{LI}^{f_1 f_2 \rightarrow tX}$ is soft function

S_{LI} satisfies the renormalization group equation

$$\left(\mu \frac{\partial}{\partial \mu} + \beta(g_s) \frac{\partial}{\partial g_s}\right) S_{LI} = -(\Gamma_S^\dagger)_{LK} S_{KI} - S_{LK} (\Gamma_S)_{KI}$$

Soft anomalous dimension Γ_S controls the evolution of the soft function which gives the exponentiation of logarithms of N

tH⁻ production



Born cross section for $bg \rightarrow tH^- \propto \alpha\alpha_s(m_b^2 \tan^2 \beta + m_t^2 \cot^2 \beta)$

$\tan \beta = v_2/v_1$ ratio of vevs of two Higgs doublets

$\Gamma_S^{bg \rightarrow tH^-} = (\alpha_s/\pi)\Gamma_S^{(1)} + (\alpha_s/\pi)^2\Gamma_S^{(2)} + \dots$, with


$$\Gamma_S^{(1)} = C_F \left[\ln \left(\frac{m_t^2 - t}{m_t \sqrt{s}} \right) - \frac{1}{2} \right] + \frac{C_A}{2} \ln \left(\frac{m_t^2 - u}{m_t^2 - t} \right)$$

$$\Gamma_S^{(2)} = \left[C_A \left(\frac{67}{36} - \frac{\zeta_2}{2} \right) - \frac{5}{18} n_f \right] \Gamma_S^{(1)} + C_F C_A \frac{(1 - \zeta_3)}{4}$$


tH^- production

Total cross sections at aNNLO

bg->tH⁻ at LHC aNNLO tanβ=30 μ=m_{H⁻}





bg->tH⁻ at LHC K-factor μ=m_{H⁻}



tH^- production


Total cross sections at aN³LO




tH^- production

Top-quark p_T and rapidity distributions


$bg \rightarrow tH^-$ at LHC top p_T aNNLO $\tan\beta=30$ $m_H=800$ GeV




$bg \rightarrow tH^-$ at LHC top rapidity aNNLO $\tan\beta=30$ $m_H=800$ GeV



H^-W^+ production




Total cross sections




H^-W^+ production

Charged-Higgs p_T and rapidity distributions


$b\bar{b} \rightarrow H^-W^+$ at LHC aNNLO $\tan\beta=1$ $m_H=500$ GeV



$b\bar{b} \rightarrow H^-W^+$ at LHC aNNLO $\tan\beta=1$ $m_H=500$ GeV



Single-top t -channel production



At one loop

$$\Gamma_{S11}^{t(1)} = C_F \left[\ln \left(\frac{t(t - m_t^2)}{m_t s^{3/2}} \right) - \frac{1}{2} \right] \quad \Gamma_{S12}^{t(1)} = \frac{C_F}{2N} \ln \left(\frac{u(u - m_t^2)}{s(s - m_t^2)} \right) \quad \Gamma_{S21}^{t(1)} = \ln \left(\frac{u(u - m_t^2)}{s(s - m_t^2)} \right)$$


$$\Gamma_{S22}^{t(1)} = C_F \ln \left(\frac{s - m_t^2}{m_t \sqrt{s}} \right) - \frac{1}{2N} \ln \left(\frac{t(t - m_t^2)}{s(s - m_t^2)} \right) + \frac{(N^2 - 2)}{2N} \ln \left(\frac{u(u - m_t^2)}{s(s - m_t^2)} \right) - \frac{C_F}{2}$$

At two loops


$$\Gamma_{S11}^{t(2)} = \left[C_A \left(\frac{67}{36} - \frac{\zeta_2}{2} \right) - \frac{5}{18} n_f \right] \Gamma_{S11}^{t(1)} + C_F C_A \frac{(1 - \zeta_3)}{4}$$

t-channel production at aNNLO


Single-top *t*-channel aNNLO cross sections $m_t=172.5$ GeV



Top p_T distributions in t -channel production at the LHC



Single-top s -channel production



At one loop

$$\Gamma_{S11}^{s(1)} = C_F \left[\ln \left(\frac{s - m_t^2}{m_t \sqrt{s}} \right) - \frac{1}{2} \right] \quad \Gamma_{S12}^{s(1)} = \frac{C_F}{2N} \ln \left(\frac{u(u - m_t^2)}{t(t - m_t^2)} \right) \quad \Gamma_{S21}^{s(1)} = \ln \left(\frac{u(u - m_t^2)}{t(t - m_t^2)} \right)$$


$$\Gamma_{S22}^{s(1)} = C_F \ln \left(\frac{s - m_t^2}{m_t \sqrt{s}} \right) - \frac{1}{N} \ln \left(\frac{u(u - m_t^2)}{t(t - m_t^2)} \right) + \frac{N}{2} \ln \left(\frac{u(u - m_t^2)}{s(s - m_t^2)} \right) - \frac{C_F}{2}$$

At two loops


$$\Gamma_{S11}^{s(2)} = \left[C_A \left(\frac{67}{36} - \frac{\zeta_2}{2} \right) - \frac{5}{18} n_f \right] \Gamma_{S11}^{s(1)} + C_F C_A \frac{(1 - \zeta_3)}{4}$$

s -channel production at aNNLO

Single-top s -channel aNNLO cross sections $m_t=172.5$ GeV



Associated tW production



At one loop


$$\Gamma_S^{tW^- (1)} = C_F \left[\ln \left(\frac{m_t^2 - t}{m_t \sqrt{s}} \right) - \frac{1}{2} \right] + \frac{C_A}{2} \ln \left(\frac{m_t^2 - u}{m_t^2 - t} \right)$$

At two loops

$$\Gamma_S^{tW^- (2)} = \left[C_A \left(\frac{67}{36} - \frac{\zeta_2}{2} \right) - \frac{5}{18} n_f \right] \Gamma_S^{tW^- (1)} + C_F C_A \frac{(1 - \zeta_3)}{4}$$


tW production at aN³LO

$tW^- + \bar{t}W^+$ aN³LO cross section $m_t=172.5$ GeV




Top quark p_T and rapidity distributions in tW production



Top p_T distribution in tW production at LHC aN^3LO $m_t=173.3$ GeV



Top rapidity distribution in tW production at LHC aN^3LO $m_t=173.3$ GeV




tZ production via anomalous couplings




$$\Delta\mathcal{L}^{eff} = \frac{1}{\Lambda} \kappa_{tqZ} e \bar{t} (i/2) (\gamma_\mu \gamma_\nu - \gamma_\nu \gamma_\mu) q F_Z^{\mu\nu} + h.c.$$


g u → t Z at LHC aNNLO $k_{tZ}=0.01$



g c → t Z at LHC aNNLO $k_{tZ}=0.01$



Top-antitop pair production



At one loop for $q\bar{q} \rightarrow t\bar{t}$

$$\Gamma_{11}^{q\bar{q}(1)} = \Gamma_{\text{cusp}}^{(1)}, \quad \Gamma_{12}^{q\bar{q}(1)} = \frac{C_F}{C_A} \ln\left(\frac{t_1}{u_1}\right), \quad \Gamma_{21}^{q\bar{q}(1)} = 2 \ln\left(\frac{t_1}{u_1}\right)$$

$$\Gamma_{22}^{q\bar{q}(1)} = \left(1 - \frac{C_A}{2C_F}\right) \Gamma_{\text{cusp}}^{(1)} + 4C_F \ln\left(\frac{t_1}{u_1}\right) - \frac{C_A}{2} \left[1 + \ln\left(\frac{sm_t^2 t_1^2}{u_1^4}\right)\right]$$

At two loops for $q\bar{q} \rightarrow t\bar{t}$


$$\Gamma_{11}^{q\bar{q}(2)} = \Gamma_{\text{cusp}}^{(2)}, \quad \Gamma_{12}^{q\bar{q}(2)} = \left(\frac{K}{2} - \frac{C_A}{2} N_{2l}\right) \Gamma_{12}^{q\bar{q}(1)}, \quad \Gamma_{21}^{q\bar{q}(2)} = \left(\frac{K}{2} + \frac{C_A}{2} N_{2l}\right) \Gamma_{21}^{q\bar{q}(1)}$$

$$\Gamma_{22}^{q\bar{q}(2)} = \frac{K}{2} \Gamma_{22}^{q\bar{q}(1)} + \left(1 - \frac{C_A}{2C_F}\right) \left(\Gamma_{\text{cusp}}^{(2)} - \frac{K}{2} \Gamma_{\text{cusp}}^{(1)}\right)$$

3×3 matrix for $gg \rightarrow t\bar{t}$


Top-antitop pair production

$pp \rightarrow t\bar{t}$ at LHC energies aN^3LO $m_t = 172.5$ GeV




Top p_T distributions in $t\bar{t}$ production


Normalized top p_T distribution at the LHC $S^{1/2}=13$ TeV




Normalized top p_T distribution at the LHC $S^{1/2}=8$ TeV







Normalized top p_T distribution at the LHC $S^{1/2}=8$ TeV



Normalized top p_T distribution at the LHC $S^{1/2}=8$ TeV



Top rapidity distributions in $t\bar{t}$ production



Summary

- cross sections and distributions for Higgs and top production
- soft-gluon corrections
- tH^- and H^-W^+ production at aNNLO and beyond
- t -channel and s -channel single top at aNNLO
- tW production at aN³LO
- tZ production via anomalous couplings
- $t\bar{t}$ production at aN³LO
- excellent agreement with collider data
- high-order corrections are very significant