

# CP Violation in $B \rightarrow \pi\pi\pi$ in QCD Factorization

Keri Vos

Universität Siegen

in collaboration with

Th. Mannel, J. Virto, R. Klein

arXiv:1708.02047




Theor. Physik 1



DFG FOR 1873

# Motivation




- CP violation in multibody decays provides more information
- Study with data-driven model-independent approach
  - using “partial” factorization Kraenkl, Mannel, Virto [2015]
  - first leading order study

# Dalitz distribution - Kinematics

Kraenkl, Mannel, Virto [2015]

- $B^+ \rightarrow \pi^+(k_1)\pi^-(k_2)\pi^+(k_3)$  Symmetric Dalitz plot
- Kinematic variables  $s_{+-}^{\text{low}} = \frac{(k_1+k_2)^2}{m_B^2}$  and  $s_{+-}^{\text{high}} = \frac{(k_2+k_3)^2}{m_B^2}$



# Naive-Factorization in three-body decays

Mannel, Virto, Klein, KKV [2017]

At leading order

$$\langle \pi^- \pi^+ \pi^- | (\bar{u}b)_{V-A} \times (\bar{d}u)_{V-A} | B^- \rangle = \langle \pi^- | (\bar{d}u)_{V-A} | 0 \rangle \langle \pi^- \pi^+ | (\bar{u}b)_{V-A} | B^- \rangle$$


$$\langle \pi^- \pi^+ \pi^- | (\bar{d}b)_{V-A} \times (\bar{u}u)_{V-A} | B^- \rangle = \langle \pi^- | (\bar{d}b)_{V-A} | B^- \rangle \langle \pi^- \pi^+ | (\bar{u}u)_{V-A} | 0 \rangle$$

**New non-perturbative input** → New strong phases!


- $B \rightarrow \pi\pi$  form factor (isoscalar and isovector)
- $2\pi$  LCDA (isovector only)
  - Normalized to time-like pion form factor  $F_\pi$
  - Experimentally from  $e^+e^- \rightarrow \pi\pi(\gamma)$  data

# Time-like pion formfactor $F_\pi(s)$

Hanhart, Kubis, Shekhovtsova, Roig, Was, Predzinski



Vector form factor




Phase


- No experimental data on the phase available

# Time-like pion form factor $F_\pi(s)$

Hanhart, Kubis, Shekhovtsova, Roig, Was, Predzinski



Vector form factor



Phase


- Isovector  $B \rightarrow \pi\pi$  form factor studied with LCSR Khodjamirian, Virto, Cheng

$$\text{Phase } F_\pi = \text{Phase } F_{B \rightarrow \pi\pi}^{I=1}$$


- Experimental information on isoscalar form factor?

# Isoscalar contribution

Daub, Hanhart, Kubis, Passemar, Cirigliano



Scalar form factor




Phase

$$\langle \pi^- \pi^+ | m_u \bar{u}u + m_d \bar{d}d | 0 \rangle = m_\pi^2 F_\pi^S .$$


- $F_\pi^S$  scalar pion form factor (analogous to  $F_\pi$ )
  - Dispersion theory, coupled Omnes-equations
  - Data on  $F_\pi^S$  only available up to 1.3 GeV

# Isoscalar contribution

Daub, Hanhart, Kubis, Passemar, Cirigliano



Scalar form factor



Phase



$$F_{B \rightarrow \pi\pi}^{I=0} \propto \beta e^{i\phi} F_{\pi}^S$$

- First study: Fit  $\beta$  and  $\phi$  to experimental data
- Information on  $\beta, \phi$  from fit to Dalitz projections





# Dalitz and CP Distributions

Mannel, Klein, Virto, KKV [2017]



# Dalitz and CP Distributions



$$A_{CP} \propto \beta \sin \gamma \sin \phi \cos \theta$$

- Our model only gives Vector-Scalar interferences
- Several extensions of our framework possible

# Outlook

- Study CPV in three-body decays in QCD factorization approach
  - Improve description of the unknown (isoscalar) inputs
  - Include  $\mathcal{O}(\alpha)$  corrections
  - Include higher-partial waves
  - Apply to  $B \rightarrow K\pi\pi, B \rightarrow D\pi\pi$
- Improved experimental data needed
  - Dalitz distributions with background and efficiency correction
  - Data in different kinematic regions
  - Connection with  $B \rightarrow \pi\pi\ell\nu$  or  $B \rightarrow \pi\pi\ell\ell$
  - Updated  $B \rightarrow \rho\pi$  measurements

# Outlook

- Study CPV in three-body decays in QCD factorization approach
  - Improve description of the unknown (isoscalar) inputs
  - Include  $\mathcal{O}(\alpha)$  corrections
  - Include higher-partial waves
  - Apply to  $B \rightarrow K\pi\pi, B \rightarrow D\pi\pi$
- Improved experimental data needed
  - Dalitz distributions with background and efficiency correction
  - Data in different kinematic regions
  - Connection with  $B \rightarrow \pi\pi\ell\nu$  or  $B \rightarrow \pi\pi\ell\ell$
  - Updated  $B \rightarrow \rho\pi$  measurements

Thank you for your attention

# Decay amplitude

At leading order, leading twist

$$\mathcal{A}_{s_{\pm}^{\text{low}} \ll 1} = \frac{G_F}{\sqrt{2}} m_B^2 \left[ f_{\pi} \frac{m_{\pi}}{m_B^2} (\lambda_u (a_1 + a_4^u) + \lambda_c a_4^c) F_t(s_{\pm}^{\text{low}}, \zeta) \right. \\ \left. + (\lambda_u (a_2 - a_4^u) - \lambda_c a_4^c) (2\zeta - 1) F_{\pi}(s_{\pm}^{\text{low}}) f_0(s_{\pm}^{\text{low}}) \right],$$

- $a_j$  as in two-body decay, contain perturbative strong phases  $\mathcal{O}(\alpha_s)$
- $\lambda_u = |\lambda_u| e^{i\gamma}$  weak phase
- Only 4 inputs that can be obtained from data
  - $B \rightarrow \pi$  form factor  $f_0$
  - Single pion DA gives the pion decay constant  $f_{\pi}$
  - $B \rightarrow \pi\pi$  form factor  $F_t$
  - $2\pi$  LCDA gives  $F_{\pi}$

# Direct CP Violation

$$\mathcal{A} \propto e^{i\gamma} |\mathcal{A}_u| e^{i\phi_u} + |\mathcal{A}_c| e^{i\phi_c}$$

- $\gamma$  weak phase from CKM
- $\mathcal{A}_u$  and  $\mathcal{A}_c$  from current-current and penguin operators with

$$\langle \pi\pi\pi | (\bar{b}u)(\bar{u}d) | B \rangle \text{ and } \langle \pi\pi\pi | (\bar{b}c)(\bar{u}c) | B \rangle$$

- CPV induced by non-perturbative phases in matrix elements
  - $B \rightarrow \pi\pi$  form factor (isoscalar and isovector)
  - $2\pi$  LCDA (isovector only)

$$A_{CP} \equiv \frac{|\mathcal{A}|^2 - |\bar{\mathcal{A}}|^2}{|\mathcal{A}|^2 + |\bar{\mathcal{A}}|^2} = \frac{2|\mathcal{A}_u||\mathcal{A}_c| \sin(\Delta\phi) \sin \Delta\gamma}{|\mathcal{A}_u|^2 + |\mathcal{A}_c|^2 + 2|\mathcal{A}_u||\mathcal{A}_c| \cos(\Delta\phi) \cos \Delta\gamma}$$

# $B \rightarrow \pi\pi$ Form factor: Isovector contributions

- Light-Cone Sum Rule Khodjamirian, Virto, Cheng

$$F_t(q^2, \zeta)^{I=1} = \frac{6m_b^2(2\zeta - 1)F_\pi(q^2)}{m_\pi f_B m_B^2} \int_{u_0}^1 \frac{du}{u} \bar{u} e^{\frac{m_B^2 - s(u)}{M^2}} (m_b^2 - m_\pi^2 + u^2 q^2)$$

$$s(u) \equiv \frac{m_b^2 - \bar{u}m_\pi^2 + u\bar{u}q^2}{u}$$

- Reduces to  $B \rightarrow \rho$  form factor in  $\rho$ -dominance, zero-width approximation

$$F_t^{I=1} \propto (2\zeta - 1)A_0^{B\rho} \frac{g_{\rho\pi\pi} m_\rho}{\sqrt{2}(m_\rho^2 - s - im_\rho\Gamma_\rho)} \propto (2\zeta - 1)F_\pi A_0^{B\rho}$$

# S-wave form factor model

$$\langle \pi^-(k_1)\pi^+(k_2)|\bar{u}u|0\rangle = \langle \pi^-(k_1)\pi^+(k_2)|S^0\rangle \text{BW}_S \langle S^0|\bar{u}u|0\rangle$$

$$\langle S^0|\bar{u}u|0\rangle = f_S m_S, \quad \langle \pi^-(k_1)\pi^+(k_2)|S^0\rangle = g_{S\pi^-\pi^+} m_S$$

$$F_\pi^S(q^2) = \frac{2m_u}{m_\pi^2} \frac{f_S m_S^2 g_{S\pi^-\pi^+}}{m_S^2 - q^2 - i\sqrt{q^2}\Gamma_S}$$



$$\begin{aligned} & \langle \pi^-(k_1)\pi^+(k_2)|J_\nu|B^-(p)\rangle \\ &= \langle \pi^-(k_1)\pi^+(k_2)|S^0\rangle \text{BW}_S \langle S^0(q)|J_\nu|B^-(p)\rangle \end{aligned}$$

Finally

$$F_t^{I=0}(q^2) = \frac{m_B^2}{m_\pi f_\pi} \beta e^{i\phi} F_\pi^S(q^2)$$



# Helicity Angle




$$k_3 \cdot (k_1 - k_2) = \frac{\beta_\pi}{2} \sqrt{\lambda} \cos \theta$$

# Discussion


- Difficult to generate rich structure in CP asymmetry
  - QCDF framework and form factors only reliable close to the edges
  - Simple scenarios to get qualitative picture
- Additional strong phases generated above charm threshold
- Scenario: Breit-Wigner shape

$$\mathcal{A}_c = \mathcal{A}_c^{(0)} + g \frac{4m_c^2}{m_B^2 s_{+-}^{\text{low}} - 4m_c^2 + im_c \Gamma}$$

# Charm model scenario



Scenario



Experimental Data

$$\mathcal{A}_c = \mathcal{A}_c^{(0)} + g \frac{4m_c^2}{m_B^2 s_{+-}^{\text{low}} - 4m_c^2 + im_c \Gamma}$$