

Search for electroweak production of supersymmetric particles in the two and three lepton final state at $\sqrt{s} = 13$ TeV with the ATLAS detector

Friday, January 5, 2018 6:10 PM (15 minutes)

A search for the electroweak production of charginos, neutralinos and sleptons decaying to final states involving two or three electrons or muons is presented. The analysis is based on 36.1 fb^{-1} of $\sqrt{s}=13$ TeV proton-proton collisions recorded by the ATLAS detector at the Large Hadron Collider. No significant deviations from the Standard Model expectation are observed and results are interpreted in a range of scenarios based on simplified models. Considered scenarios include the associated production of mass-degenerate next-to-lightest neutralino and lightest chargino, followed by their decays to final-state leptons and lightest neutralinos via either sleptons or Standard Model gauge bosons; direct production of chargino pairs, which in turn decay to leptons and lightest neutralinos via intermediate sleptons; and slepton pair production, where each slepton decays directly to the lightest neutralino and a lepton. Stringent limits at 95% confidence level are placed on the masses of relevant supersymmetric particles in each of these scenarios.

Summary

Primary author: VADLA, Knut Oddvar Hoie (University of Oslo (NO))

Presenters: VADLA, Knut Oddvar Hoie (University of Oslo (NO)); Dr GRAMSTAD, Eirik (University of Oslo (NO))

Session Classification: Friday PM

Track Classification: Particle physics - experiment