

EW, QCD, and interference contributions

Mathieu PELLE

Institute for Theoretical Physics and Astrophysics,
University Würzburg

Based on: [\[arXiv:1611.02951\]](#) *Phys.Rev.Lett.* **118** (2017) no.26, 261801,
[\[arXiv:1708.00268\]](#) *JHEP* **1710** (2017) 124

in collaboration with: Benedikt Biedermann and Ansgar Denner

[\[arXiv:1801.XXXX\]](#)

+ VBSCAN MC team (incomplete list): Brass, Dittmaier, **Grossi**, **Karlberg**,
Pelliccioli, **Rauch**, Reuter, **Rothe**, **Schwan**, **Stienemeier**, **Zaro**

Monte-Carlo description of VBS
Amsterdam, the Netherlands

16th of November 2017



Outline

- 1 Contributions to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 2 Complete NLO corrections to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 3 Large NLO EW corrections to VBS
- 4 Conclusion

Content


- 1 Contributions to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 2 Complete NLO corrections to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 3 Large NLO EW corrections to VBS
- 4 Conclusion

$pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$

→ All partonic channels taken into account

- $uu \rightarrow \mu^+ \nu_\mu e^+ \nu_e d\bar{d}$
- $u\bar{d} \rightarrow \mu^+ \nu_\mu e^+ \nu_e d\bar{u}$
- $u\bar{d} \rightarrow \mu^+ \nu_\mu e^+ \nu_e s\bar{c}$
- $u\bar{s} \rightarrow \mu^+ \nu_\mu e^+ \nu_e d\bar{c}$
- $uc \rightarrow \mu^+ \nu_\mu e^+ \nu_e s\bar{d}$
- $\bar{s}\bar{d} \rightarrow \mu^+ \nu_\mu e^+ \nu_e u\bar{c}$
- $\bar{d}\bar{d} \rightarrow \mu^+ \nu_\mu e^+ \nu_e u\bar{u}$

→ Tree amplitudes of order $\mathcal{O}(g^6)$ and $\mathcal{O}(g_s^2 g^4)$



$$pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$$

Three contributions at LO:

- $\mathcal{O}(\alpha^6)$ (= “EW contribution”)
- $\mathcal{O}(\alpha_s \alpha^5)$ (= “interference”)
- $\mathcal{O}(\alpha_s^2 \alpha^4)$ (= “QCD contribution”)

Cuts: inspired by Refs. [1405.6241, 1611.02428, 1410.6315, CMS-PAS-SMP-17-004]



charged lepton: $p_{T,\ell} > 20 \text{ GeV}$, $|y_\ell| < 2.5$, $\Delta R_{\ell\ell} > 0.3$

jets: $p_{T,j} > 30 \text{ GeV}$, $|y_j| < 4.5$, $\Delta R_{j\ell} > 0.3$

missing energy: $p_{T,\text{miss}} > 40 \text{ GeV}$,


+ Extra VBS cuts on m_{jj} and Δy_{jj}




→ “Play the experimentalist” and scan over m_{jj} and Δy_{jj}



VBSCAN MC team / full computation

→ Background contributions become very relevant on the peak






VBS CAN MC team / full computation

→ Experimentalist are doing a good job!

($m_{jj} > 500$ GeV and $|\Delta y_{jj}| > 2.5$ are good values)

- Check the validity of the VBS approximation (implemented in different variants in: BONSAY, POWHEG, VBFNLO, and MG_aMC@NLO)
- Comparison at LO: full vs. VBS approximation



VBSCAN MC team

- Reasonable agreement away from the resonance region

Message: do not use the VBS for $m_{jj} < 200 \text{ GeV}$ and $|\Delta y_{jj}| < 2.0$

→ Follow up

- NLO computations running (now):
 - full vs. VBS approximation (as implemented in BONSAI)
 - $m_{jj} > 200 \text{ GeV}$ and $|\Delta y_{jj}| > 2.0$
- Is the approximation failing?
- If yes, by how much?

→ Stay tuned

Content

- 1 Contributions to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 2 Complete NLO corrections to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 3 Large NLO EW corrections to VBS
- 4 Conclusion


→ Calculation of both NLO QCD and EW corrections to

$$pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$$

- Off-shell and non-resonant contributions
 - Realistic final state
- Full calculations vs. VBS approximation
- EW corrections can be large in certain phase space regions
 - Sudakov logarithms
- Theoretical and numerical challenge to consider $2 \rightarrow 6$ process
 - Virtual corrections involving up to 8-point functions

$$pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$$

LO cross sections at $\mathcal{O}(\alpha^6)$, $\mathcal{O}(\alpha_s \alpha^5)$, and $\mathcal{O}(\alpha_s^2 \alpha^4)$



NLO cross sections at $\mathcal{O}(\alpha^7)$, $\mathcal{O}(\alpha_s \alpha^6)$, $\mathcal{O}(\alpha_s^2 \alpha^5)$, and $\mathcal{O}(\alpha_s^3 \alpha^4)$

- Order $\mathcal{O}(\alpha_s \alpha^6)$ and $\mathcal{O}(\alpha_s^2 \alpha^5)$: QCD and EW corrections mix
- Combined measurement

Predictions for $\sqrt{s} = 13\text{TeV}$ at the LHC $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_{ejj}$

- NNPDF3.0QED [NNPDF collaboration]
- dynamical renormalisation and factorisation scale:

$$\mu_{\text{ren}} = \mu_{\text{fac}} = \sqrt{p_{T,j1} p_{T,j2}}$$

- Cuts inspired by Refs. [1405.6241, 1611.02428, 1410.6315, CMS-PAS-SMP-17-004]:

charged lepton: $p_{T,\ell} > 20 \text{ GeV}, \quad |y_\ell| < 2.5, \quad \Delta R_{\ell\ell} > 0.3$

jets: $p_{T,j} > 30 \text{ GeV}, \quad |y_j| < 4.5, \quad \Delta R_{jl} > 0.3$

missing energy: $p_{T,\text{miss}} > 40 \text{ GeV},$

→ For the two leading jet in p_T :

jet-jet: $m_{jj} > 500 \text{ GeV}, \quad |\Delta y_{jj}| > 2.5.$

→ Final state: 2 jets, missing p_T , and 2 same sign leptons

- anti- k_T jet algorithm [Cacciari, Salam, Soyez; 0802.1189]

$R = 0.4$ for jet recombination and $R = 0.1$ for photon recombination

→ LO fiducial cross sections:

Order	$\mathcal{O}(\alpha^6)$	$\mathcal{O}(\alpha_s \alpha^5)$	$\mathcal{O}(\alpha_s^2 \alpha^4)$	Sum
σ_{LO} [fb]	1.4178(2)	0.04815(2)	0.17229(5)	1.6383(2)



→ NLO fiducial cross sections: (normalised to σ_{LO})

Order	$\mathcal{O}(\alpha^7)$	$\mathcal{O}(\alpha_s \alpha^6)$	$\mathcal{O}(\alpha_s^2 \alpha^5)$	$\mathcal{O}(\alpha_s^3 \alpha^4)$	Sum
$\delta\sigma_{\text{NLO}}$ [fb]	-0.2169(3)	-0.0568(5)	-0.00032(13)	-0.0063(4)	-0.2804(7)
$\delta\sigma_{\text{NLO}}/\sigma_{\text{LO}}$ [%]	-13.2	-3.5	0.0	-0.4	-17.1

Updated with respect to Split presentation [Biedermann, Denner, MP; 1708.00268]

- Large EW corrections at $\mathcal{O}(\alpha^7)$
- Negative corrections at $\mathcal{O}(\alpha_s \alpha^6)$:
 - ~ 0.6% difference with respect to VBS approximation
(neglecting s -channel and t -/ u -channel interferences)
 - Tuned comparison against [Denner, et al.; 1209.2389] and [Jäger, et al.; 0907.0580]
 - VBS approximation in RECOLA
- Photon PDF contribution at NLO (not included in NLO definitions):
 +1.50% with LUXqed [Manohar et al.; 1607.04266]



Separated contributions



Updated with respect to Split presentation [Biedermann, Denner, MP; 1708.00268]

- Clear hierarchy of LO contributions
- Different behaviour of the NLO corrections (normalised to the full LO)

Combined predictions




Updated with respect to Split presentation [Biedermann, Denner, MP; 1708.00268]

- Large negative corrections for the full process
- Corrections dominated by EW correction to EW process
 - Bands do not overlap

Content

- 1 Contributions to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 2 Complete NLO corrections to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 3 Large NLO EW corrections to VBS
- 4 Conclusion

LO: $\mathcal{O}(\alpha^6)$	σ^{LO} [fb]	$\sigma_{\text{EW}}^{\text{NLO}}$ [fb]	δ_{EW} [%]
NLO: $\mathcal{O}(\alpha^7)$	1.5348(2)	1.2895(6)	-16.0




[Biedermann, Denner, MP; 1611.02951]

→ Huge NLO electroweak correction (!)


- Leading behaviour dominated by:
Sudakov logarithms (bosonic part of the virtual), $\log^2 \left(\frac{Q^2}{M_W^2} \right)$
 - Usually in the tail of the distribution (suppressed)
 - Usually small for total cross section
 - Usually smaller than the QCD corrections
- Large corrections not due to VBS cuts
 - remove $m_{jj} > 500 \text{ GeV}$ and $|\Delta y_{jj}| > 2.5$
 - relax $p_{T,j}$ and $p_{T,miss}$

- Double-pole approximation: [Dittmaier, Schwan; 1511.01698]
 leading contribution of expansion about the resonance poles
 → Required two W bosons for the virtual contributions



- Agree within 1% with full calculation
- Dominated by factorisable corrections
 → Large corrections driven by the scattering process

- Effective Vector Boson approximation:



- Simplify the discussion to $W^+W^+ \rightarrow W^+W^+$
- Leading logarithm approximation [Denner, Pozzorini; hep-ph/0010201]

$$\sigma_{LL} = \sigma_{LO} \left[1 - \frac{\alpha}{4\pi} 4C_W^{\text{ew}} \log^2 \left(\frac{Q^2}{M_W^2} \right) + \frac{\alpha}{4\pi} 2b_W^{\text{ew}} \log \left(\frac{Q^2}{M_W^2} \right) \right]$$

(double EW logs, collinear single EW logs, and single logs from parameter renormalisation included) (angular-dependant logarithms omitted)

$$\sigma_{LL} = \sigma_{LO} \left[1 - \frac{\alpha}{4\pi} 4C_W^{\text{ew}} \log^2 \left(\frac{Q^2}{M_W^2} \right) + \frac{\alpha}{4\pi} 2b_W^{\text{ew}} \log \left(\frac{Q^2}{M_W^2} \right) \right]$$


- For $Q = \langle m_{4\ell} \rangle \sim 390 \text{ GeV}$

$$\delta_{EW}^{LL} = -16\% (!)$$

→ Corrections 3-4 times larger than for $q\bar{q} \rightarrow W^+W^+$

- C^{ew} larger for bosons than fermions
- $\langle m_{4\ell} \rangle$ larger for VBS (massive t -channel [Denner, Hahn; hep-ph/9711302])
 NB: $\langle m_{4\ell} \rangle \sim 250 \text{ GeV}$ for $q\bar{q} \rightarrow W^+W^+$

Large NLO EW corrections:
 intrinsic feature of VBS at the LHC



[Biedermann, Denner, MP; 1611.02951]


- Near $y_{j_1 j_2} = 0$: two jets back-to-back
- Bulk of the cross section, $\sim -16\%$ corrections
- Band: $\pm 1/\sqrt{N_{\text{obs}}}$ for 3000 fb^{-1} → probe of the EW sector

Content

- 1 Contributions to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 2 Complete NLO corrections to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$
- 3 Large NLO EW corrections to VBS
- 4 Conclusion

Conclusion

- Study of the fiducial region
 - Validity of the VBS approximation
- Full NLO corrections to $pp \rightarrow \mu^+ \nu_\mu e^+ \nu_e jj$:
[Biedermann, Denner, MP; 1708.00268]
 - At NLO VBS and QCD-induced contributions mix:
 - Combined measurement
 - Full computations:
 - Small differences with respect to the VBS approximation
- NLO EW corrections to VBS:
[Biedermann, Denner, MP; 1611.02951]
 - Unexpected large EW corrections
 - Probe of the EW sector



Back-up slides

BACK-UP

- Tools

- Virtual corrections: RECOLA [Actis, Denner, Hofer, Lang, Scharf, Uccirati]
- + COLLIER [Denner, Dittmaier, Hofer]
- Private Monte Carlos (MoCANLO [Feger] + another one)
- Dipole subtraction scheme [Catani, Seymour], [Dittmaier]
- Complex-mass scheme [Denner et al.]

- Inputs

- G_μ scheme:

$$\alpha = \frac{\sqrt{2}}{\pi} G_\mu M_W^2 \left(1 - \frac{M_W^2}{M_Z^2} \right) \quad \text{with} \quad G_\mu = 1.16637 \times 10^{-5} \text{ GeV}^2$$



- Parameters:

$$\begin{aligned} m_t &= 173.21 \text{ GeV}, & \Gamma_t &= 0 \text{ GeV} \\ M_Z^{\text{OS}} &= 91.1876 \text{ GeV}, & \Gamma_Z^{\text{OS}} &= 2.4952 \text{ GeV} \\ M_W^{\text{OS}} &= 80.385 \text{ GeV}, & \Gamma_W^{\text{OS}} &= 2.085 \text{ GeV} \\ M_H &= 125 \text{ GeV} & \Gamma_H &= 4.07 \times 10^{-3} \text{ GeV} \end{aligned}$$

Validations


- Two independent Monte Carlo integrators
- Tree-level matrix elements: MADGRAPH5_AMC@NLO [Alwall et al.; 1405.0301]
- One-loop matrix elements:
 - vs. MADLOOPS [Hirschi et al.; 1103.0621]:
 - $\mathcal{O}(\alpha^7)$ and $\mathcal{O}(\alpha_s^3 \alpha^4)$
 - Two libraries in COLLIER [Denner, Dittmaier, Hofer; 1407.0087, 1604.06792]:
 - $\mathcal{O}(\alpha_s \alpha^6)$, $\mathcal{O}(\alpha_s^2 \alpha^5)$, and $\mathcal{O}(\alpha_s^3 \alpha^4)$
- NLO computations:
 - DPA for $\mathcal{O}(\alpha^7)$ (automatised in [MP et al.; 1607.05571, 1612.07138] following [Dittmaier, Schwan; 1511.01698])
 - $\mathcal{O}(\alpha_s \alpha^6)$ vs. [Denner, et al.; 1209.2389] in the VBS approximation
- IR-subtraction/finiteness:
 - Variation of α parameter [Nagy, Troscanyi; hep-ph/9806317]
 - Variation of technical cuts
 - Variation of IR-scale

Distributions extra




DPA (1) [Dittmaier, Schwan; 1511.01698]

- At LO



- At NLO



DPA (2) [Dittmaier, Schwan; 1511.01698]

- Factorisable corrections

$$\mathcal{M}_{\text{virt, fact, PA}} = \sum_{\lambda_1, \dots, \lambda_r} \left(\prod_{i=1}^r \frac{1}{K_i} \right) \left[\mathcal{M}_{\text{virt}}^{I \rightarrow N, \bar{R}} \prod_{j=1}^r \mathcal{M}_{\text{LO}}^{j \rightarrow R_j} \right. \\ \left. + \mathcal{M}_{\text{LO}}^{I \rightarrow N, \bar{R}} \sum_{k=1}^r \mathcal{M}_{\text{virt}}^{k \rightarrow R_k} \prod_{j \neq k}^r \mathcal{M}_{\text{LO}}^{j \rightarrow R_j} \right] \left\{ \bar{k}_l^2 \rightarrow \hat{k}_l^2 = M_l^2 \right\}_{l \in \bar{R}}$$

- Non-factorisable corrections:

$$2\text{Re} \{ \mathcal{M}_{\text{LO, PA}}^* \mathcal{M}_{\text{virt, nfact, PA}} \} = |\mathcal{M}_{\text{LO, PA}}|^2 \delta_{\text{nfact}}$$

- On-shell projection
- DPA applied to virtual corrections and I -operator
- Full Born and Real contributions: