


Isolated Photons with Jets in DIS at ZEUS

Oleg Kuprash
DESY (Hamburg, Germany)

Wilhelm und Else Heraeus Physics School on
Diffractive and electromagnetic processes at high energies, Heidelberg (Germany)
September 2 - 6, 2013

HERA collider



Electrons: 27.5 GeV

Protons: 920 GeV

$\sqrt{s} = 318$ GeV

Kinematics:

$$Q^2 = -q^2 = -(l - l')^2$$


$$y = \frac{Pq}{Pl}$$

$$x_{Bj} = \frac{Q^2}{2Pq}$$


$$Q^2 = x_{Bj}ys$$

$Q^2 \lesssim 1 \text{ GeV}^2$: photoproduction (PHP)

$Q^2 \gtrsim 1 \text{ GeV}^2$: DIS



Isolated photons production in DIS



- Isolated photon production:

- LL-radiation (ISR, FSR)
- QQ-radiation (incoming or outgoing quark)

- Isolated photons:

- do not undergo hadronisation process, therefore provide a direct probe of the underlying partonic process
- allow to test QCD matrix elements
- it is expected for isolated photons + jets to be more sensitive to the underlying partonic process, compared to inclusive photons
- *isolated photon data potentially could provide separation between u - and d -quark densities in proton*

Event selection and reconstruction (1/2)

- integrated luminosity of $\approx 330 \text{ pb}^{-1}$ (data taken between 2004-2007)

Observables:

- $Q^2 = -q^2 = -(l - l')^2$
- $x = \frac{Q^2}{2pq}$
- transverse energy E_T^γ and pseudorapidity η^γ of the photon
- transverse energy E_T^{jet} and pseudorapidity η^{jet} of the accompanying jet

Photon isolation:

- no tracks within $\Delta R(\eta, \phi) = 0.2$ cone around the photon candidate
- ratio of the energy of the photon candidate to the energy of the jet containing it greater than 0.9

Monte Carlo:

- Pythia for the simulation of the QQ-radiation
- Ariadne for the simulation of the LL-radiation and background

Event selection and reconstruction (2/2)

Deep inelastic scattering


- $10 < Q^2 < 350 \text{ GeV}^2$
- $-40 < Z_{\text{vtx}} < 40 \text{ cm}$
- $35 < E - p_z < 65 \text{ GeV}$
- $E_{\text{electron}} > 10 \text{ GeV}$
- $140^\circ < \theta_{el} < 180^\circ$

Photon

- $4 < E_T^\gamma / \text{GeV} < 15$
- $-0.7 < \eta^\gamma < 0.9$
- $\Delta R < 0.2$
- $\frac{E_{\text{EMC}}}{E_{\text{HAC}} + E_{\text{EMC}}} > 0.9$
- $\frac{E_T^\gamma}{E_{\text{jet containing } \gamma}} > 0.9$

Accompanying jet

- $E_T^{\text{jet}} > 2.5 \text{ GeV}$
- $-1.5 < \eta^{\text{jet}} < 1.8$



Background to isolated photons

□ Photons from decays of neutral mesons:

- $\pi_0 \rightarrow \gamma\gamma$ (98.8 %)
- $\eta \rightarrow \gamma\gamma$ (39.3 %)
- $\eta \rightarrow \pi_0\pi_0\pi_0$ (32.6 %)

→ it is the main source of the background

→ opening angle of two photons after π^0 decay:

$$\sin \frac{\phi}{2} = \frac{m}{E}$$

At $E = 5$ GeV $\phi = 1.55^\circ$ for π^0 and $\phi = 6.3^\circ$ for η -mesons

→ there is a possibility to use shower shape method



□ Photons from quark to photon fragmentation

→ this process occurs over long distances and cannot be calculated perturbatively

→ easy to suppress by applying of the isolation cut

Extraction of the signal

Following variables are using to describe the shower shape:




$$\langle \delta z \rangle = \frac{\sum |z_i - z_{cluster}| \cdot E_i}{l_{cell} \sum E_i}$$

$$f_{max} = \frac{\text{Energy in the most energetic BEMC cell}}{\text{Total energy of the cluster}}$$

- mixture of different type Monte Carlo events is used to fit the data distribution
- $\langle \delta z \rangle$ variable is used for the signal extraction, because it carries more information

Theoretical predictions: fixed order calculations

- Theoretical prediction of [A. Gehrmann-De Ridder, G. Kramer and H. Spiesberger \(Nucl. Phys. B. 578 \(2000\) 326\)](#) (GKS)
- $\text{LO}(\alpha^3)$ with three components:



- (LEFT) LL radiation, (MIDDLE) QQ radiation, (RIGHT) photon from jet fragmentation
- $\text{LO}(\alpha^3)$ and $\text{NLO}(\alpha^3\alpha_s)$ predictions are calculated

- renormalisation and factorisation scales $\mu_R = \mu_F = \sqrt{Q^2 + (p_T^{\text{jet}})^2}$

$$d\sigma = \sum_n \alpha_s^n \sum_{a=q,\bar{q}} \int dx f_a(x, \mu_F^2; \alpha_s) \cdot d\hat{\sigma}_a^{(n)}(xP, \mu_R, \mu_F)$$



- HERAPDF1.0 for PDF parametrisation

Theoretical predictions: k_T -factorisation approach

- Calculated by [S.P.Baranov, A.V.Lipatov, N.P.Zotov \(Phys.Rev.D81:094034,2010\)](#) (BLZ):
- investigation of the prompt photon production in DIS at HERA in the framework of k_T -factorisation QCD approach
- based on the off-shell partonic amplitude $eq^* \rightarrow e\gamma q$
- taken into account photon radiation from the leptons as well as from the quarks
- unintegrated proton parton densities are used in the KMR form

$$\sigma_{LL, QQ}(ep \rightarrow e\gamma X) = \sum_q \int \frac{1}{256\pi^3 x^2 s \sqrt{s} |\mathbf{p}_{\gamma T}| \exp(y_\gamma)} |\bar{\mathcal{M}}_{LL, QQ}(eq^* \rightarrow e\gamma q)|^2 \times \\ \times f_q(x, \mathbf{k}_T^2, \mu^2) d\mathbf{p}_{e'T}^2 d\mathbf{p}_{qT}^2 d\mathbf{k}_T^2 dy_{e'} dy_q \frac{d\phi_{e'}}{2\pi} \frac{d\phi_q}{2\pi} \frac{d\phi}{2\pi},$$

- In the k_T -factorisation approach the contribution from the quark radiation subprocess (QQ mechanism) is enhanced compared to the leading-order collinear approximation







$$4 < E_T^\gamma < 15 \text{ GeV}, \quad -0.7 < \eta^\gamma < 0.9, \quad \frac{E^\gamma}{E_{\text{jet containing } \gamma}} > 0.9,$$

$$10 < Q^2 < 350 \text{ GeV}^2, \quad E_{\text{elec}} > 10 \text{ GeV}, \quad 140^\circ < \theta_{\text{elec}} < 180^\circ,$$

$$E_T^{\text{jet}} > 2.5 \text{ GeV}, \quad -1.5 < \eta^{\text{jet}} < 1.8$$

- The width of the GKS NLO predictions represents theoretical uncertainty due to factorisation and renormalisation scales, varied independently by factor 2 up and down
- The width of the BLZ predictions shows the uncertainty due mainly to the procedure of fixing the rapidity of the jets from the evolution cascade in the factorisation approach
- GKS predictions systematically underestimate data and BLZ overestimate them

Cross sections (2/2) Phys. Lett. B 715 (2012) 88-97



- GKS predictions give better description of the η^{jet} shape

Including Isolated Photon + Jet Data into proton PDF fit

Under supervision of A. Geiser

Work in progress

Data: ZEUS Collaboration (Abramowicz, H. et al.). Measurement of isolated photons accompanied by jets in deep inelastic ep scattering, Phys Lett B 715 (2012) 88-97

Theory: A.Gehrmann-De Ridder, G.Kramer, H.Spiesberger. Photon plus jet cross sections in deep inelastic ep collisions at order $\mathcal{O}(\alpha^2\alpha_s)$, Nucl.Phys. B578 (2000) 326-350

+? running α_{em} from F. Jegerlehner. Electroweak Effective Couplings for Future Precision Experiments, DESY 11-117

Fitting proton PDFs: the procedure:

- HERAFitter program (www.herafitter.org) has been used to fit proton PDFs
- Input:
 - 1) Inclusive DIS HERAI data (NC + CC)
 - 2) Inclusive DIS HERAI data + data on isolated photon with jet production
- HERAFitter fits PDFs using χ^2 method, taking into account statistical and systematical uncertainties of the experimental data
- PDFs were parametrised using 13-parameter HERAPDF1.5-style function



Summary

Isolated photons with jets in DIS:





- cross sections of the production of isolated photons with jets in DIS have been measured and published by ZEUS
- predictions give adequate description of the data but systematically overestimate (for k_T -factorisation approach) or underestimate (for fixed order NLO calculations) them

Backup

Photons in DIS: comparison to MC models (1/2)



Photons in DIS: comparison to MC models (1/2)



Photons in DIS: systematics

Main sources:

- due to e , γ , jet energy scales: 5 – 7%
- the dependence on the modelling of the hadronic background by Ariadne was investigated by varying the upper limit for the $\langle \delta Z \rangle$ fit in the range 0.6, 1.0 giving typically variations of $\pm 5\%$ increasing to +12% and –14% in the most forward η^γ and highest- x bins respectively

Contributions from different flavours to the NLO cross section

