

Transverse Beam Dynamics II

The Theory of Synchrotrons:

Linear Beam Optics

The Beam as Particle Ensemble


Emittance and Beta-Function

Colliding Beams & Luminosity

„... how does it work ?“

„... does it ?“


typical values
in a strong
foc. machine:
 $x \approx \text{mm}$, $x' \leq \text{mrad}$



Resonance Problem:

Why do we have so stupid non-integer tunes ?
“Q = 64.0” sounds much better

Qualitatively spoken: Integer tunes lead to a resonant increase of the closed orbit amplitude in presence of the smallest dipole field error.




Orbit in case of a small dipole error:

$$x_{co}(s) = \frac{\sqrt{\beta(s)} * \int \frac{1}{\rho_{sl}} \sqrt{\beta_{sl}} * \cos(\psi_{sl} - \psi_s - \pi Q) ds}{2 \sin \pi Q}$$

Assume: Tune = integer $Q = 1 \rightarrow 0$

Tune and Resonances

To avoid resonance conditions the frequency of the transverse motion **must not be equal to** (or a integer multiple of) **the revolution frequency**



$$\begin{aligned} 1 * Q_x &= 1 \quad \rightarrow \quad Q_x = 1 \\ 2 * Q_x &= 1 \quad \rightarrow \quad Q_x = 0.5 \end{aligned}$$


in general:

$$m * Q_x + n * Q_y + l * Q_s = \text{integer}$$

Tune diagram up to 3rd order

Question: what will happen, if the particle performs a second turn ?

... or a third one or ... 10^{10} turns



Astronomer Hill:

*differential equation for motions with periodic focusing properties
„Hill ‘s equation“*

*Example: particle motion with
periodic coefficient*



equation of motion: $x''(s) - k(s)x(s) = 0$

*restoring force \neq const,
 $k(s)$ = depending on the position s
 $k(s+L) = k(s)$, periodic function*

*we expect a kind of quasi harmonic
oscillation: amplitude & phase will depend
on the position s in the ring.*

7.) The Beta Function

„it is convenient to see“ ... *after some beer*

... we make two statements:

1.) There exists a *mathematical function*, that defines the envelope of all particle trajectories and so can act as measure for the beam size. We call it the β – function.

2.) *Whow !!*

A particle oscillation can then be written in the form

$$x(s) = \sqrt{\varepsilon} * \sqrt{\beta(s)} * \cos(\psi(s) + \phi)$$

$\varepsilon, \Phi =$ integration *constants*
determined by initial conditions

$\beta(s)$ *periodic function* given by *focusing properties* of the lattice \leftrightarrow quadrupoles

$$\beta(s + L) = \beta(s)$$

ε *beam emittance* = *woozilycity* of the particle ensemble, *intrinsic beam parameter*,
cannot be changed by the foc. properties.

scientifically spoken: area covered in transverse x, x' phase space

... and it is constant !!!

The Beta Function

If we obtain the x, x' coordinates of a particle trajectory via



$$\begin{pmatrix} x \\ x' \end{pmatrix}_{s_2} = M(s_2, s_1) * \begin{pmatrix} x \\ x' \end{pmatrix}_{s_1}$$

The maximum size of any particle amplitude at a position “ s ” is given by

$$\hat{x}(s) = \sqrt{\varepsilon} \sqrt{\beta(s)}$$

β determines the beam size
(... the envelope of all particle trajectories at a given position “ s ” in the storage ring.

It **reflects the periodicity** of the magnet structure.



8.) Beam Emittance and Phase Space Ellipse

general solution of Hill equation

$$\left\{ \begin{array}{l} (1) \quad \mathbf{x}(s) = \sqrt{\varepsilon} \sqrt{\beta(s)} \cos(\psi(s) + \phi) \\ (2) \quad \mathbf{x}'(s) = -\frac{\sqrt{\varepsilon}}{\sqrt{\beta(s)}} \{ \alpha(s) \cos(\psi(s) + \phi) + \sin(\psi(s) + \phi) \} \end{array} \right.$$

from (1) we get

$$\cos(\psi(s) + \phi) = \frac{\mathbf{x}(s)}{\sqrt{\varepsilon} \sqrt{\beta(s)}}$$

$$\alpha(s) = \frac{-1}{2} \beta'(s)$$

$$\gamma(s) = \frac{1 + \alpha(s)^2}{\beta(s)}$$

Insert into (2) and solve for ε

$$\varepsilon = \gamma(s) \mathbf{x}^2(s) + 2\alpha(s)\mathbf{x}(s)\mathbf{x}'(s) + \beta(s) \mathbf{x}'^2(s)$$

- * ε is a *constant* of the motion ... it is independent of „s“
- * parametric representation of an *ellipse* in the $x \ x'$ space
- * shape and orientation of ellipse are given by α, β, γ

Phase Space Ellipse

particel trajectory: $x(s) = \sqrt{\varepsilon} \sqrt{\beta(s)} \cos \{ \psi(s) + \phi \}$

max. Amplitude: $\hat{x}(s) = \sqrt{\varepsilon\beta}$ \longrightarrow x' at that position ...?

... put $\hat{x}(s)$ into $\varepsilon = \gamma(s) x^2(s) + 2\alpha(s)x(s)x'(s) + \beta(s) x'^2(s)$ and solve for x'

$$\varepsilon = \gamma \cdot \varepsilon\beta + 2\alpha\sqrt{\varepsilon\beta} \cdot x' + \beta x'^2$$

\longrightarrow $x' = -\alpha \cdot \sqrt{\varepsilon / \beta}$


* A high β -function means a large beam size and a small beam divergence. !
... et vice versa !!!

* In the middle of a quadrupole $\beta = \text{maximum}$,
 $\alpha = \text{zero}$ } $x' = 0$... and the ellipse is flat

Beam Emittance and Phase Space Ellipse

In phase space x, x' a particle oscillation, observed at a given position “ s ” in the ring is running on an ellipse ... making Q revolutions per turn.


$$x(s) = \sqrt{\varepsilon} * \sqrt{\beta(s)} * \cos(\psi(s) + \varphi)$$



$$\varepsilon = \gamma(s) * x^2(s) + 2\alpha(s)x(s)x'(s) + \beta(s)x'(s)^2$$

Liouville: in reasonable storage rings area in phase space is constant.




$$A = \pi * \varepsilon = \text{const}$$



Particle Tracking in a Storage Ring


Calculate x, x' for each linear accelerator element according to matrix formalism

plot x, x' as a function of „s“





... and now the ellipse:

note for each turn x, x' at a given position „ s_1 “ and plot in the phase space diagram




Emittance of the Particle Ensemble:



... *to be very clear:*

*as long as our particle is running on an ellipse in x, x' space everything is alright, the beam is stable and **we can sleep well at nights.***


*If however we have scattering at the rest gas, or non-linear fields, or beam collisions (!) **the particle will perform a jump in x' and ϵ will increase***



Emittance of the Particle Ensemble:

$$x(s) = \sqrt{\varepsilon} \sqrt{\beta(s)} \cdot \cos(\Psi(s) + \phi)$$


$$\hat{x}(s) = \sqrt{\varepsilon} \sqrt{\beta(s)}$$



single particle trajectories, $N \approx 10^{11}$ per bunch

LHC: $\beta = 180\text{ m}$
 $\varepsilon = 5 * 10^{-10}\text{ m rad}$


$$\sigma = \sqrt{\varepsilon * \beta} = \sqrt{5 * 10^{-10}\text{ m} * 180\text{ m}} = 0.3\text{ mm}$$



**Gauß
Particle Distribution:**

$$\rho(x) = \frac{N \cdot e}{\sqrt{2\pi}\sigma_x} \cdot e^{-\frac{1}{2}\frac{x^2}{\sigma_x^2}}$$


particle at distance 1σ from centre
 $\leftrightarrow 68.3\%$ of all beam particles



aperture requirements: $r_0 = 12 * \sigma$

The „not so ideal“ World

Lattice Design in Particle Accelerators



1952: Courant, Livingston, Snyder:

Theory of strong focusing in particle beams

Recapitulation: ...the story with the matrices !!!


Equation of Motion:

$$\mathbf{x}'' + \mathbf{K} \mathbf{x} = 0 \quad K = 1/\rho^2 - k \quad \dots \text{ hor. plane:}$$


$$K = k \quad \dots \text{ vert. Plane:}$$

Solution of Trajectory Equations


$$\begin{pmatrix} \mathbf{x} \\ \mathbf{x}' \end{pmatrix}_{s_1} = \mathbf{M} * \begin{pmatrix} \mathbf{x} \\ \mathbf{x}' \end{pmatrix}_{s_0}$$



$$\mathbf{M}_{\text{drift}} = \begin{pmatrix} 1 & l \\ 0 & 1 \end{pmatrix}$$



$$\mathbf{M}_{\text{foc}} = \begin{pmatrix} \cos(\sqrt{|K|}l) & \frac{1}{\sqrt{|K|}} \sin(\sqrt{|K|}l) \\ -\sqrt{|K|} \sin(\sqrt{|K|}l) & \cos(\sqrt{|K|}l) \end{pmatrix}$$



$$\mathbf{M}_{\text{defoc}} = \begin{pmatrix} \cosh(\sqrt{|K|}l) & \frac{1}{\sqrt{|K|}} \sinh(\sqrt{|K|}l) \\ \sqrt{|K|} \sinh(\sqrt{|K|}l) & \cosh(\sqrt{|K|}l) \end{pmatrix}$$

$$\mathbf{M}_{\text{total}} = \mathbf{M}_{QF} * \mathbf{M}_D * \mathbf{M}_B * \mathbf{M}_D * \mathbf{M}_{QD} * \mathbf{M}_D * \dots$$

9.) Lattice Design: „... how to build a storage ring“

Geometry of the ring: $B^* \rho = p / e$

p = momentum of the particle,
 ρ = curvature radius

$B\rho$ = beam rigidity

Circular Orbit: bending angle of one dipole


$$\alpha = \frac{ds}{\rho} \approx \frac{dl}{\rho} = \frac{Bdl}{B\rho}$$

The angle run out in one revolution must be 2π , so for a full circle

$$\alpha = \frac{\int Bdl}{B\rho} = 2\pi$$

$$\int Bdl = 2\pi \frac{p}{q}$$

... defines the integrated dipole field around the machine.



Example LHC:




7000 GeV Proton storage ring
dipole magnets $N = 1232$
 $l = 15 \text{ m}$
 $q = +1 e$

$$\int \mathbf{B} \, dl \approx N \, l \, \mathbf{B} = 2\pi \, p / e$$

$$B \approx \frac{2\pi \, 7000 \, 10^9 \, eV}{1232 \, 15 \, m \, 3 \, 10^8 \, \frac{m}{s} \, e} = \underline{\underline{8.3 \, \text{Tesla}}}$$

LHC: Lattice Design

the ARC 90° FoDo in both planes



equipped with additional corrector coils

MB: main dipole

MQ: main quadrupole

MQT: Trim quadrupole

MQS: Skew trim quadrupole

MO: Lattice octupole (Landau damping)

MSCB: Skew sextupole

Orbit corrector dipoles


MCS: Spool piece sextupole

MCDO: Spool piece 8 / 10 pole

BPM: Beam position monitor + diagnostics

FoDo-Lattice


A magnet structure consisting of focusing and defocusing quadrupole lenses in alternating order with **nothing** in .
(**Nothing** = elements that can be neglected on first sight: drift, bending magnets, RF structures ... **and especially experiments...**)



Starting point for the calculation: in the middle of a focusing quadrupole
Phase advance per cell $\mu = 45^\circ$,
→ calculate the twiss parameters for a periodic solution


11.) The structure of matter:

Fixed target experiments:





HARP Detector, CERN

high event rate
easy track identification
asymmetric detector
limited energy reach



Collider experiments: $E=mc^2$





low event rate (luminosity)
challenging track identification
symmetric detector
 $E_{lab} = E_{cm}$

Z_0 boson discovery at the UA2 experiment (CERN).
The Z_0 boson decays
into a e^+e^- pair, shown as white dashed lines.

Problem: Our particles are *VERY* small !!

Overall cross section of the Higgs:




$$1b = 10^{-24} \text{ cm}^2$$

$$1pb = 10^{-12} * 10^{-24} \text{ cm}^2 = 1 / \text{mio} * 1 / \text{mio} * 1 / \text{mio} * 1 / \text{mio} * 1 / \text{mio} * 1 / 10000 \text{ mm}^2$$

The only chance we have:
 compress the transverse beam size ... at the IP

The particles are “very small”




LHC typical:

$$\sigma = 0.1 \text{ mm} \rightarrow 16 \mu\text{m}$$

II.) A Bit of Theory


The big storage rings: „Synchrotrons“

12.) Insertions



β -Function in a Drift


$$\beta(\ell) = \beta_0 + \frac{\ell^2}{\beta_0}$$



At the end of a long symmetric drift space the beta function reaches its maximum value in the complete lattice.

-> here we get the largest beam dimension.


-> keep l as small as possible



7 sigma beam size inside a mini beta quadrupole

... clearly there is an



*... unfortunately ... in general
high energy detectors that are
installed in that drift spaces
are a little bit bigger than a few centimeters ...*



13.) The Mini- β Insertion & Luminosity:

production rate of events is determined by the cross section Σ_{react} and a parameter L that is given by the design of the accelerator:
 ... the luminosity

$$R = L * \Sigma_{react} \approx 10^{-12} b \cdot 25 \frac{1}{10^{-15} b} = \text{some } 1000 H$$




remember:
 $1b = 10^{-24} cm^2$

The luminosity is a storage ring quality parameter and depends on beam size ($\beta !!$) and stored current

$$L = \frac{1}{4\pi e^2 f_0 b} * \frac{I_1 * I_2}{\sigma_x^* * \sigma_y^*}$$


yes ... yes ... there is NO talk without it ...

The Higgs



ATLAS event display: Higgs \Rightarrow two electrons & two muons

Luminosity



Example: Luminosity run at LHC

$$\beta_{x,y} = 0.55 \text{ m}$$

$$f_0 = 11.245 \text{ kHz}$$

$$\varepsilon_{x,y} = 5 * 10^{-10} \text{ rad m}$$

$$n_b = 2808$$

$$\sigma_{x,y} = 17 \text{ } \mu\text{m}$$

$$L = \frac{1}{4\pi e^2 f_0 n_b} * \frac{I_{p1} I_{p2}}{\sigma_x \sigma_y}$$

$$I_p = 584 \text{ mA}$$

$$L = 1.0 * 10^{34} \text{ } 1/\text{cm}^2 \text{ s}$$

The Tune ...

...is the number of these transverse oscillations per turn and corresponds to the „Eigenfrequency“ or sound of the particle oscillations. As in any oscillating system (e.g. pendulum) we have to avoid resonance conditions between the eigenfrequency of the system (= particle) and any external frequency that might act on the beam. Most prominent external frequency is the revolution frequency itself !! -> avoid integer tunes.

The Beta function shows the overall effect of all focusing fields; it has a certain value (m) that depends on the actual position in the ring.

The beam emittance describes the quality of the particle ensemble. It measures the area in phase space and can be considered like the temperature of a gas. The lower the emittance the better the beam quality. Together with the beta function it defines the beam dimension.

The lattice cell is the special magnet arrangement of the principle building block in an accelerator. Most appropriate for high energy accelerators is the FoDo.

The Higgs particle is very small, 10^{-36} cm², and so it is difficult to produce.